

# GALLATIN COUNTY EMS Patient Care Guidelines



EMS Medical Director: Bret M Birrer, MD FAAEM FACEP

Effective Date: October 20, 2025

Corrections, addition of Crashing Medical Patient guideline, addition of formulary section: December 4, 2025

These Gallatin County EMS Patient Care Guidelines were developed as a field guide for easy use by the Emergency Care Provider based on the NASEMSO National Model EMS Clinical Guidelines version 3.0 and Montana ECP Scope of Practice Document. This document is to be used by all agencies in Gallatin County under the medical direction of Bret M Birrer, MD FAAEM FACEP including:

American Medical Response  
Amsterdam Fire Department  
Big Sky Fire Department  
Bozeman Fire Department  
Bridger Canyon Fire Department  
Central Valley Fire District  
Clarkston Fire Service Area  
Fort Ellis Fire/Rescue  
Gallatin County 911  
Gallatin County Sheriff Office – deputy EMT program  
Gallatin County Sheriff Search and Rescue  
Gallatin Gateway Rural Fire District  
Gallatin River Ranch Fire Department  
Hebgen Basin Rural Fire District  
Hyalite Fire District  
Manhattan Fire Department  
Montage Big Sky  
Mountain Western Medical, LLC  
Three Forks Area Ambulance  
Willow Creek Fire District  
Yellowstone Club Ski Patrol  
Yellowstone Mountain Club Rural Fire District

### **ACKNOWLEDGEMENTS**

Global Medical Response for an edible version of the Clark County EMS System Emergency Medical Care Protocols to use as a base for creation of this document. Dr. Josh Gaither, Chief of Division of EMS at The University of Arizona College of Medicine – Tucson, for permission to use the Shock or Crashing Medical Patient Administrative Guideline. Dr. Eric Lowe, former EMS Medical Director for Gallatin County, for his original work on most of the Operations Guidelines adopted from the previous version of the Gallatin County EMS Guidelines.

## FOREWORD

These Patient Care Guidelines have been adapted from the National Association of State EMS Officials (NASEMSO) Model EMS Clinical Guidelines published online in March 2022. These algorithms include specific recommendations for evaluation and treatment.

The recommendations within each guideline are listed in order by provider level scope of practice. It is assumed that more advanced levels of ECPs will perform all recommended evaluations and treatments included in the preceding level of care.

The guidelines include specific pediatric recommendations, highlighted by the EMS for Children bear logo, where specific pediatric recommendations differ from those for adults. It is assumed that children will receive the evaluation and care recommended for all patients, unless specific pediatric recommendations are included in the algorithm.

A pediatric patient is defined as age less than 15 years. Pediatric treatment guidelines are to be used on children who have not yet experienced puberty. Signs of puberty include chest or underarm hair on males, and any breast development in females. Age 15 and above, or signs of puberty, is considered an adult patient in regard to treatment guidelines.

The General Adult Assessment and General Pediatric Assessment guidelines should be applied to all patient encounters. All initial patient care is included in this guideline to reduce the need for extensive reiteration of basic assessment and other considerations in every guideline.

Online medical direction may be utilized at any time during the patient encounter per local protocols.

The NASEMSO model guidelines include additional information that medical direction authorities may find helpful for education, training, and quality improvement activities, including patient safety considerations, educational pearls, performance measures, and literature references: [nasemso.org](http://nasemso.org).

### General EMS Medical Director Expectations

Paramedics are expected to maintain National Registry.

All paramedics new to Gallatin County (newly licensed as well as experienced) will need to successfully complete an EMS Medical Director clearance meeting between completing their agency specific FTO time and being allowed to practice paramedicine independently in Gallatin County.

Failure to abide by the following cardinal rules will result in de-credentialling and inability to practice prehospital medicine in Gallatin County:

1. Don't intentionally harm a patient.
2. Don't intentionally withhold needed treatment.
3. Don't falsify the medical record.
4. Don't practice while intoxicated.
5. Failure to remediate.

Bret M Birrer, MD FAAEM FACEP  
EMS Medical Director  
Gallatin County

## TABLE OF CONTENTS

Effective Date / Agencies / Acknowledgements .....	2
Foreword / Medical Director Expectations .....	3
Terms and Conventions .....	8
<b>ADULT TREATMENT GUIDELINES .....</b>	<b>10</b>
General Adult Assessment .....	11
General Adult Trauma Assessment .....	12
Abdominal/Flank Pain, Nausea & Vomiting .....	14
Allergic Reaction .....	16
Altitude Illness .....	18
Altered Mental Status/Syncope .....	20
Behavioral Emergency .....	22
Bradycardia .....	24
Burns .....	26
Cardiac Arrest (Non-Traumatic) .....	29
High Performance CPR .....	32
Chest Pain (Non-Traumatic) and Suspected Acute Coronary Syndrome .....	33
Childbirth/Labor .....	35
Cold-Related Illness .....	37
Crashing Medical Patient .....	39
Drowning .....	41
Epistaxis .....	43
Heat-Related Illness .....	45
Head Injury (EPIC-TBI) .....	47
Hyperkalemia (Suspected) .....	48
Obstetrical Emergency .....	50
Overdose/Poisoning .....	52
Pain Management .....	54
Pulmonary Edema/CHF .....	56
Respiratory Distress .....	58
Seizure .....	60
Sepsis (Suspected) .....	62
Shock .....	64
Smoke Inhalation .....	66
STEMI (Suspected) .....	68
Stroke (CVA) .....	70
Tachycardia/Stable .....	73
Tachycardia/Unstable .....	75
Ventilation Management .....	77
<b>PEDIATRIC TREATMENT GUIDELINES .....</b>	<b>79</b>
General Pediatric Assessment .....	80
General Pediatric Trauma Assessment .....	82
Abdominal Pain, Nausea & Vomiting .....	84
Allergic Reaction .....	86
Altered Mental Status .....	88
Bradycardia .....	90
Brief Resolved Unexplained Event (BRUE) .....	92
Burns .....	94
Cardiac Arrest (Non-Traumatic) .....	97
Cold-Related Illness .....	99



## **PEDIATRIC TREATMENT GUIDELINES (Cont.)**

Drowning .....	101
Heat-Related Illness .....	103
Head Injury (EPIC-TBI) .....	105
Neonatal Resuscitation .....	106
Overdose/Poisoning .....	108
Pain Management .....	110
Respiratory Distress .....	112
Seizure .....	114
Shock .....	116
Smoke Inhalation .....	118
Tachycardia/Stable .....	120
Tachycardia/Unstable .....	122
Ventilation Management .....	124

## **OPERATIONS GUIDELINES .....126**

Air Medical Activation .....	127
BHDRAMC Trauma Activation .....	128
Big Sky Area Trauma Field Triage .....	129
BSMC Trauma Activation .....	130
Defining a Patient .....	131
Destination Hospital Choice .....	132
Interfacility Transfers .....	133
Medication List .....	134
Prehospital Death Determination .....	135
Prehospital Report .....	136
POLST/Palliative Care .....	137
Refusal .....	139
Termination of Resuscitation .....	141

## **PROCEDURES GUIDELINES .....142**

Continuous Positive Airway Pressure (CPAP) .....	143
Electrical Therapy/Defibrillation .....	144
Electrical Therapy/Synchronized Cardioversion .....	145
Electrical Therapy/Transcutaneous Pacing .....	146
Endotracheal Intubation .....	147
Hemorrhage Control .....	148
High Flow Nasal Cannula (HFNC) .....	149
Legal Blood Draw .....	151
LUCAS Device .....	152
Medication Administration .....	153
Needle Cricothyroidotomy .....	154
Needle Decompression .....	155
Push Dose Epinephrine .....	156
Spinal Motion Restriction .....	157
Supraglottic Airway Device .....	158
Surgical Cricothyroidotomy .....	159
Tracheostomy Tube Replacement .....	160
Traction Splint .....	161
Ultrasound .....	162
Vagal Maneuvers .....	165
Vascular Access .....	166

<b>FORMULARY .....</b>	<b>167</b>
Acetaminophen .....	168
Activated charcoal .....	168
Adenosine .....	168
Albuterol .....	169
Amiodarone .....	169
Aspirin .....	170
Atropine .....	170
Brilinta .....	171
Calcium chloride .....	171
Calcium gluconate .....	171
Dextrose .....	171
Diazepam .....	172
Diphenhydramine .....	172
Droperidol .....	173
Epinephrine .....	174
Fentanyl .....	175
Glucagon .....	175
Haloperidol .....	176
Heparin .....	176
Hydroxocobalamin .....	176
Ibuprofen .....	177
Ipratropium .....	177
Ketamine .....	178
Ketorolac .....	178
Lidocaine .....	179
Lorazepam .....	179
Magnesium sulfate .....	180
Methylprednisolone .....	180
Metoclopramide .....	181
Metoprolol .....	181
Midazolam .....	182
Morphine .....	182
Naloxone .....	183
Nitroglycerin .....	183
Norepinephrine .....	183
Ondansetron .....	184
Oral glucose .....	184
Oxygen .....	185
Oxymetazoline .....	185
Oxytocin .....	185
Sodium bicarbonate .....	186
Tranexamic acid .....	186

Patient Care Guideline Key:



Caution / Warning / Alert



Pediatric Treatment Consideration (for patients  $\leq 15$  years of age)



Online Medical Control



Specific Guideline



EMR Licensed Attendant and above may perform these steps



EMT Licensed Attendant and above may perform these steps



EMT with Endorsement(s) [airway, IV/IO initiation, IV/IO maintenance, medications] and above may perform these steps



AEMT Licensed Attendant and above may perform these steps



Paramedic Licensed Attendant

## TERMS AND CONVENTIONS


<b>AED</b>	means Automated External Defibrillator
<b>AMPLE</b>	means Allergies; Medications; Prior history; Last meal eaten; Events leading up to injury/illness
<b>AMS</b>	means Altered Mental Status
<b>ASA</b>	means Acetylsalicylic Acid
<b>BG</b>	means Blood Glucose
<b>BP</b>	means Blood Pressure
<b>BVM</b>	means Bag-Valve-Mask
<b>CCC</b>	means Continuous Cardiac Compressions
<b>CHF</b>	means Congestive Heart Failure
<b>COPD</b>	means Chronic Obstructive Pulmonary Disease
<b>CP</b>	means Chest Pain
<b>CPR</b>	means Cardiopulmonary Resuscitation
<b>CVA</b>	means Cerebrovascular Accident
<b>DCAP-BTLS</b>	means Deformities; Contusions; Abrasions; Punctures/Penetrations; Burns; Tenderness; Lacerations; Swelling
<b>DKA</b>	means Diabetic Ketoacidosis
<b>ECG</b>	means Electrocardiogram
<b>ETA</b>	means Estimated Time of Arrival
<b>ETT</b>	means Endotracheal Tube
<b>GCS</b>	means Glasgow Coma Scale
<b>GU</b>	means Genitourinary
<b>HACE</b>	means High Altitude Cerebral Edema
<b>HAPE</b>	means High Altitude Pulmonary Edema
<b>HEENT</b>	means Head, Ears, Eyes, Nose, Throat
<b>HPI</b>	means History of Present Illness
<b>HR</b>	means Heart Rate
<b>ICP</b>	means Intracranial Pressure
<b>IM</b>	means Intramuscular
<b>IN</b>	means Intranasal
<b>IO</b>	means Intraosseous
<b>IV</b>	means Intravenous
<b>IVP</b>	means Intravenous Push




<b>IVPB</b>	means Intravenous Piggyback
<b>JVD</b>	means Jugular Venous Distention
<b>MAD</b>	means Mucosal Atomizer Device
<b>MI</b>	means Myocardial Infarction
<b>MOI</b>	means Mechanism of Injury
<b>NRB</b>	means Non-rebreather
<b>NS</b>	means Normal Saline
<b>NV</b>	means Nausea/Vomiting
<b>OPQRST</b>	means Onset; Provokes; Quality; Radiates; Severity; Time (used in evaluating localized pain)
<b>PCI</b>	means Percutaneous Coronary Intervention
<b>PCR</b>	means Patient Care Record/Report
<b>PO</b>	means By Mouth
<b>PRN</b>	means As Needed
<b>q</b>	means Every
<b>ROSC</b>	means Return of Spontaneous Circulation
<b>RR</b>	means Respiratory Rate
<b>RUQ</b>	means Right Upper Quadrant
<b>SAMPLE</b>	means Symptoms; Allergies; Medications; Prior history; Last meal eaten; Events leading up to injury/illness
<b>SL</b>	means Sublingual
<b>SOB</b>	means Shortness of Breath
<b>S/P</b>	means Status/Post
<b>SQ</b>	means Subcutaneous
<b>S/S</b>	means Signs/Symptoms
<b>SVN</b>	means Small Volume Nebulizer
<b>SVT</b>	means Supraventricular Tachycardia
<b>TCAs</b>	means Tricyclic Antidepressants
<b>TFTC</b>	means Trauma Field Triage Criteria
<b>TIA</b>	means Transient Ischemic Attack
<b>TKO/KVO</b>	means To Keep Open/Keep Vein Open
<b>VF</b>	means Ventricular Fibrillation
<b>VT</b>	means Ventricular Tachycardia
<b>VS</b>	means Vital Signs
<b>WPW</b>	means Wolff-Parkinson-White Syndrome

# **ADULT TREATMENT GUIDELINES**

# General Adult Assessment



# General Adult Trauma Assessment





**History**

- Time and mechanism of injury
- Damage to structure or vehicle
- Location in structure or vehicle
- Others injured or dead
- Speed and details of MVC
- Restraints/protective equipment
- Past medical history
- Medications

**Signs and Symptoms**

- DCAP-BTLS
- AMS or unconscious
- Hypotension or shock
- Arrest


**Differential (life threatening)**

- Tension pneumothorax
- Flail chest
- Pericardial tamponade
- Open chest wound
- Hemothorax
- Intra-abdominal bleeding
- Pelvis/femur fracture
- Spine fracture/cord injury
- Head injury
- Extremity fracture
- HEENT (airway obstruction)
- Hypothermia

**Pearls**

- Recommended exam: Mental Status, Skin, HEENT, Heart, Lung, Abdomen, Extremities, Back, Neuro.
- Transport destination is based on the Trauma Field Triage Criteria Guideline.
- Transport should not be delayed for procedures; ideally procedures should be performed enroute when possible. Target scene time less than 10 minutes for unstable patients or those likely to need surgical intervention.
- BVM is an acceptable method of ventilating and managing an airway if pulse oximetry can be maintained  $\geq 90\%$ . Consider airway adjuncts as appropriate avoiding nasal airway adjuncts in patients with oral or other facial injuries.
- Head injury: target SBP greater than 110 mmHg. Hypotension should be avoided to maintain cerebral perfusion.
- TRANEXAMIC ACID (TXA) administration may be considered within three hours of injury and signs of hemorrhagic shock.
- If pelvis is unstable, place pelvic binder or sheet to stabilize pelvis .
- Geriatric patients should be evaluated with a high index of suspicion; occult injuries may be present and geriatric patients can decompensate quickly.

# Abdominal / Flank Pain, Nausea & Vomiting



**History**

- Age
- Medical/surgical history
- Onset
- Quality
- Severity
- Fever
- Menstrual history

**Signs and Symptoms**

- Pain location
- Tenderness
- Nausea
- Vomiting
- Diarrhea
- Dysuria
- Constipation
- Vaginal bleeding/discharge
- Pregnancy

**Differential**

- Liver (Hepatitis)
- Gastritis, PUD
- Gallbladder
- MI
- Pancreatitis
- Kidney stone
- Abdominal aneurysm
- Appendicitis
- Bladder/prostate disorder
- Pelvic (PID, ectopic pregnancy, ovarian cyst)
- Spleen enlargement
- Bowel obstruction
- Gastroenteritis
- Ovarian and testicular torsion

**Pearls**

- Recommended Exam: Mental Status, Skin, HEENT, Heart, Lung, Abdomen, Back, Extremities, Neuro.
- Neuro disorders or signs of hypoperfusion/shock in the presence of abdominal pain may indicate an aneurysm.
- Document mental status and vital signs prior to administration of antiemetics & pain management.
- Repeat vital signs after each fluid bolus
- In patients  $\geq 35$  years old consider cardiac origin. Perform a 12-Lead ECG.
- Consider retroperitoneal palpation for kidney pain.
- Abdominal pain in women of childbearing age should be considered pregnancy until proven otherwise.
- Nausea and vomiting are symptoms of illness – in addition to treating the patient's nausea and vomiting a thorough history and physical are key to identifying what may be a disease in need of emergent treatment (e.g., bowel obstruction, myocardial infarction, pregnancy).
- For nausea/vomiting during pregnancy, preference is METOCLOPRAMIDE and/or DIPHENHYDRAMINE (both FDA pregnancy category B).
- Abdominal pain in older adults, patients with bleeding disorders, patients on anticoagulation medications, and patients that are immunocompromised may be a harbinger for severe illness.

**Dystonic Reaction / Extrapyrimal Symptoms**

- Condition causing involuntary muscle movements or spasms typically of the face, neck and upper extremities.
- Typically an adverse reaction to drugs such as METOCLOPRAMIDE or DROPERIDOL.
- When recognized, administer DIPHENHYDRAMINE 25-50 mg IM/IV.

# Allergic Reaction



General Adult Assessment

Evidence of anaphylaxis?

No (Mild)

Yes (Moderate/Severe)

E\*



Vascular Access

E\*

A

DIPHENHYDRAMINE  
25-50 mg PO

P

DIPHENHYDRAMINE  
25-50 mg IM/IV

Reassess patient q 5 min

Allergic Reaction – mild immune response to an allergen with symptoms such as hives or local swelling/itching.

Anaphylaxis – moderate/severe reaction meeting either of these criteria:

- Exposure to known or likely allergen with hypotension OR respiratory compromise
- Two or more of the following after exposure to likely allergen: skin/mucosal changes (hives, flushing, edema); respiratory compromise (SOB, wheezing, stridor, hypoxia); hypotension or signs of shock; persistent GI symptoms (abdominal pain, vomiting)



Refer to Push Dose  
Epinephrine Guideline pg 156  
for mixing instructions



Continue General Adult Assessment

EPINEPHRINE

FR

Patient prescribed EpiPen

E

EPINEPHRINE auto-injector

E\*

EPINEPHRINE 1:1000, 0.3 - 0.5 mg  
IM; may repeat q 15 min

BREATHING TREATMENT

E

ALBUTEROL 2.5 mg in 3 ml SVN,  
repeat as needed  
+/-  
IPRATROPIUM 500 mcg SVN  
once

E\*

A

DIPHENHYDRAMINE  
50 mg PO

E\*



Vascular access  
500 ml LR or NS bolus IV/IO;  
may repeat up to 2000 ml

P

DIPHENHYDRAMINE  
50 mg IM/IV

P



Ventilation Management

Cardiac monitor

Pt in persistent shock? If Yes, then

P

PUSH DOSE EPINEPHRINE  
1:100,000 10 mcg IV/IO, may  
repeat q 2-5 min to maintain  
SBP>90 (1 ml of 1:100,000  
solution) AND/OR EPINEPHRINE  
0.5 mcg/kg/min IV drip



### History

- Onset and location
- Insect sting or bite
- Food allergy/exposure
- Medication allergy/exposure
- New clothing, soap, detergent
- Past history of reactions
- Past medical history
- Medication history

### Signs and Symptoms

- Itching or hives
- Coughing/wheezing or respiratory distress
- Throat or chest constriction
- Difficulty swallowing
- Hypotension/shock
- Edema
- Nausea/vomiting

### Differential


- Urticarial (rash only)
- Anaphylaxis (systemic effect)
- Shock (vascular effect)
- Angioedema (drug induced)
- Aspiration/airway obstruction
- Asthma/COPD
- CHF

### Pearls


- Recommended Exam: Mental Status, Skin, Heart, Lung.
- Anaphylaxis is an acute and potentially lethal multisystem allergic reaction.
- EPINEPHRINE is the first-line drug that should be administered in acute anaphylaxis (moderate / severe symptoms).
- IM EPINEPHRINE (1:1,000) should be administered in priority before or during attempts at IV or IO access.
- Never give EPINEPHRINE 1:1000 (IM concentration) through IV/IO route.
- IM EPINEPHRINE should be administered in the anterolateral thigh (vastus lateralis) as produces the most rapid rise in serum level.
- There are no absolute contraindications to EPINEPHRINE. Use with caution in elderly patients, patients with known cardiovascular disease, or significant tachycardia or hypertension.
- Contact Medical Control for refractory anaphylaxis.
- Consider ETCO2 monitoring.
- Hypovolemia or distributive shock should be addressed with a fluid bolus prior to the administration of push-dose pressors.
- Remove trigger if still present (sting, food, etc)
- Always perform ECG monitoring when administering EPINEPHRINE.
- Contrary to common belief that all cases of anaphylaxis present with cutaneous manifestations, such as urticaria or mucocutaneous swelling, a significant portion of anaphylactic episodes may not involve these signs and symptoms on initial presentation.
- There is no proven benefit to using steroids in the management of allergic reactions and/or anaphylaxis

### QI Metrics:

- EPINEPHRINE given appropriately.
- Airway assessment documented.



# Altitude Illness



**History**

- Altitude of home residence
- Duration of ascent to elevation
- Prophylaxis against altitude (ibuprofen, acetazolamide, sildenafil)
- Past medical history

**Signs and Symptoms**

- Headache
- Anorexia
- Nausea, vomiting
- Fatigue, weakness
- Dizziness, lightheadedness
- Difficulty sleeping
- Dyspnea, cough
- Altered mentation
- Ataxia
- Stupor, coma

**Differential**

- Carbon monoxide poisoning
- Dehydration
- Exhaustion
- Hypoglycemia
- Hyponatremia
- Tension headache, migraine headache
- CHF
- Flash pulmonary edema
- Metabolic encephalopathy

**Pearls**

- Recommended Exam: Mental Status, Heart, Lung, Neuro.
- Descent is the mainstay of therapy and is the definitive therapy for all altitude related illnesses.
- Descent should always be the primary treatment strategy for patients suffering from altitude illness, especially patients suffering from HACE and HAPE.
- Descent of 500–1000 feet is often enough to see improvements in patient conditions.
- The high-altitude environment is inherently dangerous. Rescuers must balance patient needs with patient safety and safety for the responders.
- Rapid descent by a minimum of 500–1000 feet is a priority, however rapidity of descent must be balanced by current environmental conditions and other safety considerations.
- Patients with HAPE are suffering from non-cardiogenic pulmonary edema and may benefit from positive pressure ventilation via either bag assisted ventilation, CPAP, or other means of positive pressure ventilation.
- Patients suffering from altitude illness are commonly dehydrated and require IV fluids — once resuscitation is complete and the patient requires no further fluid boluses, maintain IV fluids at 125 mL/hr.
- HAPE is the most lethal of all altitude illnesses.

**Acute mountain sickness:**

- Headache plus one or more of the following: anorexia, nausea or vomiting, fatigue or weakness, dizziness or lightheadedness or difficulty sleeping.
- In infants and young children, symptoms include pallor, fussiness, vomiting, decreased appetite, poor sleep, decreased playfulness.
- These symptoms must occur in the setting of recent arrival to high altitude (generally considered greater than 5000 – 7000 feet).


**High altitude pulmonary edema (HAPE):**

- Progressive dyspnea, cough, hypoxia, and weakness in high altitude environments (considered greater than 8000 feet).
- In infants and young children, symptoms again include pallor, fussiness, vomiting, decreased appetite, poor sleep, decreased playfulness.
- Patients may or may not exhibit new symptoms if acute mountain sickness precedes symptoms of HAPE.

**High altitude cerebral edema (HACE):**

- Heralded by mental status changes in patients with symptoms of acute mountain sickness including altered mentation, ataxia, or stupor and progressing to coma.
- Typically seen in high altitude environments (greater than 8000 feet).

# Altered Mental Status / Syncope





### History

- Known diabetic, Medic Alert tag
- Drugs or drug paraphernalia
- Report of drug use or toxic ingestion
- Past medical history
- Medications
- History of trauma
- Change in condition
- Changes in feeding or sleep habits

### Signs and Symptoms

- Decreased mental status or lethargy
- Changes in baseline mental status
- Bizarre behavior
- Hypoglycemia
- Hyperglycemia
- Irritability
- Breath odor
- Track marks
- Evidence of trauma

### Differential

- Head trauma
- CNS (stroke, tumor, seizure, infection)
- Cardiac (MI, CHF)
- Hypothermia
- Infection
- Thyroid
- Shock (septic, metabolic, traumatic)
- Diabetes
- Toxicological or ingestion
- Acidosis/Alkalosis
- Environmental exposure
- Hypoxia
- Electrolyte abnormality
- Psychiatric disorder

### Pearls

- Recommended Exam: Mental Status, HEENT, Skin, Heart, Lung, Abdomen, Back Extremities, Neuro.
- Preference is to administer D10. May administer D50 if D10 is not available. DEXTROSE 50% can cause local tissue damage if it extravasates from vein and may cause hyperglycemia.
- Hypoglycemic patients who have had a seizure should be transported to the hospital regardless of their mental status and response to therapy.
- Sulfonylureas (e.g., glyburide, glipizide) have long half-lives ranging from 12–60 hrs. Patients with corrected hypoglycemia who are taking these agents are at particular risk for recurrent symptoms and frequently require hospital admission.
- Pay careful attention to the head exam for signs of injury.
- With depressed mental status, initial focus is on airway protection, oxygenation, ventilation, and perfusion.
- Be aware of AMS as presenting sign of an environmental toxin or Haz-Mat exposure, and protect personal safety and that of other responders.
- Do not let alcohol confuse the clinical picture; alcohol is not commonly a cause of total unresponsiveness to pain.
- If narcotic overdose or hypoglycemia is suspected, administer NARCAN 0.4-2mg or GLUCOSE prior to advanced airway procedures.

### AMS differential – mnemonic **AEIOU-TIPS**:

**A** – Alcohol, Abuse, Atypical migraine

**E** – Epilepsy, Electrolytes

**I** – Insulin (hypoglycemia)

**O** – Oxygen, Overdose

**U** – Uremia (kidney failure)


**T** – Trauma, Tumor

**I** – Infection

**P** – Psych, Poisoning

**S** – Seizure, Subarachnoid hemorrhage, Sepsis

# Behavioral Emergency



### History

- Situational crisis
- Psychiatric illness/medications
- Injury to self or threats to others
- Medic Alert tag
- Substance abuse/overdose
- Diabetes
- Seizure history

### Signs and Symptoms

- Anxiety, agitation, confusion
- Affect change, hallucinations
- Delusional thoughts, bizarre behavior
- Combative, violent
- Expression of suicidal/homicidal thoughts

### Differential

- AMS differential
- Hypoglycemia
- Postictal state
- Alcohol intoxication
- Toxin/substance abuse
- Medication effect or overdose
- Withdrawal syndromes
- Depression
- Bipolar
- Schizophrenia
- Anxiety disorder

### Pearls

- **Pharmacological sedation is a medical procedure that results from a medical assessment. Sedation is never to be utilized to control behavior for the purpose of law enforcement initiatives or assistance.**
- **Law enforcement assistance should be requested on all calls involving potentially violent patients.**
- **Under no circumstances are patients to be transported restrained in the prone position.**
- **Patients may not be transported with their arms restrained behind their back or in an ankle-to-wrist (hog-tied) manner.**
- The clinician should be ready to resuscitate the patient in case of inadvertent changes in respiratory or hemodynamic status. Patients should be continuously monitored with all available adjuncts when possible, including HR, ECG, RR, SpO2, BP, ETCO2, perfusion state, mental state.
- Physical restraints, including gurney straps, should never restrict chest wall movement.
- Law enforcement **MUST** accompany the patient in the back of the ambulance if in handcuffs and/or The Wrap.
- Patients expressing suicidal or homicidal ideation or who are otherwise a danger to themselves or others may not refuse transport. Contact law enforcement if necessary to initiate legal hold.
- EMS providers are not to remove taser darts unless there is a need to do so to administer medical care. Dart removal is part of the education to use the device and is the responsibility of the person or agency who deploys the device.


### Dystonic Reaction

- Condition causing involuntary muscle movements or spasms typically of the face, neck and upper extremities.
- Typically an adverse reaction to drugs such as HALOPERIDOL (may occur with administration).
- When recognized, administer DIPHENYDRAMINE 50 mg IM/IV/IO.

### S.A.F.E.R.

- Stabilize the situation by containing and lowering the stimuli.
- Assess and acknowledge the crisis.
- Facilitate the identification and activation of resources (chaplain, family, friends or police).
- Encourage patient to use resources and take actions in his/her best interest.
- Recovery or referral – leave patient in care of responsible person or professional, or transport to appropriate facility.

# Bradycardia



**History**

- Past medical history
- Medications (AV nodal blockers)
- Pacemaker

**Signs and Symptoms**

- HR <60/min with hypotension, acute AMS, chest pain, acute CHF, seizures, syncope, or shock secondary to bradycardia
- Respiratory distress

**Differential**

- Acute myocardial infarction
- Hypoxia
- Pacemaker failure
- Hypothermia
- Sinus bradycardia
- Athletic
- Head injury (elevated ICP) or stroke
- Spinal cord lesion
- Sick sinus syndrome
- AV block
- Overdose / Toxin exposure
- Shock
- Electrolyte disorder


**Pearls**

- Recommended Exam: Mental Status, HEENT, Heart, Lung, Neuro.
- Identifying signs and symptoms of poor perfusion caused by bradycardia are paramount. Treatment should only be given when patient is symptomatic due to bradycardia. Athletes, patients on beta blockers, and young healthy patients may have a slow resting heart rate at baseline.
- Do not delay pacing while waiting for IV access.
- Ensure adequate oxygenation and provide ventilatory support as needed.
- Consider hyperkalemia in patients with wide complex bradycardia.
- Atropine is unlikely to be effective and should be avoided in patients who have had a heart transplant.

**QI Metrics**

- High degree blocks correctly identified.
- Pacer pads on patient if ATROPINE given.
- Patient paced if appropriate.

# Burns



### History

- Type of exposure (heat, gas, chemical)
- Inhalational injury
- Time of injury
- Past medical history & medications
- Other trauma
- Loss of consciousness
- Tetanus/immunization status

### Signs and Symptoms

- Burns, pain, swelling
- Dizziness
- Loss of consciousness
- Hypotension/shock
- Airway compromise/distress
- Wheezing
- Singed facial or nasal hair
- Hoarseness or voice changes

### Differential

- Superficial (1<sup>st</sup> degree) – red and painful
- Partial Thickness (2<sup>nd</sup> degree) – blistering
- Full Thickness (3<sup>rd</sup> degree) – painless/charred or leathery skin
- Thermal
- Chemical
- Electrical
- Radiation
- Lightning

### Pearls

- Burn patients are trauma patients; evaluate for multisystem trauma.
- Assure whatever has caused the burn, is no longer contacting the injury. (Stop the burning process!)
- Recommended Exam: Mental Status, HEENT, Neck, Heart, Lungs, Abdomen, Extremities, Back, Neuro.
- Consider early intubation with patients experiencing significant inhalation injuries. Consider air ambulance for airway management needs beyond the scope of the responding ground medic.
- Potential CO exposure should be treated with 100% oxygen. Pulse oximetry may not be accurate. (For patients in which the primary event is CO inhalation, transport to a hospital equipped with a hyperbaric chamber is indicated [when reasonably accessible].)
- Circumferential burns to extremities are dangerous due to potential vascular compromise secondary to soft tissue swelling. Elevate extremity.
- Burn patients are prone to hypothermia - never apply ice or cool burns; must maintain normal body temperature.
- Consider ETCO<sub>2</sub> monitoring.
- Have a high index of suspicion for cyanide poisoning in a patient with depressed GCS, respiratory difficulty, and cardiovascular collapse in the setting of an enclosed-space fire. Give antidote (hydroxocobalamin), if available, in this circumstance

### Early Intubation Indications

- Signs of Airway Obstruction
- Hoarseness, Stridor, Dysphagia
- Extensive Deep Facial Burns
- Significant Risk of Edema
- Burns in Mouth
- Total BSA  $\geq 40\%$
- Signs of Respiratory Compromise
  - Accessory Muscle Use
  - Inability to Clear Secretions
  - Poor Oxygenation
- Altered Mentation

### Patients meeting the following criteria shall be transported to the closest appropriate Burn Care Center:


1. Second degree burns  $>10\%$  body surface area (BSA).
2. Any Third degree burns.
3. Burns that involve the face, hands, feet, genitalia, perineum, or major joints.
4. Electrical burns including lightning injury.
5. Chemical burns.
6. Circumferential burns.
7. Inhalation burns.
8. Burn injury with concomitant trauma

### Pearls (Electrical)

- Do not contact the patient until you are certain the source of the electric shock has been disconnected.
- Attempt to locate contact points, (entry wound where the AC source contacted the patient; an exit at the ground point); both sites will generally be full thickness.
- Cardiac monitor; anticipate ventricular or atrial irregularity to include V-Tach, V-Fib, heart blocks, etc.
- Attempt to identify the nature of the electrical source (AC vs DC), the amount of voltage and the amperage the patient may have been exposed to during the electrical shock.

### Pearls (Chemical)

- Certainly 0.9% NaCl Sol'n or Sterile Water is preferred; however if it is not readily available, do not delay; use tap water for flushing the affected area or other immediate water sources. Flush the area as soon as possible with the cleanest, readily available water or saline solution using copious amounts of fluids.






\*Fluid of choice LR/NS, DO NOT use dextrose containing fluids

Wt (lbs)	Wt (kg)	% TBSA	/Hr for 1 <sup>st</sup> 8 Hrs of care	60 gtt set, gtt/min	20 gtt set, gtt/min	15 gtt set, gtt/min	10 gtt set, gtt/min
66	30	10	75	75	25.0	18.8	12.5
66	30	20	150	150	50.0	37.5	25.0
66	30	30	225	225	75.0	56.3	37.5
66	30	40	300	300	100.0	75.0	50.0
66	30	50	375	375	125.0	93.8	62.5
66	30	60	450	450	150.0	112.6	75.0
88	40	10	100	100	33.3	25.0	16.7
88	40	20	200	200	66.7	50.0	33.3
88	40	30	300	300	100.0	75.0	50.0
88	40	40	400	400	133.3	100.0	66.7
88	40	50	500	500	166.7	125.00	83.3
88	40	60	600	600	200.0	150.0	100.0
110	50	10	125	125	41.7	31.3	20.8
110	50	20	250	250	83.3	62.5	41.7
110	50	30	375	375	125.0	93.8	62.5
110	50	40	500	500	166.7	125.0	83.3
110	50	50	625	625	208.3	156.3	104.2
110	50	60	750	750	250.0	187.6	125.0
132	60	10	150	150	50.0	37.5	25.0
132	60	20	300	300	100.0	75.0	50.0
132	60	30	450	450	150.0	112.5	75.0
132	60	40	600	600	200.0	150.0	100.0
132	60	50	750	750	250.0	187.5	125.0
132	60	60	900	900	300.0	225.0	150.0
154	70	10	175	175	58.3	43.8	29.2
154	70	20	350	350	116.7	87.5	58.3
154	70	30	525	525	175.0	131.3	87.5
154	70	40	700	700	233.3	175.0	116.7
154	70	50	875	875	291.7	218.8	145.8
154	70	60	1050	1050	350.0	262.6	175.0
176	80	10	200	200	66.7	50.0	33.3
176	80	20	400	400	133.3	100.0	66.7
176	80	30	600	600	200.0	150.0	100.0
176	80	40	800	800	266.7	200.0	133.3
176	80	50	1000	1000	333.3	250.0	166.7
176	80	60	1200	1200	400.0	300.0	200.0
198	90	10	225	225	75.0	56.3	37.5
198	90	20	450	450	150.0	112.5	75.0
198	90	30	675	675	225.0	168.8	112.5
198	90	40	900	900	300.0	225.0	150.0
198	90	50	1125	1125	375.0	281.3	187.5
198	90	60	1350	1350	450.0	337.6	225.0
220	100	10	250	250	83.3	62.5	41.7
220	100	20	500	500	166.7	125.0	83.3
220	100	30	750	750	250.0	187.5	125.0
220	100	40	1000	1000	333.3	250.0	166.7
220	100	50	1250	1250	416.7	312.5	208.3
220	100	60	1500	1500	500.0	375.0	250.0
242	110	10	275	275	91.6	68.7	45.9
242	110	20	550	550	183.4	137.5	91.6
242	110	30	825	825	275	206.2	137.5
242	110	40	1100	1100	366.6	275.0	183.4
242	110	50	1375	1375	458.4	343.7	229.1
242	110	60	1650	1650	550.0	412.4	275
264	120	10	300	300	99.9	74.9	50.1
264	120	20	600	600	200.1	150.0	99.9
264	120	30	825	825	300.0	224.9	150.0
264	120	40	1200	1200	399.9	300.0	200.1
264	120	50	1500	1500	500.1	374.9	249.9
264	120	60	1650	1650	600.0	449.8	300



# Cardiac Arrest (Non-Traumatic)



#### History

- Events leading to arrest
- Estimated down time
- Past medical history
- Medications
- Existence of terminal illness

#### Signs and Symptoms

- Unresponsive
- Apneic
- Pulseless

#### Differential

- Medical vs. Trauma
- VF vs. Pulseless VT
- Asystole
- PEA
- Primary cardiac event vs. respiratory or drug overdose

#### Pearls

- Efforts should be directed at high quality and continuous compressions with limited interruptions and early defibrillation when indicated. Crews should consider using a “pit crew” approach with predefined roles and crew resource management principles. Strongly consider initial anterior-posterior pad placement due to superior rates of successful defibrillation.
- Consider early IO placement if IV is difficult.
- Ventilation rate should be 8-10 breaths per minute. Hyperventilation can worsen patient outcomes.
- Continuous waveform capnography should be monitored throughout resuscitation for confirmation and monitoring of advanced airways (when present), as well as monitoring effectiveness of chest compressions.
- Routine use of mechanical CPR devices is not recommended. If utilized, the mechanical CPR device should be applied in a manner that minimizes interruptions in compressions, keeping breaks in CPR to less than 10 seconds. Use mechanical CPR devices per manufacturer’s guidelines. Manual CPR must be initiated before the application of a CPR device. Ideally, complete 2 rounds of manual compressions before application.
- **Do not give further doses of EPINEPHRINE beyond 3 mg in cardiac arrest.**
- Consider vector change or dual sequential defibrillation for refractory v-fib. See diagram on next page.
- If obtain ROSC, take a moment to stabilize the patient (i.e. NOREPINEPHRINE drip, advanced airway, IV access, etc.) before initiating transport.
- If obtain ROSC in v-fib/v-tach arrest PRIOR to giving first dose of AMIODARONE, administer AMIODARONE 150 mg IV infusion over 10 minutes to decrease risk of recurrent v-fib/v-tach.
- If a patient is pregnant at or over 20 weeks estimated gestational age OR if the fundus is palpable above the umbilicus, apply the following interventions: During CPR, an additional rescuer should apply continuous manual leftward lateral displacement of the uterus to reduce pressure on the inferior vena cava and improve venous return. Vascular access should be obtained above the diaphragm. If no ROSC after two rounds of BLS/ACLS, consider immediate transport to the nearest Emergency Department for possible Resuscitative Cesarean Delivery (RCD).




### H's & T's (reversible causes)

- Hypovolemia – Volume infusion
- Hypoxia – Oxygenation & ventilation, CPR
- Hydrogen ion (acidosis) – Ventilation, CPR
- Hypokalemia
- Hyperkalemia – Calcium, Albuterol, Sodium bicarbonate
- Hypothermia – Warming
- Tension pneumothorax – Needle decompression
- Tamponade, cardiac – Volume infusion
- Toxins – Agent specific antidote
- Thrombosis, pulmonary – Volume infusion
- Thrombosis, coronary – Emergent PCI

### ENVIRONMENTAL EXPOSURE EXCEPTION:

All victims of electrocution and lightning strikes should have resuscitative efforts begun. Any decision to determine death in the field in these cases should be made **ONLY** after consultation with online medical control.



# High Performance CPR

## 1 IMMEDIATELY UPON ARRIVAL (or arrest, if witnessed) verify circulatory arrest by the absence of:

- consciousness
- carotid pulse
- normal/regular respiration (ignore agonal respirations)

2

## CHOOSE

2

If the cardiac arrest **IS NOT WITNESSED** by EMS providers:

- **PROVIDE 2 MIN OF CHEST COMPRESSIONS**
- **Compression Person** will immediately begin
- Continuous chest compressions for 2 minutes at a rate of 100-120 compressions/minute.
- **Do NOT interrupt chest compressions.**

If the cardiac arrest **IS WITNESSED** by EMS providers:

- **Compression Person** begin immediate chest compressions while **AED Person** attaches defibrillator.
- Once defibrillator is connected STOP CPR and push "ANALYZE" button.
- Continue starting as step 4 below.

3

## DEFIBRILLATION

**Designated AED Person will:**

1. TURN ON the AED as soon as cardiac arrest has been verified.
2. Cut clothing as necessary.
3. Place the AED pads on the patient in the appropriate locations:
  - Preferable anterior-posterior placement
  - Alternatively anterior-lateral placement

**Do NOT interrupt chest compressions**

4

## CHOOSE

4

**If "SHOCK ADVISED":**

- AED will charge.
- Deliver shock and immediately begin 2 min of CPR.
- Do NOT check for a pulse after shock.

**If "NO SHOCK ADVISED":**

- Check for a pulse (<10 seconds)
- If no pulse, immediately begin 2 min of CPR

5

## AIRWAY / VENTILATION

The **Designated Ventilation Person** will begin ventilations.

- Ventilate the patient at a rate of 30 compressions to 2 breaths (peds 15:2) unless an advanced airway is in place.
- Prepare suction equipment.
- Do NOT interrupt chest compressions.

**\*\*If there is no Designated Ventilation Person available, the Designated AED Person will immediately assume the responsibilities of the Designated Ventilation Person after the AED is operational.**

6

## IV/IO PLACEMENT


- If ALS personnel or BLS personnel with an IV/IO initiation endorsement are available, the **IV Person** will prepare the patient for IV/IO placement and begin to assemble the IV set.
- Consider IO access as this has less risk of interfering with chest compressions.

7

## ALS ARRIVAL

- Upon arrival of ALS, continue the current 2 min CPR cycle.
- Paramedics will pre-charge defibrillator and analyze/shock at the end of 2 min of CPR.
- After each 2 min CPR cycle paramedics should analyze or treat in <10 seconds.
- Continue cycles of 2 min CPR with breaks <10 seconds for analysis and/or treatment by paramedics.

# Chest Pain (Non Traumatic) and Suspected Acute Coronary Syndrome



### History

- Age
- Medications: Viagra, Levitra, Cialis, Revatio, Staxyn, etc
- Past medical history: MI, angina, diabetes, HTN, HLD
- Allergies
- Recent physical exertion
- Smoking history
- Time of onset, duration, repetition

### Signs and Symptoms

- CP, pressure, ache, vise-like pain, tight
- Location: substernal, epigastric, arm, jaw, neck, shoulder
- Radiation of pain
- Pale, diaphoresis
- Shortness of breath
- Nausea, vomiting
- Dizziness, lightheadedness

### Differential

- Trauma versus medical
- Anginal versus MI
- Pericarditis
- Pulmonary embolism
- Asthma, COPD
- Pneumothorax
- Aortic dissection or aneurysm
- GE reflux or hiatal hernia
- Esophageal spasm
- Chest injury or pain
- Pleural pain
- Drug overdose (cocaine, methamphetamine)

### Pearls


- Recommended exam: Mental Status, Skin, HEENT, Heart, Lungs, Abdomen, Back, Extremities, Neuro.
- Diabetics, geriatrics, and female patients often have atypical pain. Have a high index of suspicion.
- Perform a 12-Lead ECG on all patients 35 years old or older experiencing vague jaw/ chest/ abdominal discomfort.
- Perform a 12-Lead ECG within 10 minutes of patient contact. EMT and AEMT acquire and transmit for interpretation by a physician.
- Performance of serial ECGs is encouraged for symptomatic patients with initial ECG nondiagnostic for STEMI.
- The administration of NITROGLYCERIN is contraindicated for any patient who has used erectile dysfunction medications within the last 48-72 hours.
- NITROGLYCERIN is contraindicated in any patient with hypotension, bradycardia, or tachycardia in the absence of heart failure and evidence of a right ventricular infarction.
- Avoid the use of NITROGLYCERIN in patients with a suspected aortic dissection.
- MORPHINE should be used with caution in unstable angina (UA)/non-STEMI due to an association with increased mortality.
- A complete medication list should be obtained from each patient. It is especially important for the treating physician and healthcare providers to be informed if the patient is taking beta-blockers, calcium channel blockers, clonidine, digoxin, blood thinners (anticoagulants), and medications for the treatment of erectile dysfunction or pulmonary hypertension.

### QI Metrics


- 12-Lead ECG within 10 minutes of patient contact.
- Pain reassessed after every intervention.
- Pain control documented.

### 12-Lead ECG Lead Placement

- V1 – 4<sup>th</sup> ICS R-side
- V2 – 4<sup>th</sup> ICS L-side
- V3 – straight line between V2 and V4
- V4 – 5<sup>th</sup> ICS MCL
- V5 – same level as V4 at AAL
- V6 – same level as V4 at MAL



# Childbirth / Labor





### History

- Due date
- Time contractions started/ duration/frequency
- Rupture of membranes (meconium)
- Time and amount of any vaginal bleeding
- Sensation of fetal movement
- Pre-natal care
- Past medical and delivery history
- Medications
- Gravida/Para status
- High risk pregnancy

### Signs and Symptoms

- Spasmodic pain
- Contractions
- Vaginal discharge or bleeding
- Crowning, urge to push, or urge to move bowels
- Meconium

### Differential

- Abnormal presentation (breech, limb)
- Prolapsed cord
- Placenta previa
- Abruptio placenta
- Shoulder dystocia

### Pearls

- Recommended exam (of mother): Mental Status, Heart, Lungs, Abdomen, Neuro.
- Document all times (delivery, contraction duration and frequency).
- Some bleeding is normal; copious amounts of blood or free bleeding is abnormal. Deliveries complicated by maternal bleeding (placenta previa, vas previa, or placental abruption) place the infant at risk for hypovolemia secondary to blood loss.
- Resuscitation takes priority over recording APGAR scores. Record APGAR at 1 and 5 minutes after birth.
- APGAR of 7-10 is normal, while 4-6 requires resuscitative measures.
- Supine Hypotension Syndrome: if mother has hypotension before delivery, place patient in left lateral recumbent position or manually displace gravid uterus to the left in supine position.
- Do not routinely suction the infant's airway (even with a bulb syringe) during delivery.
- Dry, warm and stimulate all newborns to facilitate respirations and prevent hypothermia.
- Hypothermia is common in newborns and worsens outcomes of nearly all post-natal complications.
  - Ensure heat retention by drying the infant thoroughly, covering the head, and wrapping the baby in dry cloth.
  - When possible utilize "kangaroo care" (i.e., placing the infant skin-to-skin directly against mother's chest and wrapping them together) as an effective warming technique.

### APGAR

#### • Activity/Muscle Tone

#### Score=0

Absent

#### Score=1

Arms/legs flexed

#### Score=2

Active movement

#### • Pulse

Absent

Below 100

Above 100

#### • Grimace/Reflex Irritability

No response

Grimace

Sneeze, cough, pulls away

#### • Appearance/Skin Color

Blue-Grey, pale all over

Normal, except extremities

Normal over entire body

#### • Respiration

Absent


Slow, irregular

Good, crying

### Shoulder dystocia


If delivery fails to progress after head delivers, quickly attempt the following:

- Hyperflex mother's hips to severe supine knee-chest position (i.e., McRoberts' maneuver).
- Apply firm suprapubic pressure to attempt to dislodge shoulder. This often requires two EMS clinicians to perform and allows for delivery in up to 75% of cases.
- Attempt to angle baby's head as posteriorly as possible but NEVER pull.
- Continue with delivery as normal once the anterior shoulder is delivered.





# Cold-Related Illness



**History**

- Age, very young and old
- Exposure to decreased temperatures, but may occur in normal temperatures
- Past medical history/medications
- Drug or alcohol use
- Time of exposure/wetness/wind chill

**Signs and Symptoms**

- AMS/coma
- Cold, clammy
- Shivering
- Extremity pain
- Bradycardia
- Hypotension or shock
- Decreased respirations

**Differential**

- Sepsis
- Environmental exposure
- Hypoglycemia
- Stroke
- Head injury
- Spinal cord injury
- Trauma

**Pearls**

- Recommended exam: Mental Status, Heart, Lung, Abdomen, Extremities, Neuro.
- Pulse checks for patients suffering hypothermia should be performed for up to 60 seconds.
- Handle the patient gently to prevent triggering a ventricular arrest.
- Extremes of age are more prone to cold emergencies.
- Obtain and document patient temperature.
- If temperature is unknown, treat the patient based on suspected temperature.
- Active warming includes hot packs that can be used on the armpit and groin (care should be taken not to place the packs directly on the skin) and/or Ready Heat II thermal blanket or equivalent.
- Warm saline or lactated ringers IV may be used.

**Hypothermia Categories**

- Mild 90°- 95° F (33°- 35° C)
- Moderate 82°- 90° F (28°- 32° C)
- Severe 75°- 82° F (24°- 28° C)
- Profound <75 degrees F (<24° C)


**Hypothermia Mechanisms**

- Radiation
- Convection
- Conduction
- Evaporation

**Cardiac Arrest Resuscitation Guidelines for the Hypothermic Patient**

- Contraindications for initiation of resuscitation in the hypothermic patient:
  - Obvious fatal injuries (such as decapitation)
  - The patient exhibits signs of being frozen (such as ice formation in the airway)
  - Chest wall rigidity such that compressions are impossible
  - Danger to rescuers or rescuer exhaustion
  - Avalanche victims buried for 35 minutes or longer with airway obstruction by ice or snow
- Fixed and dilated pupils, apparent rigor mortis, and dependent lividity may not be contraindication for resuscitation in the severely hypothermic patient.
- The mainstay of therapy in severe hypothermia and cardiac arrest should be effective chest compressions and attempts at rewarming.
- Defibrillation may be attempted once. Do not make further attempts at defibrillation until the core temperature has increased to greater than 30°C (86°F).
- Medications should be withheld until the patient's core temperature is greater than 30°C (86°F). Intervals between medication provision should be doubled until the patient reaches 35°C (95°F), at which time, normal medication intervals may be adopted.
- If a hypothermic patient clearly suffered cardiac arrest and subsequently became hypothermic afterward with prolonged down time between arrest and rescue, there is no rationale for initiating resuscitation and warming the patient.

# Crashing Medical Patient



### History

- Blood loss – vaginal or gastrointestinal bleeding, AAA, ectopic
- Fluid loss – vomiting, diarrhea, fever
- Infection
- Cardiac ischemia (MI, CHF)
- Medications
- Allergic reaction/Anaphylaxis
- Pregnancy
- History of poor oral intake

### Signs and Symptoms

- Tachycardia out of proportion to temperature
- AMS
- Delayed capillary refill >2 sec
- Tachypnea
- Hypotension
- Cool/mottled or flushed/ruddy skin
- Tarry stool/hematochezia/hematemesis

### Differential

- Ectopic pregnancy
- Dysrhythmias
- Pulmonary embolus
- Tension pneumothorax
- Medication effect/Overdose
- Vasovagal
- Physiologic (pregnancy)
- Sepsis
- Post-ROSC


### Pearls

- Recommended exam: Mental Status, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro.
- Both EMT and paramedic training have traditionally emphasized the "load and go" approach for critically ill patients, often resulting in delayed resuscitation until arrival at the hospital. Historically, this was due to the lack of tools such as IOs and vasopressors in the prehospital setting, which necessitated rapid transport. However, new tools are now available, prompting a transition to a "stay and play" model for medical patients. Our goal is to reduce the incidence of EMS-witnessed cardiac arrests by prioritizing early IV/IO access, administering PUSH DOSE EPINEPHRINE, and utilizing pressure bags for intravenous fluids, all of which allows for patient stabilization on scene. This strategy aligns with current best practices in cardiac arrest management, as on-scene resuscitation typically yields better patient outcomes than interventions conducted in the back of a moving ambulance.
- It is crucial to note this approach does not extend to hypotensive trauma patients, for whom we continue to emphasize rapid transport to ensure timely, definitive care by trauma surgery.
- Shock describes a state of tissue under perfusion. There are multiple etiologies of shock, including hypovolemic (e.g. dehydration, blood loss), obstructive (e.g. tension pneumothorax, cardiac tamponade), distributive (e.g. sepsis, severe burns), and cardiogenic (e.g. heart failure, acute papillary muscle rupture). Rapid vasopressors and fluid resuscitation are the mainstay of treatment, as the duration of shock coincides with the extent of tissue damage.
- Signs of shock include:
  - Altered mental status
  - Respiratory distress
  - Pallor
  - Clammy/diaphoretic skin
- Treatment priorities for hypotensive medical patients:
  - 2 large bore (18 gauge or larger) IVs are preferred.
  - If unsuccessful in obtaining IV access, early IO placement is crucial. Distal femur and proximal humerus are both preferred over tibial IOs due to better flow rates.
  - PUSH DOSE EPINEPHRINE should be given early in patients with SBP <70 or signs of shock present.
  - Fluid resuscitation: Although often essential in shock, fluid may worsen the clinical picture in certain conditions, including cardiogenic shock. Monitor patients for signs of fluid overload when administering rapid fluid boluses.
    - Discontinue fluids if patient is developing signs of pulmonary edema or respiratory insufficiency/failure.
    - Utilize pressors early in suspected cardiogenic shock and signs of pulmonary edema.

### Other Considerations:

- Adrenal insufficiency: Patients may have a history of adrenal insufficiency related to congenital adrenal hyperplasia or from long-term daily steroid use. These patients may require stress dose steroids to maintain blood pressure in the setting of shock. Administer patient's own HYDROCORTISONE (SOLU-CORTEF) as prescribed or METHYLPREDNISOLONE 2 mg/kg IV/IO/IM, max 125 mg.
- Pregnancy: Pregnant patients can be assisted by shifting their gravid uterus left-ward, off of their inferior vena cava.

# Drowning



### History

- Submersion in liquid, regardless of depth
- Possible history of trauma (dive)
- Duration of immersion
- Temperature of water or possibility of hypothermia
- Degree of water contamination

### Signs and Symptoms

- Unresponsive
- Mental status changes
- Decreased or absent vital signs
- Vomiting
- Coughing, wheezing, rales, stridor, rhonchi
- Apnea
- Frothy/foamy sputum


### Differential

- Trauma
- Pre-existing medical condition
- Barotrauma
- Decompression illness
- Post-immersion syndrome

### Pearls

- Recommended exam: Trauma survey, Head, Neck, Chest, Abdomen, Back, Extremities, Skin, Neuro.
- Ensure scene safety.
- Resuscitation efforts should follow an **ABCs (Airway, Breathing, Circulation)** strategy for drowning victims in cardiac arrest. Initiate 5 rescue breaths followed by a ratio of 30 chest compressions to 2 breaths.
- Hypothermia is often associated with submersion incidents.
- All patients should be transported for evaluation because of potential for worsening over the next several hours.
- Decompression illness may have a variety of presentations depending on system affected (e.g., skin, joint(s), pulmonary, neurologic), and can occur even when a diver does not exceed dive table limits.
- Resuscitation recommendations based on water temperature and submersion time:
  - If water temperature is less than 43°F (6°C) and the patient is submerged with evidence of cardiac arrest: Survival is possible for submersion time less than 90 minutes and resuscitative efforts should be initiated.
  - If water temperature is greater than 43°F (6°C) and the patient is submerged with evidence of cardiac arrest: Survival is possible for submersion time less than 30 minutes and resuscitative efforts should be initiated.

# Epistaxis



### History

- Age
- Past Medical History (HTN)
- Medications (anticoagulants, antiplatelets, aspirin, NSAIDS)
- Previous episodes of epistaxis
- Trauma
- Duration of bleeding
- Quantity of bleeding

### Signs and Symptoms

- Bleeding from nasal passages
- Dizziness/Lightheadedness
- Chest pain
- SOB
- Pain
- Nausea
- Vomiting

### Differential


- Trauma
- Infection (viral URI or Sinusitis)
- Allergic rhinitis
- Lesions (polyps, ulcers)
- Hypertension

### Pearls

- Recommended exam: Mental Status, HEENT, Lungs, Neuro.
- It is very difficult to quantify the amount of blood loss with epistaxis.
- Bleeding may be also occurring posteriorly. Evaluate for posterior blood loss by examining the posterior pharynx.
- Anticoagulants include warfarin (Coumadin), heparin, enoxaparin (Lovenox), dabigatran (Pradaxa), rivaroxaban (Xarelto), apixaban (Eliquis) and many other over the counter headache relief medications.
- Anti-platelet agents like aspirin, clopidogrel (Plavix), aspirin/dipyridamole (Aggrenox), and ticlopidine (Ticlid) can contribute to bleeding.



# Heat-Related Illness



**History**

- Age, very old and young
- Exposure to increased temperatures and/or humidity
- Past medical history/medications
- Time and duration of exposure
- Poor PO intake, extreme exertion
- Fatigue and/or muscle cramping

**Signs and Symptoms**

- AMS/coma
- Hot, dry, or sweaty skin
- Hypotension or shock
- Seizures
- Nausea

**Differential**

- Fever, Sepsis
- Dehydration
- Medications
- Hyperthyroidism
- DTs
- Heat cramps, heat exhaustion, heat stroke
- CNS lesions or tumors

**Pearls**

- Recommended exam: Mental Status, Skin, Heart, Lung, Abdomen, Extremities, Neuro.
- Extremes of age are more prone to heat emergencies.
- Cocaine, amphetamines, and salicylates may elevate body temperatures.
- Sweating generally disappears as body temperatures rise over 104° F (40° C).
- Intense shivering may occur as patient is cooled. Paramedics can consider administering a low dose benzodiazepine.
- Active cooling includes application of cold packs or ice (not directly on skin), fanning, or air conditioning.
- Cold saline is not to be administered for the treatment of hyperthermia unless directed by online medical control.
- There is no evidence supporting EMS obtaining orthostatic vital signs as a clinical indicator.
- Cold water immersion is the preferred method of active cooling. Some providers such as certified athletic trainers and event medical personnel are prepared to initiate cold water immersion prior to EMS arrival. If cold water immersion was initiated due to documented hyperthermia, these patients should not be removed from cold water immersion prior to their rectal temperature reaching 102.2F (39C) or mental status returning to baseline unless it is required to manage other emergent issues such as airway.

**Heat Cramps**

- Consist of benign muscle cramping caused by dehydration and is not associated with an elevated temperature.

**Heat Exhaustion**

- Consists of dehydration, salt depletion, dizziness, fever, AMS, headache, cramping, N/V. Vital signs usually consist of tachycardia, hypotension and elevated temperature.

**Heat Stroke**


- Consists of dehydration, tachycardia, hypotension, temperature >104° F (40° C), and AMS. Potential for seizures or coma.

**Heat Syncope**


- Transient loss of consciousness attributed to heat exposure.

# Head Injury

## EPIC ALGORITHM FOR ADULTS



# Hyperkalemia (Suspected)



**History**

- History of renal failure
- History of dialysis
- Trauma, crush injury

**Signs and Symptoms**

- Cardiac conduction disturbances
- Irritability
- Abdominal distension
- Nausea
- Diarrhea
- Oliguria
- Weakness


**Differential**

- Cardiac disease
- Renal failure
- Dialysis
- Trauma, crush injury

**Pearls**

- Patients must have suspected hyperkalemia *OR* electrocardiographic findings consistent with hyperkalemia (bradycardia with widening QRS complexes) **BEFORE** initiating treatment.
- Hyperkalemia is defined as a potassium level higher than 5.5 mmol/L.
  - Potassium of 5.5 - 6.5 mmol/L - Tall peaked T waves.
  - Potassium of 6.5 - 7.5 mmol/L - Loss of P waves.
  - Potassium of 7.5 - 8.5 mmol/L - Widening QRS.
  - Potassium of >8.5 mmol/L - QRS continues to widen, approaching sine wave.

# Obstetrical Emergency



**History**

- Medical history
- Hypertension medication
- Prenatal care
- Prior pregnancies/births
- Previous pregnancy complications

**Signs and Symptoms**

- Vaginal bleeding
- Abdominal pain (epigastric or RUQ)
- Seizures
- Hypertension
- Severe headache
- Visual changes
- Edema of the hands or face
- Shortness of breath/Pulmonary edema
- Confusion/AMS
- Ecchymosis (bruising, petechiae)


**Differential**

- Pre-eclampsia/eclampsia
- Placenta previa
- Placenta abruptio
- Spontaneous abortion
- Ectopic pregnancy
- Chronic hypertension (PMHx)
- Pregnancy induced hypertension

**Pearls**

- Recommended exam: Mental Status, Heart, Lung, Abdomen, Neuro.
- Severe headache, vision changes or RUQ pain may indicate pre-eclampsia.
- In the setting of pregnancy: hypertension is defined as >140 systolic or >90 diastolic or a relative increase of 30 systolic or 20 diastolic from the patient's normal pre-pregnancy BP.
- Maintain left lateral position.
- Ask patient to quantify bleeding - number of pads used per hour.
- Any pregnant patient involved in a MVC should be seen by a physician for evaluation.
- Postpartum eclampsia/pre-eclampsia commonly presents up to 48 hours after childbirth. Often the presenting symptom is headache or SOB. If symptomatic, treat as eclampsia/pre-eclampsia.
- May present up to 6 weeks after childbirth, assess for history or pre-eclampsia/eclampsia during pregnancy or delivery.
- Magnesium toxicity presents as (progression): Hypotension followed by, Loss of deep tendon reflexes followed by, Somnolence, slurred speech followed by, Respiratory paralysis followed by, Cardiac arrest. Treatment includes stopping the MAGNESIUM drip and giving CALCIUM GLUCONATE 3 g IV or CALCIUM CHLORIDE 1 g IV in cases of pending respiratory arrest.

# Overdose / Poisoning





**History**

- Ingestion or suspected ingestion of a potentially toxic agent
- Substance ingested, route, quantity
- Time of ingestion
- Reason (suicidal, accidental, criminal)
- Available medications in home
- Past medical history, medications

**Signs and Symptoms**

- Mental status changes
- Hypotension/hypertension
- Decreased respiratory rate
- Tachycardia, dysrhythmias
- Seizures
- SLUDGE+BBB
- Malaise, weakness
- GI symptoms
- Dizziness
- Syncope
- Chest pain

**Differential**

- TCA overdose
- Acetaminophen
- Aspirin
- Depressants
- Stimulants
- Anticholinergic
- Cardiac medications
- Solvents, alcohols, cleaning agents, insecticides


**Pearls**

- Recommended exam: Mental Status, Skin, HEENT, Heart, Lung, Abdomen, Extremities, Neuro.
- NALOXONE should be administered in small increment doses IV to address respiratory depression and ensure adequate ventilation. Monitor patient to watch for any signs of respiratory depression reoccurring. IV/IM are preferred routes for predictability.
- Overdose or toxin patients with significant ingestion/exposure should be closely monitored and aggressively treated. Do not hesitate to contact medical control if needed.
- In the case of cyanide poisoning, altered mental status may be profound. Profound altered mental status can be defined as a deficit that includes disorientation, bewilderment and difficulty following commands.
- If patient is suspected to have narcotic overdose/hypoglycemia, administer NARCAN/GLUCOSE prior to supraglottic device/intubation.
- Poison Control: 1-800-222-1222.

**Agents**

- Acetaminophen: Initially normal or N/V. Tachypnea and AMS may occur later. Renal dysfunction, liver failure and/or cerebral edema may manifest.
- Depressants: Decreased HR, BP, temp and RR.
- Anticholinergic: Increased HR, increased temperature, dilated pupils and AMS changes.
- Insecticides: May include S/S of organophosphate poisoning.
- Solvents: N/V, cough, AMS.
- Stimulants: Increased HR, BP, temperature, dilated pupils, seizures, and possible violence.
- TCA: Decreased mental status, dysrhythmias, seizures, hypotension, coma, death.

# Pain Management



**History**

- Age
- Location, duration
- Severity (1-10)
- Past medical history
- Pregnancy status
- Drug allergies and medications

**Signs and Symptoms**

- Severity (pain scale)
- Quality
- Radiation
- Relation to movement, respiration
- Increased with palpation of area

**Differential**

- Musculoskeletal
- Visceral (abdominal)
- Cardiac
- Pleural, respiratory
- Neurogenic
- Renal (colic)


**Pearls**

- Recommended exam: Respiratory Status, Mental Status, Area of pain, Neuro.
- Pain severity (1-10) is to be recorded before and after medication administration and upon arrival at hospital.
- Monitor BP and respirations closely as sedative and pain control agents may cause hypotension and/or respiratory depression.
- Consider patient's age, weight, clinical condition, use of drugs/alcohol, exposure to opiates when determining initial opiate dosing. Weight based dosing may provide a standard means of dosing calculation, but it does not predict response. Consider starting at a lower initial dose and titrating to effect is recommended.
- Administer opioids with caution to patients with Glasgow Coma Score (GCS) less than 15, hypotension, identified medication allergy, hypoxia (SPO2 less than 90%) after maximal supplemental oxygen therapy, or signs of hypoventilation.
- Exercise care when administering opiates and benzodiazepines; this combination results in deeper anesthesia with significant risk of respiratory compromise. Utilize ETCO2 when giving combination.
- Burn patients may require more aggressive dosing.
- ACETAMINOPHEN and KETOROLAC are not to be used as the primary pain management medication for Chest Pain/ Suspected ACS or STEMI patients.
- ACETAMINOPHEN and/or KETOROLAC should be considered the primary treatment for severe pain for patients that do not wish to receive narcotic analgesia.
- Avoid non-steroidal anti-inflammatory medications such as IBUPROFEN and KETOROLAC in patients with NSAID allergy, aspirin-sensitive asthma, renal insufficiency, pregnancy, or known peptic ulcer disease. KETOROLAC should not be used in patients with hypotension (due to renal toxicity).
- Consider FENTANYL as the preferred opioid agent for traumatic pain.
- ACETAMINOPHEN may be given if it has been  $\geq 4$  hours since last dose of  $\leq 650$  mg or  $\geq 6$  hours since last dose of 1 gm. IBUPROFEN/KETOROLAC may be given if it has been  $\geq 6$  hours since last dose of  $\leq 600$  mg NSAID,  $\geq 8$  hours since last dose of 800 mg NSAID,  $\geq 12$  hours since last dose of Naproxen/Naprosyn, or  $\geq 24$  hours since last dose of Meloxicam.

**• QI Metrics**

- Vital signs with O<sub>2</sub> sats recorded.
- Pain scale documented before and after intervention.
- Vital signs repeated after intervention.
- If considering repeat administration of pain medications, nasal cannula capnography should be utilized. ETCO2 when giving combination opiates and benzodiazepines.

# Pulmonary Edema / CHF



**History**

- Congestive heart failure
- Past medical history
- Medications
- Cardiac history
- Home oxygen

**Signs and Symptoms**

- Respiratory distress, bilateral rales
- Apprehension, orthopnea
- JVD
- Pink, frothy sputum
- Peripheral edema
- Diaphoresis
- Hypotension, shock
- Chest pain

**Differential**

- MI
- Congestive heart failure
- Asthma
- Anaphylaxis
- Aspiration
- COPD
- Pleural effusion
- Pneumonia
- Pericardial tamponade
- Toxic exposure
- High altitude pulmonary edema (HAPE)
- Opioid overdose
- Kidney failure or dialysis noncompliance


**Pearls**

- Recommended exam: Mental Status, HEENT, Skin, Neck, Heart, Lungs, Abdomen, Extremities, Neuro.
- Allow patient to maintain position of comfort.
- The administration of NITROGLYCERIN is contraindicated for any patient who has used erectile dysfunction medications within the last 48-72 hours.
- Pulse oximetry and EtCO<sub>2</sub> should be routinely used as adjuncts to other forms of monitoring in patients with respiratory complaints.
- 12-lead ECG may be indicated to assess for dysrhythmia or ischemia, particularly in patients with risk factors for coronary artery disease and/or presentation consistent with CHF.
- Pulmonary edema is more commonly a problem of volume distribution than total body fluid overload, so administration of diuretics provide no immediate benefit for most patients and can cause significant harm.
- Nitrates provide both subjective and objective improvement, and might decrease intubation rates, incidence of MIs, and mortality. High-dose nitrates can reduce both preload and afterload and potentially increase cardiac output and blood pressure.

**QI Metrics**

- Blood pressure reassessed after each NITROGLYCERIN dose.
- ETCO<sub>2</sub> monitored.

# Respiratory Distress



**History**

- PMHx: asthma, COPD, CHF, chronic bronchitis, emphysema
- Home treatment (oxygen, nebulizers)
- Medication
- Toxic exposure

**Signs and Symptoms**

- Shortness of breath
- Pursed lip breathing
- Decreased ability to speak
- Increased respiratory rate and effort
- Wheezing, rhonchi
- Use of accessory muscles
- Fever, cough
- Tachycardia


**Differential**

- Asthma
- Anaphylaxis
- Aspiration
- COPD
- Pleural effusion
- Pneumonia
- Pulmonary embolus
- Pneumothorax
- Cardiac (MI or CHF)
- Pericardial tamponade
- Hyperventilation
- Inhaled toxin

**Pearls**

- Recommended exam: Mental Status, HEENT, Skin, Neck, Heart, Lungs, Abdomen, Extremities, Neuro.
- Pulse oximetry and end tidal continuous waveform capnography should be monitored.
- Consider MI.
- Allow the patient to assume a position of comfort.
- In the setting of severe bronchoconstriction, wheezing may not be heard. Patients with known asthma with severe dyspnea should be empirically treated, even if wheezing is absent.
- A “shark fin” on waveform capnography suggests significant bronchospasm and obstructive physiology.
- Invasive airways do not improve bronchospasm. The airway should be managed in the least invasive way possible. Supraglottic devices and endotracheal intubation should be considered only if BVM ventilation fails.
- Positive pressure ventilation in the setting of bronchoconstriction, either via a supraglottic airway or intubation, increases the risk of air trapping which can lead to pneumothorax and cardiovascular collapse. These interventions should be reserved for situations of respiratory failure.

# Seizure





**History**

- Reported or witnessed seizure activity
- Previous seizure history
- Seizure medications
- History of trauma
- History of diabetes
- History of pregnancy
- Time of seizure onset
- Number of seizures
- Alcohol use, abuse, or abrupt cessation
- Fever

**Signs and Symptoms**

- Decreased mental status
- Sleepiness
- Tongue trauma
- Incontinence
- Observed seizure activity
- Evidence of trauma
- Unconsciousness

**Differential**

- CNS trauma
- Tumor
- Metabolic, hepatic or renal failure
- Hypoxia
- Electrolyte abnormality (Na, Ca, Mg)
- Drugs, medication non-compliance
- Infection, fever
- Alcohol withdrawal
- Eclampsia
- Stroke
- Hyperthermia
- Hypothermia

**Pearls**

- Recommended exam: Mental Status, HEENT, Heart, Lungs, Extremities, Neuro.
- Benzodiazepines are effective in terminating seizures; do not delay IM/IN administration while initiating an IV.
- Recent evidence supports the use of MIDAZOLAM IM as an intervention that is at least as safe and effective as intravenous LORAZEPAM for prehospital seizure cessation.
- Status epilepticus is defined as two or more seizures successively without an intervening lucid period, or a seizure lasting over five minutes.
- Grand mal seizures (generalized) are associated with loss of consciousness, incontinence and oral trauma.
- Focal seizures affect only part of the body and are not usually associated with a loss of consciousness.
- Be prepared to address airway issues and support ventilations as needed. Many airway/breathing issues in seizing patients can be managed without intubation or placement of an advanced airway. Reserve these measures for patients that fail less invasive maneuvers as noted above.
- Consider ETCO<sub>2</sub> monitoring.
- For new onset seizures or seizures that are refractory to treatment, consider other potential causes including, but not limited to, trauma, stroke, electrolyte abnormality, toxic ingestion, pregnancy with eclampsia, hyperthermia.

# Sepsis (Suspected)



General Adult Assessment

FR

Oxygen Keep SpO<sub>2</sub> >94%

E\*

Vascular Access

P

Cardiac Monitoring/Capnography

Does the patient have a known or suspected source of infection AND TWO of the following criteria?

SBP <90 mm Hg

HR >90/min

Respiratory rate >20/min

Altered mental status

Temperature > or = 100.4 F or < or = 96.8 F

Persistent ETCO<sub>2</sub> <25 on waveform capnography

YES

NO

CODE SEPSIS notification immediately upon recognition of patient meeting Sepsis criteria

Alternate appropriate treatment guidelines as indicated

E\*

LR or NS bolus 500 ml IV/IO, reassess criteria and re-examine. May repeat x3 for SBP <90, with no rales on lung exam. Max dose 2000 ml.

If SBP <90 after 2000 ml IV fluids:

NOREPINEPHRINE 0.05-0.5 mcg/kg/min IV drip  
titrate to maintain SBP >90 or MAP >65 OR;

P

PUSH DOSE EPINEPHRINE 1:100,000  
10 mcg IV/IO, may repeat q 2-5 min to maintain SBP >90  
(1 ml of a 1:100,000 solution)  
AND/OR EPINEPHRINE 0.05-0.3 mcg/kg/min IV drip



Refer to Push Dose Epinephrine Guideline pg 156 for mixing instructions



Continue General Adult Assessment

### History

- Age (Common in elderly and very young)
- Presence and duration of fever
- Previously documented infection or illness (UTI, Pneumonia, meningitis, encephalitis, cellulitis, abscesses, etc)
- Recent surgery or invasive procedure
- Any recent hospitalization
- Immunocompromised (transplant, HIV, diabetes, cancer)
- Bedridden or immobile patients
- Prosthetic or indwelling devices
- Immunization status
- Open wounds, even minor ones

### Signs and Symptoms

- Hyper or hypothermia
- Rash and/or excessive bruising
- Fever, Chills
- Myalgia
- Markedly decreased urine output
- Altered mentation
- Delayed capillary refill
- Elevated blood glucose (unless diabetic)

### Differential

- Cardiogenic Shock
- Hypovolemic Shock
- Dehydration
- Hyperthyroidism
- Medication/drug interaction
- Non-septic infection
- Allergic reaction/anaphylaxis
- Toxicological emergency
- Withdrawal syndromes


### Pearls

- Early recognition of sepsis allows for attentive care and early administration of antibiotics.
- Aggressive IV fluid therapy is the most important prehospital treatment for sepsis. Suspected septic patients should receive repeated fluid boluses (to a max total of 2 liters) while being checked frequently for signs of pulmonary edema, especially those patients with known history of CHF or ESRD on dialysis. STOP fluid resuscitation in the setting of pulmonary edema.
- Time IV fluid bolus was initiated and total amount given is to be recorded and reported to hospital staff at patient hand off.
- Septic patients are especially susceptible to traumatic lung injury and ARDS. If artificial ventilation is necessary, avoid ventilating with excessive tidal volumes. If CPAP is utilized, airway pressure should be limited to 5 cm H2O.
- Attempt to identify source of infection (skin, respiratory, etc.) and relay previous treatments and related history to the ED physician and nursing staff.
- Elevated serum lactate levels are a useful marker of hypoperfusion in sepsis and often become elevated prior to the onset of hypotension. ETCO2 levels are inversely proportional to serum lactate levels.
- Disseminated intravascular coagulation (DIC) is an ominous, late stage manifestation of sepsis characterized by frank, extensive bruising, bleeding from multiple sites, and finally tissue death.
- Conditions such as Crohn's, psoriasis, rheumatoid arthritis and other autoimmune disorders are now being treated with medications that impair the immune system. These patients need to be considered as immunocompromised.
- Hypovolemia or distributive shock should be addressed with a fluid bolus prior to the administration of vasopressors.
- NOREPINEPHRINE is the preferred vasopressor drip for septic shock.

### QI Metrics

- Vital signs to include blood pressure, heart rate, respiratory rate, SpO2, and EtCO2 documented throughout transport.
- Vital signs before, during, and after fluid administration.
- Documentation of the time IV fluid was started and total amount given.

# Shock



### History

- Blood loss - vaginal bleeding, ectopic, GI bleeding or AAA
- Fluid loss - vomiting, diarrhea, fever
- Infection
- Cardiac problems
- Medications
- Allergic reaction
- Pregnancy
- History of poor oral intake
- Adrenal insufficiency

### Signs and Symptoms

- Restlessness, confusion
- Weakness, dizziness
- Weak rapid pulse
- Pale, cool, clammy skin
- Delayed capillary refill
- Hypotension
- Coffee-ground emesis
- Tarry stools

### Differential

- Hypovolemic shock
- Cardiogenic shock
- Septic shock
- Neurogenic shock
- Anaphylactic shock
- Ectopic pregnancy
- Dysrhythmias
- Pulmonary embolism
- Tension pneumothorax
- Medication effect or overdose
- Vasovagal
- Physiologic (pregnancy)

### Pearls

- Recommended exam: Mental Status, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro.
- Shock often is present with normal vital signs and may develop insidiously. Tachycardia may be the only manifestation.
- Hypotension can be defined as a systolic BP of <90. This is not always reliable and should be interpreted in context and patient's typical BP, if known. Shock may present with a normal BP initially.
- Normal Shock Index (SI) is less than 0.7. A Shock Index (SI) greater than 1.0 is indicative of shock and need for aggressive resuscitation. [SI = HR / SBP]
- An ETCO<sub>2</sub> measurement of <25 mm/hg is indicative of shock.
- Hypovolemia or distributive shock should be addressed with a fluid bolus prior to the administration of vasopressors.
- If the patient's condition deteriorates after fluid administration, rales or hepatomegaly develop, then consider cardiogenic shock and withholding further fluid administration.
- While there are no absolute contraindications to epinephrine, it should be used with caution in elderly patients, patients with known cardiovascular disease, or significant tachycardia or hypertension, and should be administered only when the patient's signs and symptoms are severe.
- Consider all possible causes of shock and treat per appropriate protocol.

### Hypovolemic shock

- Hemorrhage, trauma, GI bleeding, ruptured aortic aneurysm, or pregnancy related bleeding

### Cardiogenic shock

- Heart failure, MI, cardiomyopathy, myocardial contusion, toxins, dysrhythmia, valvular disease

### Distributive shock

- Sepsis, anaphylaxis, neurogenic, toxins, overdose, endocrine

### Obstructive shock

- Pericardial tamponade, pulmonary embolus, tension pneumothorax

**For patients with known adrenal insufficiency, administer patient's own  
HYDROCORTISONE (SOLU-CORTEF) as prescribed OR;**



**METHYLPREDNISOLONE 2 mg/kg IV/IO/IM, max 125 mg**

### Causes of Adrenal Insufficiency:


**Addison's Disease**

**Congenital Adrenal Hyperplasia**

**Long term administration of steroids**

**Others**

# Smoke Inhalation



### History

- Exposed to smoke in a structure fire
- Exposed to smoke in a vehicle fire
- Exposed to smoke from other sources, industrial, confined space, wilderness fire, etc.

### Signs and Symptoms

- Facial burns
- Singed nasal hairs or facial hair
- Shortness of breath
- Facial edema
- Stridor
- Grunting respirations

### Differential

- COPD
- CHF
- Toxic inhalation injury
- Caustic inhalation injury
- Pneumonitis

### Pearls

- Recommended exam: Mental Status, HEENT, Heart, Lungs, Abdomen, Back, Extremities, Neuro.
- Protect yourself and your crew.
- Have a high index of suspicion when treating patients at the scene of a fire.
- If the medication is not available on scene do not delay transport waiting for it.
- Carefully monitor respiratory effort and correct life threats immediately.
- Decide early on if you want to intubate as burned airways swell, making intubation difficult.
- Profound altered mental status can be defined as a deficit that includes disorientation, bewilderment and difficulty following commands.

### Preparation and Administration of HYDROXOCOBALAMIN


#### Complete Starting Dose: 5 g

**1. Reconstitute:** Place the vial in an upright position. Add **200 mL** of 0.9% Sodium Chloride Injection\* to the vial using the transfer spike. **Fill to the line.**


\* 0.9% Sodium Chloride Injection is the recommended diluent (diluent not included in the kit). Lactated Ringer's Solution and 5% Dextrose Injection have also been found to be compatible with HYDROXOCOBALAMIN.

**2. Mix:** The vial should be repeatedly inverted or rocked, **NOT** shaken, for at least **60 seconds** prior to infusion.

**3. Infuse Vial:** Use vented intravenous tubing, hang and infuse over **15 minutes**.



# STEMI (Suspected)





### History

- Age
- Medication: Viagra, Levitra, Cialis, Revatio, Staxyn, etc.
- Past Medical History: MI, angina, diabetes, HTN, HLD
- Allergies
- Recent physical exertion
- Time of onset, duration

### Signs and Symptoms

- CP, pressure, ache, vice-like pain, tight, heavy
- Location: substernal, epigastric, arm, jaw, neck, shoulder
- Radiation of pain
- Pale, diaphoresis
- Shortness of breath
- Nausea, vomiting
- Dizziness, lightheadedness

### Differential

- Trauma versus medical
- Angina versus MI
- Pericarditis
- Pulmonary embolism
- Asthma, COPD
- Pneumothorax
- Aortic dissection or aneurysm
- GE reflux or hiatal hernia
- Esophageal spasm
- Chest injury or pain
- Pleural pain
- Drug overdose (cocaine, methamphetamines)

### Pearls

- Recommended exam: Mental Status, Skin, HEENT, Heart, Lungs, Abdomen, Back, Extremities, Neuro.
- Diabetics, geriatrics, and female patients often have atypical pain. Have a high index of suspicion.
- Perform a 12-Lead ECG on all patients 35 years old and older experiencing vague jaw/ chest/ abdominal discomfort.
- Perform a 12-Lead ECG within 10 minutes of patient contact.
- The administration of NITROGLYCERIN is contraindicated for any patient who has used erectile dysfunction medications within the last 48-72 hours.


### STEMI BYPASS PROCEDURE

- Applicable to BSFD, YMCRFD, and HBRFD *only* when bypassing BSMC and transporting (ground or flight) directly to hospital with cardiac catheterization capability (i.e. BHDRMC, EIRMC).
- Perform a 12-Lead ECG within 10 minutes of patient contact and transmit to base hospital or receiving facility.
- Once physician confirms STEMI on 12-Lead ECG, administer:
  - BRILINTA 180 mg PO
  - HEPARIN 60 units/kg IV bolus (max 4000 units)

### QI Metrics

- 12-Lead ECG within 10 minutes of patient contact.
- Pain reassessed with every intervention.
- Pain control documented.

# Stroke (CVA)



### History

- Previous CVA, TIAs
- Previous cardiac/vascular surgery
- Associated diseases: diabetes, HTN, CAD
- Atrial Fibrillation
- Medications: anticoagulant, antiplatelet
- History of recent trauma
- History of recent surgery
- History of recent hemorrhage

### Signs and Symptoms

- Facial droop
- Weakness, paralysis
- Blindness or other vision change
- Aphasia, dysarthria
- Vertigo, dizziness
- Vomiting
- Headache
- Neck pain/stiffness
- Seizures
- Hypertension, hypotension

### Differential

- AMS
- TIA
- Seizure
- Hypoglycemia
- Sepsis
- Migraine
- Intoxication
- Tumor
- Trauma
- Dialysis/ Renal Failure

### Pearls

- Recommended exam: Mental Status, HEENT, Heart, Lungs, Abdomen, Extremities, Neuro.
- Determine time of “last known well” (onset) or last time patient was seen normal.
- Three most predictive findings for diagnosis of stroke: facial droop, arm drift or weakness, abnormal speech.
- Activate CODE STROKE for “last known well” under 4.5 hours or signs of LVO stroke under 24 hours. Note – a ‘wake up stroke’ is an indication for CODE STROKE as still may be a thrombolytics candidate based on MRI showing DWI-positive and FLAIR-negative mismatch.
- Transport to an appropriate Stroke Center or Endovascular Treatment Center if possible.
- Prevent aspiration by elevating head of stretcher 15–30 degrees if SBP greater than 100 mmHg.
- Avoid multiple IV attempts.

### QI Metrics

- Complete the prehospital stroke scale and RACE or VAN assessment in less than five minutes.
- Time of “last known well” (onset) documented.
- Blood glucose documented.
- 12-Lead EKG completed.
- Scene time <10 minutes.
- Telemetry to receiving facility.

## Cincinnati Pre-hospital Stroke Scale

**1. FACIAL DROOP:** Have patient show teeth or smile.



**Normal:**  
both sides  
of the face  
move equally



**Abnormal:**  
one side of  
face does not  
move as well  
as the other  
side

**2. ARM DRIFT:** Patient closes eyes & holds both arms out for 10 sec.



**Normal:**  
both arms  
move the  
same or both  
arms do not  
move at all



**Abnormal:**  
one arm does  
not move or  
drifts down  
compared to  
the other

**3. ABNORMAL SPEECH:** Have the patient say “you can’t teach an old dog new tricks.”

**Normal:** patient uses correct words with no slurring

**Abnormal:** patient slurs words, uses the wrong words, or is unable to speak

**INTERPRETATION:** If any 1 of these 3 signs is abnormal, the probability of a stroke is 72%.

# If you suspect a stroke, BE FAST:



## Balance

Does the person have a loss of balance?



## Eyes

Has the person lost vision in one or both eyes?



## Face

Does the person's face look uneven?



## Arms

Can the person raise both arms for 10 seconds?



## Speech

Is the person's speech slurred?



## Time

Time is brain. Call 911 if you suspect a stroke.

**Weakness  
PLUS...**

**V Vision**

**A Aphasia**

**N Neglect**

## Rapid Arterial Occlusion Evaluation (RACE) Scale

### An EMS Assessment Tool for Acute Ischemic Stroke

(Sensitivity 85%, Specificity 68%)

Test Item	Score = 0	Score = 1	Score = 2	Patient Score
Facial Palsy	Absent	Mild	Moderate/Severe	
Arm Motor	Normal/Mild	Moderate	Severe	
Leg Motor	Normal/Mild	Moderate	Severe	
Head/Gaze Deviation	Absent	Present	N/A	
Aphasia* (if righthemiparesis)	Performs Both Tasks	Performs 1 Task	Performs Neither Tasks	
Agnosia* (if lefthemiparesis)	Patient Recognizes Arm and Impairment	Unable to Recognize Arm or Impairment	Unable to Recognize BOTH Arm and Impairment	
			<b>TOTAL SCORE = (0-9)</b>	

\*Aphasia: Ask the patient to: 1. "Close your Eyes" AND 2. "Make a Fist"


\*Agnosia: Ask the patient and evaluate recognition of deficit:

1. While showing paretic arm: "Whose arm is this?"
2. Ask patient: "Can you lift both arms and clap?"

**If RACE Score = 5 or greater, patient may have an ischemic stroke with a large vessel occlusion**

Reference:  
Natalia Pérez de la Ossa, et al. (2014). Design and Validation of a Prehospital Stroke Scale to Predict Large Arterial Occlusion: The Rapid Arterial Occlusion Evaluation Scale. *Stroke*, 45, 87-91. Retrieved from <http://stroke.ahajournals.org/content/45/1/87.full>

# Tachycardia / Stable (Normal Mental Status, Palpable Radial Pulse)



**History**

- Medications (aminophylline, diet pills, thyroid supplements, decongestants, digoxin)
- Diet (caffeine)
- Drugs (cocaine, methamphetamines)
- Past medical history: CHF, dysrhythmia

**Signs and Symptoms**

- Heart rate >150
- Dizziness, lightheadedness
- CP
- SOB, dyspnea
- Diaphoresis
- Palpitations
- Syncope/near syncope
- AMS
- Hemodynamic compromise

**Differential**


- Heart disease (WPW, valvular)
- Sick sinus syndrome
- MI
- Electrolyte imbalance (acidosis, hypokalemia, hyperkalemia)
- Exertion, fever, pain, emotional stress
- Hypoxia
- Hypovolemia
- Drug effect, overdose
- Hyperthyroidism
- Hypoglycemia
- Hypothermia
- Tension pneumothorax
- Cardiac tamponade
- Thrombus
- Trauma

**Pearls**

- Recommended exam: Mental Status, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro.
- Carefully monitor patients as you treat them; stable tachycardia may convert to unstable rhythms/conditions quickly.
- A-fib rarely requires cardioversion in the field. As it is difficult to ascertain the onset of this rhythm, the risk of stroke needs to be considered prior to cardioversion.
- A wide-complex irregular rhythm should be considered pre-excited A-fib (e.g., Wolff-Parkinson-White Syndrome, Lown-Ganong-Levine Syndrome). Extreme care must be taken in these patients. Avoid AV nodal blocking agents such as ADENOSINE, calcium channel blockers, digoxin, and beta-blockers in patients with pre-excitation A-fib because these drugs may cause a paradoxical increase in the ventricular response and potential ventricular fibrillation. Treatment in order of preference is: PROCAINAMIDE, synchronized cardioversion, and potentially AMIODARONE.
- Use of METOPROLOL can worsen CHF, chronic obstructive pulmonary disease (COPD), asthma, as well as hypotension and bradycardia.
- Patients who receive beta-blockers (e.g., METOPROLOL) with calcium channel blockers (e.g., DILTIAZEM) are at increased risk for hypotension and bradycardia.
- Sedate patients prior to cardioversion, if time allows. Preference for anterior-posterior pad placement. Preference for single high energy (at least 200 J) resulting in highest success rate for single shock.



# Tachycardia / Unstable (Mental Status Changes, No Palpable Radial Pulse)



**History**

- Medications (aminophylline, diet pills, thyroid supplements, decongestants, digoxin)
- Diet (caffeine)
- Drugs (cocaine, methamphetamines)
- Past medical history: CHF, dysrhythmia

**Signs and Symptoms**

- Heart rate >150
- Dizziness, lightheadedness
- CP
- SOB, dyspnea
- Diaphoresis
- Palpitations
- Syncope/near syncope
- AMS
- Hemodynamic compromise

**Differential**

- Heart disease (WPW, valvular)
- Sick sinus syndrome
- MI
- Electrolyte imbalance (acidosis, hypokalemia, hyperkalemia)
- Exertion, fever, pain, emotional stress
- Hypoxia
- Hypovolemia
- Drug effect, overdose
- Hyperthyroidism
- Hypoglycemia
- Hypothermia
- Tension pneumothorax
- Cardiac tamponade
- Thrombus
- Trauma

**Pearls**

- Recommended exam: Mental Status, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro.
- If patient is in arrest, efforts should focus on quality chest compressions and rhythm correction.
- Administer ADENOSINE at a proximal large bore IV site, rapidly followed by a saline flush.
- Sedate patients prior to cardioversion/defibrillation, if time allows. Preference for anterior-posterior pad placement. Preference for single high energy (at least 200 J) resulting in highest success rate for single shock.



# Ventilation Management

Use supplemental oxygen to maintain an oxygen saturation of >94% or >90% for patients on home oxygen for chronic conditions

**FR**

**E**

**Basic Airway Maneuvers**

- Open Airway: Chin Lift/Jaw Thrust
- NPA (EMT) or OPA (EMR) as needed
- Suction as needed

Consider spinal motion restriction (SMR)

Consider Altered Mental Status/Syncope

**Respiratory Distress and/or Tracheostomy Tube Replacement Guidelines if needed**

**FR**

**Administer oxygen**

**BVM as needed**

No Yes

**Intervention effective?**

**E\***

**Supraglottic Airway**

**Obtain Vascular Access**

**ETCO2 numerical monitoring**

**Endotracheal Intubation**

**ECG Monitor / Capnography**

**P**


**Consider Post-Intubation Sedation: Administer**

**MIDAZOLAM 0.1 mg/kg IM/IV/IO, max dose 5 mg. May repeat every 5 min OR;**

**KETAMINE 2 mg/kg IV, max 100 mg. May repeat every 10 min.**

**Consider Post-Intubation Analgesia: Administer**

**FENTANYL 1 mcg/kg, max dose 100 mcg. May repeat every 10 min.**



Always weigh the risks and benefits of endotracheal intubation in the field against transport. All prehospital endotracheal intubations are considered high risk. If ventilation/oxygenation is adequate, transport may be the best option. The most important airway device and the most difficult to use correctly and effectively is the Bag Valve Mask (not the laryngoscope). Few prehospital airway emergencies cannot be temporized or managed with proper BVM techniques.

#### **DIFFICULT AIRWAY ASSESSMENT:**

**Difficult BVM Ventilation – MOANS:** Difficult Mask seal due to facial hair, anatomy, blood or secretions/trauma; **O**beses or late pregnancy; **A**ge >55; **N**o teeth (roll gauze and place between gums and cheeks to improve seal); **S**tiff or increased airway pressures (asthma, COPD, obese, pregnant).

**Difficult Laryngoscopy – LEMON:** Look externally for anatomical distortions (small mandible, short neck, large tongue); Evaluate 3-3-2 Rule (Mouth open should accommodate 3 patient fingers, mandible to neck junction should accommodate 3 patient fingers, chin-neck junction to thyroid prominence should accommodate 2 patient fingers); **M**allampati (difficult to assess in the field); **O**bstuction / **O**beses or late pregnancy; **N**eck mobility.

**Difficult Supraglottic Device Placement – RODS:** Restricted mouth opening; **O**bstuction / **O**beses or late pregnancy; Distorted or disrupted airway; Stiff or increased airway pressures (asthma, COPD, obese, pregnant).

#### **Pearls**

- **The Montana Board of Medical Examiners DOES NOT allow drug assisted intubation (DAI) or rapid sequence intubation (RSI) for standard ground ALS paramedics.**
- Consider preoxygenation/lung denitrogenating with a non-rebreather, a nasal cannula at 15 LPM, CPAP, or BVM prior to intubation (as patient condition allows).
- Severe hypotension (SBP<90) should be addressed with IV fluids and/or pressors (as appropriate) prior to intubation in order to reduce the likelihood of post-intubation cardiovascular decline.
- Capnometry (color) or capnography (waveform) is mandatory with all methods of intubation. Document results.
- Continuous capnography (ETCO<sub>2</sub>) is mandatory for the monitoring of all patients with an advanced airway.
- If an effective airway is being maintained by BVM and/or basic airway adjuncts (e.g. nasopharyngeal airway) with continuous pulse oximetry values of ≥90% or values expected based on pathophysiologic condition with otherwise reassuring vital signs (e.g. pulse oximetry of 85% with otherwise normal vitals in a post-drowning patient), it is acceptable to continue with basic airway measures instead of using a supraglottic airway device or intubation. Consider CPAP or HFNC as indicated by guidelines and patient condition.
- For the purposes of this guideline, a secure airway is achieved when the patient is receiving appropriate oxygenation and ventilation.
- An intubation attempt is defined as passing the laryngoscope blade or endotracheal tube past the teeth or inserted into the nasal passage.
- An appropriate ventilatory rate is one that maintains an ETCO<sub>2</sub> of 35 - 45. Avoid hyperventilation.
- Paramedics should use a supraglottic airway device if oral-tracheal intubation is unsuccessful.
- Maintain C-spine stabilization for patients with suspected spinal injury.
- Gastric tube placement should be considered in all intubated patients if time allows.

#### **QI Metrics**


- Mandatory notification of Medical Director within 24 hours of any failed intubation attempt. For simplicity, may use this online form: <https://forms.gle/G3RPJQ8fEugRafMTA>
- Mandatory notification of Medical Director within 24 hours of any attempted, successful, and/or failed cricothyroidotomy in the field. For simplicity, may use this online form: <https://forms.gle/G3RPJQ8fEugRafMTA>

# **PEDIATRIC TREATMENT GUIDELINES**

**For patients who weigh up to 40 kg or  
< 15 years of age.**

**Pediatric treatment guidelines are to be used on children who have not yet experienced puberty. Signs of puberty include chest or underarm hair on males, and any breast development in females.**


# General Pediatric Assessment




#### Pearls

- For all scenes where patient needs exceed available EMS resources, initial assessment and treatment shall be in accordance with an approved triage methodology.
- Correct life-threatening problems as identified.
- If the ability to adequately ventilate a patient cannot be established, the patient must be transported to the nearest emergency department.
- Never withhold oxygen from a patient in respiratory distress.

## PEDIATRIC ASSESSMENT TRIANGLE



# General Pediatric Trauma Assessment



**History**

- Time and mechanism of injury
- Damage to structure or vehicle
- Location in structure or vehicle
- Others injured or dead
- Speed and details of MVC
- Restraints/protective equipment
- Past medical history
- Medications

**Signs and Symptoms**

- DCAP-BTLS
- AMS or unconscious
- Hypotension or shock
- Arrest


**Differential (Life threatening)**

- Tension pneumothorax
- Flail chest
- Pericardial tamponade
- Open chest wound
- Hemothorax
- Intra-abdominal bleeding
- Pelvis/femur fracture
- Spine fracture/cord injury
- Head injury
- Extremity fracture
- HEENT (airway obstruction)
- Hypothermia

**Pearls**

- Recommended exam: Mental Status, Skin, HEENT, Heart Lung, Abdomen, Extremities, Back, Neuro.
- Transport destination is based on the Trauma Field Triage Criteria Guideline.
- Transport should not be delayed for procedures; ideally procedures should be performed enroute when possible. Target scene time less than 10 minutes for unstable patients or those likely to need surgical intervention.
- BVM is an acceptable method of ventilating and managing an airway if pulse oximetry can be maintained  $\geq 90\%$ .
- Pediatric patients should be evaluated with a high index of suspicion; occult injuries may be present and pediatric patients can decompensate quickly.

# Pediatric Abdominal Pain, Nausea & Vomiting



**\*\*Round up to nearest ½ pill**



**History**

- Age
- Medical/surgical history
- Onset
- Quality
- Severity
- Fever
- Travel history
- Other sick contacts

**Signs and Symptoms**

- Pain location
- Tenderness
- Nausea
- Vomiting
- Diarrhea
- Dysuria
- Constipation
- Lack of appetite
- AMS


**Differential**

- Liver (Hepatitis)
- Gastritis
- Pancreatitis
- Kidney stone
- Appendicitis, mesenteric adenitis
- Bladder infection
- Bowel obstruction
- Intussusception
- Gastroenteritis
- Diabetic ketoacidosis (DKA)
- CNS (Increased pressure, headache, stroke, CNS lesions, trauma or hemorrhage)

**Pearls**

- Recommended Exam: Mental Status, Skin, HEENT, Heart, Lung, Abdomen, Back, Extremities, Neuro.
- Document mental status and vital signs prior to administration of antiemetics & pain management.
- Repeat vital signs after each fluid bolus.
- Consider retroperitoneal palpation for kidney pain.
- Pediatric fluid bolus is 20 ml/kg; may repeat to a maximum of 60 ml/kg.
- New onset DKA in pediatric patients commonly presents with nausea, vomiting, abdominal pain, and/or urinary frequency. Overly aggressive administration of fluid in hyperglycemic patients may cause cerebral edema or dangerous hyponatremia. Cerebral edema is a leading cause of death in children with DKA.
- If there is suspicion that the patient is in DKA, do not exceed 10 ml/kg NS bolus x1.
- MORPHINE is not recommended in children for abdominal pain.
- Consider cardiac and ETCO<sub>2</sub> monitoring.
- Abdominal pain in children less than 2 years old may be a harbinger for severe illness.

# Pediatric Allergic Reaction



### History

- Onset and location
- Insect sting or bite
- Food allergy/exposure
- Medication allergy/exposure
- New clothing, soap, detergent
- Past history of reactions
- Past medical history
- Medication history

### Signs and Symptoms

- Itching or hives
- Coughing/wheezing or respiratory distress
- Throat or chest constriction
- Difficulty swallowing
- Hypotension/shock
- Edema
- Nausea/vomiting

### Differential


- Urticarial (rash only)
- Anaphylaxis (systemic effect)
- Shock (vascular effect)
- Angioedema (drug induced)
- Aspiration/airway obstruction
- Asthma/RAD
- CHF
- Scorpion Envenomation

### Pearls


- Recommended exam: Mental status, skin, heart, lung
- Anaphylaxis is an acute and potentially lethal multisystem allergic reaction.
- EPINEPHRINE is a first-line drug that should be administered immediately in acute anaphylaxis. IM EPINEPHRINE (1:1000) should be administered before or during attempts at IV or IO access.
- Remove trigger if still present (stinger, food, etc).
- Never give EPINEPHRINE 1:1000 (IM concentration) through IV/IO route.
- IM EPINEPHRINE should be administered in the anterolateral thigh (vastus lateralis) as produces the most rapid rise in serum level.
- Contact Medical Control for refractory anaphylaxis.
- Consider monitoring ETCO<sub>2</sub>.
- Hypovolemia or distributive shock should be addressed with a fluid bolus prior to the administration of push-dose pressors.
- Contrary to common belief that all cases of anaphylaxis present with cutaneous manifestations, such as urticaria or mucocutaneous swelling, a significant portion of anaphylactic episodes may not involve these signs and symptoms on initial presentation.
- There is no proven benefit to using steroids in the management of allergic reactions and/or anaphylaxis.

### QI Metrics

- EPINEPHRINE given appropriately.
- Airway assessment documented.



# Pediatric Altered Mental Status



**History**

- Past medical history
- Medications
- Recent illness
- Irritability
- Lethargy
- Changes in feeding/sleeping
- Diabetes
- Potential ingestion
- Trauma
- Last wet diaper/urine output

**Signs and Symptoms**

- Decrease in mentation
- Change in baseline mentation
- Bizarre behavior
- Hypoglycemia (cool, diaphoretic skin)
- Hyperglycemia (warm, dry, skin; fruity breath; Kussmaul respirations; dehydration)

**Differential**

- Hypoxia
- CNS (trauma, stroke, seizure, infection)
- Thyroid (hyper/hypo)
- Shock (septic-infection, metabolic, traumatic)
- Diabetes (hyper/hypoglycemia)
- Toxicologic
- Acidosis/Alkalosis
- Environmental exposure
- Electrolyte abnormalities
- Psychiatric disorder

**Pearls**

- Recommended Exam: Mental Status, HEENT, Skin, Heart, Lung, Abdomen, Back, Extremities and Neuro.
- Pay careful attention to the head exam for signs of injury.
- With depressed mental status, initial focus is on airway protection, oxygenation, ventilation, and perfusion.
- Be aware of AMS as presenting sign of an environmental toxin or Haz-Mat exposure and protect personal safety and that of other responders.
- Consider alcohol, prescription drugs, illicit drugs and over the counter preparations as possible etiology.
- If narcotic overdose or hypoglycemia is suspected, administer NARCAN 0.1 mg/kg or GLUCOSE prior to advanced airway procedures.
- New onset DKA in pediatric patients commonly presents with nausea, vomiting, abdominal pain, and/or urinary frequency. Overly aggressive administration of fluid in hyperglycemic patients may cause cerebral edema or dangerous hyponatremia. Cerebral edema is a leading cause of death in children with DKA. Limit IV fluid bolus to single 10 ml/kg bolus.
- Dextrose 10% can be safely used in all ages of patient. Dextrose 10% works as effectively and quickly as other concentrations.

# Pediatric Bradycardia

 General Pediatric Assessment



 Ventilation Management

Bradycardia causing hypotension, altered mental status, poor perfusion or shock?

No


Yes

Identify underlying cause

<b>E</b>	Blood glucose testing
<b>E*</b> 	Consider Vascular Access
<b>P</b>	Cardiac monitor
	Consider Overdose/Poisoning

Monitor, Reassess, Transport to the hospital

Identify underlying cause



<b>E</b>	Blood glucose testing
<b>E*</b> 	Vascular Access
<b>P</b>	Cardiac monitor


HR <60 bpm?

Yes

No

<b>FR</b>	<b>CPR</b>
<b>P</b>	<b>EPINEPHRINE</b> 1:10,000 0.01 mg/kg IV/IO (max 1 mg); repeat q 3-5 min
<b>P</b>	<b>ATROPINE</b> 0.02 mg/kg IV/IO (min dose 0.1 mg; max 0.5 mg) may repeat once after 5 min

<b>E*</b>	LR or NS bolus 20 ml/kg; may repeat up to 60 ml/kg
<b>P</b>	<b>PUSH DOSE EPINEPHRINE</b> 1:100,000 10 mcg IV/IO, may repeat q 3-5 min
<b>P</b>	<b>ATROPINE</b> 0.02 mg/kg IV/IO (min dose 0.1 mg, max 0.5 mg); may repeat once after 5 min
 	Consider Transcutaneous Pacing

 Refer to Push Dose Epinephrine Guideline pg 156 for mixing instructions

 Cardiac Arrest

Pulseless

Reassess

 Notify Receiving Hospital

### History

- Respiratory insufficiency
- Past medical history
- Medications
- Pacemaker


### Signs and Symptoms

- HR <60/min with hypotension, acute altered mental status, chest pain, acute CHF, seizures, syncope, or shock secondary to bradycardia
- Respiratory distress


### Differential

- Hypoxia
- Hypothermia
- Sinus bradycardia
- Athletic
- Head injury (elevated ICP)
- Spinal cord lesion
- Overdose

### Pearls

- Pediatric pacing is by Medical Control order only. 
- Recommended Exam: Mental Status, HEENT, Heart, Lung, Neuro.
- Bradycardia causing symptoms is typically <60/minute. Rhythm should be interpreted in the context of symptoms and pharmacological treatment given only when symptomatic; otherwise, monitor and reassess.
- Identifying signs and symptoms of poor perfusion caused by bradycardia are paramount.
- Hypoxemia is a common cause of bradycardia; be sure to oxygenate the patient and provide ventilatory support as needed.

# Brief Resolved Unexplained Event (BRUE)





### History

- History of circumstances and symptoms before, during, and after the event
- Concurrent symptoms (e.g., infectious symptoms, fussy, less active, poor sleep, poor feeding)
- Prior history of BRUE
- Past medical history (e.g., prematurity, prenatal/birth complications, gastric reflux, congenital heart disease, developmental delay, airway abnormalities)
- Family history of sudden unexplained death or cardiac arrhythmia
- Social history

### Signs and Symptoms

- Signs of respiratory distress (e.g., tachypnea, grunting or other abnormal breath sounds, nasal flaring, retracting, or head bobbing)
- Skin color (pallor, cyanosis, redness)
- Altered mental status (tired, lethargic, unresponsive, or irritable)
- Trauma or neglect


### Differential

- Gastric reflux
- Swallowing dysfunction
- Nasal congestion or excessive secretions
- Periodic breathing of the newborn
- Breath-holding spell
- Seizure
- Hypoglycemia
- Abuse or neglect

### Pearls

- Recommended Exam: Mental Status, HEENT, Skin, Heart, Lung, Abdomen, Back, Extremities and Neuro.
- Definition of BRUE: an event in an infant less than 1 year old reported by a bystander as sudden, brief (less than 1 minute), unexplained, and completely resolved upon EMS arrival that includes one or more of the following:
  - Breathing change (absent, decreased, or irregular)
  - Color change (central cyanosis or pallor)
  - Marked change in muscle tone (hyper- or hypotonia)
  - Altered level of responsiveness (increased, irritability, or decreased)
- Historically called an ALTE (apparent life-threatening event).
- BRUE is a group of symptoms, not a disease process.
- If the infant is not completely well upon EMS arrival, this excludes possible BRUE event.
- Routine IVs should not be placed on all suspected BRUE patients.
- Regardless of the patient's well appearance, all infants with a history of signs or symptoms suggestive of BRUE should be transported for further evaluation. Transport to a facility with at least baseline pediatric readiness. Consider transport to facility with pediatric critical care capability for any high-risk criteria:
  - Less than 2 months of age
  - History of prematurity (less than or equal to 32 weeks gestation)
  - More than one BRUE, now or in the past
  - Event duration greater than 1 minute
  - CPR or resuscitation by caregivers or trained rescuers
- EMS clinicians play a unique and important role in obtaining an accurate history soon after the event and in observing, documenting, and reporting environmental, scene and social indicators that may point to an alternate diagnosis.

# Pediatric Burns



### History

- Type of exposure (heat, gas, chemical)
- Inhalational injury
- Time of injury
- Past medical history & medications
- Other trauma
- Loss of consciousness
- Tetanus/immunization status

### Signs and Symptoms

- Burns, pain, swelling
- Dizziness
- Loss of consciousness
- Hypotension/shock
- Airway compromise/distress
- Wheezing
- Singed facial or nasal hair
- Hoarseness or voice changes


### Differential

- Superficial (1<sup>st</sup> degree) – red and painful
- Partial Thickness (2<sup>nd</sup> degree) – blistering
- Full Thickness (3<sup>rd</sup> degree) – painless/charred or leathery skin
- Thermal
- Chemical
- Electrical
- Radiation
- Lightning

### Pearls

- Burn patients are trauma patients; evaluate for multisystem trauma.
- Assure whatever has caused the burn, is no longer contacting the injury. (Stop the burning process!)
- Recommended Exam: Mental Status, HEENT, Neck, Heart, Lungs, Abdomen, Extremities, Back, Neuro.
- Consider early intubation with patients experiencing significant inhalation injuries. Consider air ambulance for airway management needs beyond the scope of the responding ground medic.
- Potential CO exposure should be treated with 100% oxygen. Pulse oximetry may not be accurate. (For patients in which the primary event is CO inhalation, transport to a hospital equipped with a hyperbaric chamber is indicated [when reasonably accessible].)
- Circumferential burns to extremities are dangerous due to potential vascular compromise secondary to soft tissue swelling. Elevate extremity.
- Burn patients are prone to hypothermia - never apply ice or cool burns; must maintain normal body temperature.
- Consider ETCO<sub>2</sub> monitoring.
- Evaluate the possibility of child abuse with children and burn injuries

Note: Each arm totals 9 percent (front of arm 4½ percent, back of arm 4½ percent)



### Patients meeting the following Criteria shall be transported to the closest appropriate Burn Care Center:

1. Second degree burns >10% body surface area (BSA).
2. Any Third degree burns.
3. Burns that involve the face, hands, feet, genitalia, perineum, or major joints.
4. Electrical burns including lightning injury.
5. Chemical burns.
6. Circumferential burns.
7. Inhalation burns.
8. Burn injury with concomitant trauma

### Early Intubation Indications

- Signs of Airway Obstruction
- Hoarseness, Stridor, Dysphagia
- Extensive Deep Facial Burns
- Significant Risk of Edema
- Burns in Mouth
- Total BSA ≥ 40%
- Signs of Respiratory Compromise
  - Accessory Muscle Use
  - Inability to Clear Secretions
  - Poor Oxygenation
- Altered Mentation

### Pearls (Electrical)

- Do not contact the patient until you are certain the source of the electric shock has been disconnected.
- Attempt to locate contact points, (entry wound where the AC source contacted the patient; an exit at the ground point); both sites will generally be full thickness.
- Cardiac monitor; anticipate ventricular or atrial irregularity to include V-Tach, V-Fib, heart blocks, etc.
- Attempt to identify the nature of the electrical source (AC vs DC), the amount of voltage and the amperage the patient may have been exposed to during the electrical shock.

### Pearls (Chemical)


- Certainly 0.9% NaCl Sol'n or Sterile Water is preferred; however if it is not readily available, do not delay; use tap water for flushing the affected area or other immediate water sources. Flush the area as soon as possible with the cleanest, readily available water or saline solution using copious amounts of fluids.

**Burn Injury IV Fluid Rates**  
**Infusion Rate < 30 KG**  
**Parkland Formula**

\*Fluid of choice LR/NS, DO NOT use dextrose containing fluids

Wt (lbs)	Wt (kg)	% TBSA	/Hr for 1 <sup>st</sup> 8 Hrs of care	60 gtt set, gtt/min	20 gtt set, gtt/min	15 gtt set, gtt/min	10 gtt set, gtt/min
11	5	10	12.5	12.5	4.2	3.2	2.1
11	5	20	25	25	8.3	6.3	4.2
11	5	30	37.5	37.5	12.5	9.5	6.3
11	5	40	50	50	16.7	12.5	8.3
11	5	50	62.5	62.5	20.8	15.7	10.5
11	5	60	75	75	25	18.7	12.5
22	10	10	25	25	8.4	6.4	4.1
22	10	20	50	50	16.6	12.5	8.4
22	10	30	75	75	25	18.9	12.5
22	10	40	100	100	33.3	25	16.6
22	10	50	125	125	41.6	31.4	20.9
22	10	60	150	150	50	37.4	25
27.5	12.5	10	31.3	31.3	10.5	7.5	5.2
27.5	12.5	20	62.5	62.5	20.8	15.7	10.5
27.5	12.5	30	93.8	93.8	31.3	23.6	15.7
27.5	12.5	40	125	125	41.7	31.7	21
27.5	12.5	50	156.2	156.2	52.1	39.8	26.3
27.5	12.5	60	187.4	187.4	62.5	47.9	31.6
33	15	10	37.5	37.5	12.6	8.5	6.2
33	15	20	75	75	25	18.8	12.6
33	15	30	112.5	112.5	37.5	28.3	18.8
33	15	40	150	150	50	37.5	25
33	15	50	187.5	187.5	62.5	46.7	31.2
33	15	60	225	225	75	55.9	37.4
38.5	17.5	10	43.8	43.8	14.7	10.6	7.3
38.5	17.5	20	87.5	87.5	29.2	21.9	14.7
38.5	17.5	30	131.3	131.3	43.8	33	21.9
38.5	17.5	40	175	175	58.3	44.2	29.2
38.5	17.5	50	218.7	218.7	72.8	55.4	36.5
38.5	17.5	60	262.4	262.4	87.3	66.6	43.8
44	20	10	50	50	16.7	12.6	8.3
44	20	20	100	100	33.3	25	16.7
44	20	30	150	150	50	37.6	25
44	20	40	200	200	66.7	50	33.3
44	20	50	250	250	83.3	62.6	41.7
44	20	60	300	300	100	75	50
49.6	22.5	10	56.3	56.3	18.8	14.2	9.4
49.6	22.5	20	112.5	112.5	37.5	28.1	18.8
49.6	22.5	30	168.8	168.8	56.3	42.3	28.2
49.6	22.5	40	225	225	75	56.4	37.6
49.6	22.5	50	281.2	281.2	93.7	70.5	47
49.6	22.5	60	337.4	337.4	112.5	84.6	56.4
55.1	25	10	62.5	62.5	20.9	15.7	10.4
55.1	25	20	125	125	41.7	31.2	20.9
55.1	25	30	187.5	187.5	62.5	47	31.3
55.1	25	40	250	250	83.4	62.5	41.8
55.1	25	50	312.5	312.5	104.2	78	52.3
55.1	25	60	375	375	125	93.5	62.8
60.6	27.5	10	68.8	68.8	23	17.3	11.5
60.6	27.5	20	137.5	137.5	45.9	34.4	23
60.6	27.5	30	206.2	206.2	68.8	51.7	34.4
60.6	27.5	40	274.9	274.9	91.7	79.7	53.3
60.6	27.5	50	343.6	343.6	114.6	96.9	64.8
60.6	27.5	60	412.4	412.4	137.5	114.1	76.3
66	30	10	75	75	25.0	18.8	12.5
66	30	20	150	150	50.0	37.5	25.0
66	30	30	225	225	75.0	56.3	37.5
66	30	40	300	300	100.0	75.0	50.0
66	30	50	375	375	125.0	93.8	62.5
66	30	60	450	450	150.0	112.6	75.0

# Pediatric Cardiac Arrest (Non-Traumatic)





### History

- Events leading to arrest
- Estimated down time
- Past medical history
- Medications
- Existence of terminal illness

### Signs and Symptoms

- Unresponsive
- Apneic
- Pulseless

### Differential

- Medical vs. Trauma
- VF/Pulseless VT
- Asystole
- PEA
- Primary cardiac event vs. respiratory or drug overdose

### Pearls

- Respiratory failure resulting in cardiac arrest should be addressed as it is identified.
- Efforts should be directed at high quality chest compressions with limited interruptions and early defibrillation when indicated.
- Consider early IO placement if IV is difficult.
- DO NOT HYPERVENTILATE.
- Once an advanced airway is in place, compressions should be continuous with ventilations every three seconds.
- Reassess and document ETT placement using auscultation and ETCO2 capnography.
- Switch compressors every two minutes.
- Try to maintain patient modesty.
- Mechanical chest compression devices should be used if available and appropriate for patient age/size in order to provide for consistent uninterrupted chest compressions and crew safety.
- Adult paddles/pads may be used on children weighing greater than 10 kg.
- Pre-assignment of pit crew roles is recommended. When this is not possible, tasks may be assigned by order of arrival: 1<sup>st</sup>: Airway; 2<sup>nd</sup>: Compressions; 3<sup>rd</sup>: IV/IO access, medication administration; 4<sup>th</sup>: Measure, Monitor/AED placement; 5<sup>th</sup>: family liaison/history gathering.
- Pre-plan drug dosing based on weight estimations while en route and verify with a height based tape once reaching the patient.
- Proper BVM selection: <5 kg = infant BVM. 5-30 kg = pediatric BVM. >30 kg = adult BVM.


### H's & T's (reversible causes)

- Hypovolemia – Volume infusion
- Hypoxia – Oxygenation & ventilation, CPR
- Hydrogen ion (acidosis) – Ventilation, CPR
- Hyperkalemia – Calcium, Albuterol, Sodium Bicarbonate
- Hypokalemia
- Hypothermia – Warming
- Hypoglycemia – Glucose
- Tension pneumothorax – Needle decompression
- Tamponade, cardiac – Volume infusion
- Toxins – Agent specific antidote
- Thrombosis, pulmonary – Volume infusion
- Thrombosis, coronary – Emergent PCI

### ENVIRONMENTAL EXPOSURE EXCEPTION:

All victims of electrocution and lightning strikes should have resuscitative efforts begun. Any decision to determine death in the field in these cases should be made **ONLY** after consultation with online medical control.

# Pediatric Cold-Related Illness



**History**

- Age, very young and old
- Exposure to decreased temperatures, but may occur in normal temperatures
- Past medical history/medications
- Drug or alcohol use
- Infections/sepsis
- Time of exposure/wetness/wind chill

**Signs and Symptoms**

- AMS/coma
- Cold, clammy
- Shivering
- Extremity pain
- Bradycardia
- Hypotension or shock
- Decreased respirations

**Differential**

- Sepsis
- Environmental exposure
- Hypoglycemia
- Stroke
- Head injury
- Spinal cord injury
- Trauma

**Pearls**

- Recommended exam: Mental Status, Heart, Lung, Abdomen, Extremities, Neuro.
- Pulse checks for patients suffering hypothermia should be performed for up to 60 seconds.
- Handle the patient gently to prevent triggering a ventricular arrest.
- Extremes of age are more prone to cold emergencies.
- Obtain and document patient temperature.
- If temperature is unknown, treat the patient based on suspected temperature.
- Active warming includes hot packs that can be used on the armpit and groin (care should be taken not to place the packs directly on the skin) and/or Ready Heat II thermal blanket or equivalent.
- Warm saline or lactated ringers IV may be used.

**Hypothermia Categories**

- Mild 90° - 95° F (33° - 35° C)
- Moderate 82° - 90° F (28° - 32° C)
- Severe 75° - 82° F (24° - 28° C)
- Profound <75 degrees F (<24° C)

**Hypothermia Mechanisms**


- Radiation
- Convection
- Conduction
- Evaporation

**Cardiac Arrest Resuscitation Guidelines for the Hypothermic Patient**

- Contraindications for initiation of resuscitation in the hypothermic patient:
  - Obvious fatal injuries (such as decapitation)
  - The patient exhibits signs of being frozen (such as ice formation in the airway)
  - Chest wall rigidity such that compressions are impossible
  - Danger to rescuers or rescuer exhaustion
  - Avalanche victims buried for 35 minutes or longer with airway obstruction by ice or snow
- Fixed and dilated pupils, apparent rigor mortis, and dependent lividity may not be contraindication for resuscitation in the severely hypothermic patient.
- The mainstay of therapy in severe hypothermia and cardiac arrest should be effective chest compressions and attempts at rewarming.
- Defibrillation may be attempted once. Do not make further attempts at defibrillation until the core temperature has increased to greater than 30°C (86°F).
- Medications should be withheld until the patient's core temperature is greater than 30°C (86°F). Intervals between medication provision should be doubled until the patient reaches 35°C (95°F), at which time, normal medication intervals may be adopted.
- If a hypothermic patient clearly suffered cardiac arrest and subsequently became hypothermic afterward with prolonged down time between arrest and rescue, there is no rationale for initiating resuscitation and warming the patient.



# Pediatric Drowning



### History

- Submersion in liquid, regardless of depth
- Possible history of trauma (dive)
- Duration of immersion
- Temperature of water or possibility of hypothermia
- Degree of water contamination

### Signs and Symptoms

- Unresponsive
- Mental status changes
- Decreased or absent vital signs
- Vomiting
- Coughing, wheezing, rales, stridor, rhonchi
- Apnea
- Frothy/foamy sputum


### Differential

- Trauma
- Pre-existing medical condition
- Barotrauma
- Decompression illness
- Post-immersion syndrome

### Pearls

- Recommended exam: Trauma survey, Head, Neck, Chest, Abdomen, Back, Extremities, Skin, Neuro.
- Ensure scene safety.
- Resuscitation efforts should follow an **ABCs (Airway, Breathing, Circulation)** strategy for drowning victims in cardiac arrest. Initiate 5 rescue breaths followed by a ratio of 15 chest compressions to 2 breaths.
- Drowning is a common cause of death in children. Risk factors for drowning include male gender, age less than 14 years old, alcohol use, lack of supervision, and risky behavior.
- Hypothermia is often associated with submersion incidents.
- All patients should be transported for evaluation because of potential for worsening over the next several hours.
- Resuscitation recommendations based on water temperature and submersion time:
  - If water temperature is less than 43°F (6°C) and the patient is submerged with evidence of cardiac arrest: Survival is possible for submersion time less than 90 minutes and resuscitative efforts should be initiated.
  - If water temperature is greater than 43°F (6°C) and the patient is submerged with evidence of cardiac arrest: Survival is possible for submersion time less than 30 minutes and resuscitative efforts should be initiated.

# Pediatric Heat-Related Illness



**History**

- Age, very old and young
- Exposures to increased temperatures and/or humidity
- Past medical history/medications
- Time and duration of exposure
- Poor PO intake, extreme exertion
- Fatigue and/or muscle cramping

**Signs and Symptoms**

- AMS/coma
- Hot, dry, or sweaty skin
- Hypotension or shock
- Seizures
- Nausea

**Differential**

- Fever
- Dehydration
- Medications
- Hyperthyroidism
- Heat cramps, heat exhaustion, heat stroke
- CNS lesions or tumors

**Pearls**

- Recommended exam: Mental Status, Skin, Heart, Lung, Abdomen, Extremities, Neuro.
- Extremes of age are more prone to heat emergencies.
- Salicylates may elevate body temperatures.
- Sweating generally disappears as body temperatures rise over 104° F (40° C).
- Intense shivering may occur as patient is cooled. Paramedics can consider administering a low dose benzodiazepine.
- Active cooling includes application of cold packs or ice (not directly on skin), fanning, or air conditioning.
- Cold saline is not to be administered for the treatment of hyperthermia unless directed by online medical control.
- There is no evidence supporting EMS obtaining orthostatic vital signs as a clinical indicator.
- Cold water immersion is the preferred method of active cooling. Some providers such as certified athletic trainers and event medical personnel are prepared to initiate cold water immersion prior to EMS arrival. If cold water immersion was initiated due to documented hyperthermia, these patients should not be removed from cold water immersion prior to their rectal temperature reaching 102.2F (39C) or mental status returning to baseline unless it is required to manage other emergent issues such as airway.

**Heat Cramps**

- Consist of benign muscle cramping caused by dehydration and is not associated with an elevated temperature.

**Heat Exhaustion**

- Consists of dehydration, salt depletion, dizziness, fever, AMS, headache, cramping, N/V. Vital signs usually consist of tachycardia, hypotension and elevated temperature.

**Heat Stroke**

- Consists of dehydration, tachycardia, hypotension, temperature >104° F (40° C), and AMS. Potential for seizures or coma.

**Heat Syncope**

- Transient loss of consciousness attributed to heat exposure.

# Pediatric Head Injury

## EPIC4KIDS ALGORITHM



### EPIC4Kids Algorithm: For Children Any Suspicion of TBI (Mechanism, GCS, Exam)

Automatically: 15 L/min O<sub>2</sub> NRB, IV Access  
q3-5 min: Monitor O<sub>2</sub>, BP, HR

#### Airway/Breathing

O<sub>2</sub> sat <90 &/or hypoventilation  
(despite NRB)

No

Continue careful  
monitoring of O<sub>2</sub>  
sat and airway

Yes

-BLS airway  
maneuvers  
-BVM \*

O<sub>2</sub> sat <90  
despite BLS?

Yes

Consider ALS airway if experienced provider available:

- Place advanced airway:
  - Pre-oxygenate: BVM with 100% O<sub>2</sub> @ age-appropriate rate \*
  - Check placement using ETCO<sub>2</sub> monitor/detector
- AVOID even MILD hyperventilation with Ventilation Rate Timer and Pressure-controlled Bag:
  - Carefully keep rate @ age-appropriate rate \*
  - ETCO<sub>2</sub> available: Target ETCO<sub>2</sub> 40 mmHg (range: 35-45)
  - Control Ventilatory volume:
    - Ventilator available: utilize as soon as possible
    - Tidal volume = 7 cc/kg
    - Ventilator not available: Continue Pressure-controlled BVM
- Monitor: O<sub>2</sub> sat and airway every 3-5 minutes:
  - If O<sub>2</sub> sat <90% despite above interventions, consider:
    - Tension pneumothorax & needle thoracostomy

**NOTE:** NO ONE (not even Respiratory Therapists) can manually ventilate at the proper rate without ventilatory adjuncts (EVERYONE inadvertently hyperventilates unless meticulously preventing it):

- Ventilation Rate timers
- Pressure-controlled bags
- ETCO<sub>2</sub> monitoring with someone watching the level continuously
- Mechanical ventilator with careful ETCO<sub>2</sub> monitoring

#### Circulation

Hypotension\*\* or other signs of shock

No

-Continue careful  
Monitoring BP/HR  
-Pay attention for  
early signs of shock:  
-Tachycardia  
-Dropping SBP

Yes

-20ml/kg bolus NS  
-Repeat until hypotension  
resolves

#### Disability

Evaluate  
Mental  
Status/GCS


#### \* Ventilation Rates:

- Infants: (0-24 mo.):  
-25 breaths/min (bpm)
- Children: (2-14 yrs):  
-20 bpm
- Adolescents: (15-17 yrs):  
-10 bpm (same as adults)

#### \*\* Identifying Hypotension in children

- 0-9 yrs: 70 + (age x 2)
- ≥10 yrs: <90mmHg
- Rules of thumb:  
Newborn: <70mmHg  
5 year old: <80mmHg  
10 and older: <90mmHg →  
(Same as adults)

# Neonatal Resuscitation



### History

- Due date
- Time contractions started/duration/frequency
- Rupture of membranes (meconium)
- Time and amount of any vaginal bleeding
- Sensation of fetal movement
- Prenatal care
- Past medical and delivery history
- Medications
- Gravida/Para Status
- High risk pregnancy

### Signs and Symptoms

- Respiratory rate and effort
- Signs of respiratory distress
- Heart rate
- Muscle tone
- Color, appearance
- APGAR score

### Differential

- Abnormal presentation (breech, limb)
- Prolapsed cord
- Placenta previa
- Abruptio placenta

### Pearls

- Recommended Exam: Mental Status, Skin, HEENT, Neck, Chest, Heart, Lungs, Abdomen, Neuro.
- Document all times (delivery, contraction, duration, frequency).
- Some bleeding is normal; copious amounts of blood or free bleeding is abnormal.
- Record APGAR at one and five minutes after birth.
- APGAR of 7-10 is normal, while 4-7 require resuscitative measures.
- Pulse oximetry should be considered if resuscitative efforts are initiated or if supplemental oxygen is administered. Place pulse oximeter probe on right hand (preductal).
- If immediate resuscitation is required and the newborn is still attached to the mother, clamp the cord in two places and cut between the clamps. If no resuscitation is required, warm/dry/stimulate the newborn, and then cut/clamp the cord after 60 seconds or the cord stops pulsating.
- Most newborns require only drying, warming, and stimulating to help them transition from fetal respiration to newborn respiration. The resuscitation sequence can be remembered as **Dry, Warm, and Stimulate – Ventilate – Evaluate – and Resuscitate.**
- Transport mother and infant together whenever possible.

		INTERVENTION INDICATED		
		Blow-by Oxygen	Bag-Mask-Ventilation (BVM)	BVM and Chest compressions
ASSESSMENT	Heart Rate (BPM)	> 100	60–100	< 60
	Respiratory Distress/Apnea	No	Yes	
	Central Cyanosis Present	Yes	Yes/No	

### APGAR


	<u>Score=0</u>	<u>Score=1</u>	<u>Score=2</u>
• <u>Activity/Muscle Tone</u>	Absent	Arms/legs flexed	Active Movement
• <u>Pulse</u>	Absent	Below 100	Above 100
• <u>Grimace/Reflex Irritability</u>	No response	Grimace	Sneeze, cough, pulls away
• <u>Appearance/Skin Color</u>	Blue-Grey, pale all over	Normal, except extremities	Normal over entire body
• <u>Respiration</u>	Absent	Slow, irregular	Good, crying

### Caveats:

- Deep airway suctioning no longer recommended.
- Traditional CPR 3:1 ratio is standard for newborns.
- Most newborns requiring resuscitation will respond to BVM, compressions and Epinephrine; for those that don't, consider hypovolemia, pneumothorax, and/or hypoglycemia (BG <40).



# Pediatric Overdose / Poisoning





**History**

- Ingestion or suspected ingestion of a potentially toxic agent
- Substance ingested, route, quantity
- Time of ingestion
- Reason (suicidal, accidental, criminal)
- Available medications in home
- Past medical history, medications

**Signs and Symptoms**

- Mental status changes
- Hypotension/hypertension
- Decreased respiratory rate
- Tachycardia, dysrhythmias
- Seizures
- SLUDGE+BBB
- Malaise, weakness
- GI symptoms
- Dizziness
- Syncope
- Chest pain

**Differential**

- TCA overdose
- Acetaminophen
- Aspirin
- Depressants
- Stimulants
- Anticholinergic
- Cardiac medications
- Solvents, alcohols, cleaning agents, insecticides


**Pearls**

- Recommended exam: Mental Status, Skin, HEENT, Heart, Lung, Abdomen, Extremities, Neuro.
- Pediatric patients should be evaluated by a physician if an overdose/poisoning is suspected regardless of agent, amount or time.
- 4.2% SODIUM BICARBONATE should be used for all neonatal patients.
- NARCAN should be administered in small increment doses IV to address respiratory depression and ensure adequate ventilation. Monitor patient to watch for any signs of respiratory depression reoccurring. IV/IM are preferred routes for predictability.
- Overdose or toxin patients with significant ingestion/exposure should be closely monitored and aggressively treated. Do not hesitate to contact medical control if needed.
- In the case of cyanide poisoning, altered mental status may be profound. Profound altered mental status can be defined as a deficit that includes disorientation, bewilderment and difficulty following commands.
- Poison Control: 1-800-222-1222

**Agents**

- Acetaminophen: Initially normal or N/V. Tachypnea and AMS may occur later. Renal dysfunction, liver failure and/or cerebral edema may manifest.
- Depressants: Decreased HR, BP, temp and RR.
- Anticholinergic: Increased HR, increased temp, dilated pupils and mental status changes.
- Insecticides: May include S/S of organophosphate poisoning.
- Solvents: N/V, cough, AMS.
- Stimulants: Increased HR, BP, temp, dilated pupils, seizures and possible violence.
- TCA: Decreased mental status, dysrhythmias, seizures, hypotension, coma, death.

# Pediatric Pain Management



### History

- Age
- Location, duration
- Severity (1-10)
- Past medical history
- Pregnancy status
- Drug allergies and medications

### Signs and Symptoms

- Severity (pain scale)
- Quality
- Radiation
- Relation to movement, respiration
- Increased with palpation of area

### Differential

- Musculoskeletal
- Visceral (abdominal)
- Cardiac
- Pleural, respiratory
- Neurogenic
- Renal (colic)

### Pearls

- Recommended exam: Respiratory Status, Mental Status, Area of pain, Neuro.
- Pain severity (1-10) is to be recorded before and after medication administration and upon arrival at the hospital. Use FLACC for < 4 yo. Use Wong-Baker FACES for 4-12 yo.
- Monitor BP and respirations closely as sedative and pain control agents may cause hypotension and or respiratory depression.
- Consider patient's age, weight, clinical condition, use of drugs/alcohol, exposure to opiates when determining initial opiate dosing. Weight based dosing may provide a standard means of dosing calculation but it does not predict response. Consider starting at a lower initial dose and titrating to effect is recommended.
- Intranasal routes of opioid analgesia are preferred as the initial dosing route in pediatrics where IV access may be problematic.
- Exercise caution when administering opiates and benzodiazepines; this combination results in deeper anesthesia with significant risk of respiratory compromise.
- Burn patients may require more aggressive dosing.
- Consider FENTANYL as the primary opioid medication for pediatric traumatic pain.
- ACETAMINOPHEN may be given if it has been  $\geq 4$  hours since last dose. IBUPROFEN may be given if it has been  $\geq 6$  hours since last dose and age > 6 months. KETOROLAC may be given if it has been  $\geq 6$  hours since last NSAID dose and age > 2 years old.

### QI Metrics

- Vital signs with O<sub>2</sub> sats documented.
- Pain scale documented before and after each intervention.
- Repeat vital signs after each intervention.
- If considering repeat administration of pain medications, nasal cannula capnography must be utilized.

### FLACC Behavioral Scale


Categories	Scoring		
	0	1	2
Face	No particular expression or smile	Occasional grimace or frown, withdrawn, disinterested	Frequent to constant frown, clenched jaw, quivering chin
Legs	Normal position or relaxed	Uneasy, restless, tense	Kicking, or legs drawn up
Activity	Lying quietly, normal position, moves easily	Squirming, shifting back and forth, tense	Arched, rigid, or jerking
Cry	No cry (awake or asleep)	Moans or whimpers, occasional complaint	Crying steadily, screams or sobs, frequent complaints
Consolability	Content, relaxed	Reassured by occasional touching, hugging, or being talked to, distractable	Difficult to console or comfort

Each of the five categories (F) Face; (L) Legs; (A) Activity; (C) Cry; (C) Consolability is scored from 0-2, which results in a total score between zero and ten.

### Wong-Baker FACES® Pain Rating Scale



# Pediatric Respiratory Distress



**History**

- Asthma, RAD
- Home treatment (oxygen, nebulizers)
- Medication
- Prematurity
- Family history of asthma, eczema, or allergies
- Change in feeding patterns and/or number of wet diapers
- Sick contacts
- Toxic exposure

**Signs and Symptoms**

- Change in mental status
- Decreased ability to speak, weak cry
- Increased respiratory rate and effort
- Wheezing, rhonchi
- Use of accessory muscles, retractions, grunting, nasal flaring
- Fever, nasal congestion, cough
- Tachycardia

**Differential**

- Bronchiolitis
- Croup
- Asthma, RAD
- Epiglottitis
- Anaphylaxis
- Aspiration, foreign body
- Pneumonia
- Pneumothorax
- Hyperventilation
- Inhaled toxin

**Pearls**

- Recommended exam: Mental Status, HEENT, Skin, Neck, Heart, Lungs, Abdomen, Extremities, Neuro.
- Be prepared to assist ventilations as needed.
- Pulse oximetry and end tidal continuous waveform capnography should be monitored.
- Allow the patient to assume a position of comfort.


**Bronchiolitis**

- Typically presents with diffuse rhonchi and/or wheezing with a viral or other undifferentiated illness characterized by fever, rhinorrhea, nasal congestion, cough, tachypnea, and/or respiratory distress.
- Suctioning can be a very effective intervention to alleviate distress since infants are obligate nose breathers.
- Insufficient data exist to recommend the use of inhaled steam or nebulized saline
- Although albuterol and steroids have previously been a consideration, the most recent evidence does not demonstrate a benefit in routine use of albuterol or steroids for bronchiolitis.
- Ipratropium and other anticholinergic agents should not be given to children with bronchiolitis.
- Although nebulized hypertonic saline has been shown to decrease hospital length of stay when used for bronchiolitis, it does not provide immediate relief of distress and should not be administered to children in respiratory distress in the prehospital setting.

**Croup**

- Typically presents with barking cough and/or stridor with a viral or other undifferentiated illness characterized by fever, rhinorrhea, nasal congestion, tachypnea, and/or respiratory distress.
- Symptoms often improve with being upright and breathing cold night air.
- Foreign bodies can mimic croup, it is important to ask about a possible choking event.

# Pediatric Seizure



**History**

- Reported or witnessed seizure activity
- Previous seizure history
- Seizure medications
- History of trauma
- History of diabetes
- Time of seizure onset
- Number of seizures
- Alcohol ingestion
- Fever

**Signs and Symptoms**

- Decreased mental status
- Sleepiness
- Oral trauma
- Incontinence
- Observed seizure activity
- Evidence of trauma
- Unconsciousness

**Differential**

- CNS trauma
- Tumor
- Metabolic, hepatic or renal failure
- Hypoxia
- Electrolyte abnormality (Na, Ca, Mg)
- Drugs, medications non-compliance
- Infection
- Febrile seizure
- Hyperthermia
- Hypothermia
- Nonaccidental trauma (abuse)

**Pearls**

- Recommended exam: Mental Status, HEENT, Heart, Lungs, Extremities, Neuro.
- Benzodiazepines are effective in terminating seizures; do not delay IM/IN administration while initiating an IV.
- Status epilepticus is defined as two or more seizures successively without an intervening lucid period, or a seizure lasting over five minutes.
- Grand mal seizures (generalized) are associated with loss of consciousness, incontinence and oral trauma.
- Focal seizures affect only part of the body and are not usually associated with a loss of consciousness.
- Be prepared to address airway issues and support ventilations as needed.
- Consider cardiac and ETCO<sub>2</sub> monitoring.


**Febrile Seizures**


- Seizure activity in the setting of a febrile illness occurring between the age of 6 months and under 6 years old.
- The following interventions provide symptomatic relief for fevers, but do not stop the seizure:
  - ACETAMINOPHEN and/or
  - IBUPROFEN/KETOROLAC and/or
  - Removing excessive layers of clothing and/or
  - Applying cool compresses to the body
- ACETAMINOPHEN may be given if it has been  $\geq 4$  hours since last dose. IBUPROFEN may be given if it has been  $\geq 6$  hours since last dose and age  $> 6$  months. KETOROLAC may be given if it has been  $\geq 6$  hours since last NSAID dose and age  $> 2$  years old.

# Pediatric Shock

## General Pediatric Assessment

Estimated Minimum  
Systolic  
BP Calculation  
(Age in Years x 2) + 70

 **BP calculation  
applies up to age  
10 years**


FR	Oxygen 15 L NRB
E*	 Vascular Access
P	Cardiac monitor/Capnography

 Alternative appropriate treatment  
protocols as indicated

Trauma -  
related

Non-Trauma  
related

 General Pediatric Trauma

 **For patients with  
known adrenal insufficiency,  
administer patient's own  
HYDROCORTISONE (SOLU-  
CORTEF) as prescribed OR;  
METHYLPREDNISOLONE 2  
mg/kg IV/IO/IM, max 125 mg**

Continue General Pediatric  
Assessment

**E\*** LR or NS bolus  
20 ml/kg IV/IO;  
may repeat x 2 with no rales on  
lung exam

BG <60 mg/dl  
BG <40 mg/dl  
in newborn

**E**

Blood glucose testing

BG >250 mg/dl

BG normal

**E** ORAL GLUCOSE if patient  
protecting airway


**A** D10, 5-10 ml/kg IV/IO;  
max single dose 25 g

**E\*** GLUCAGON  
0.5 mg IM/IN (<20 kg);  
1 mg IM/IN (>20 kg);  
for no IV access

**E\*** NS bolus  
10 ml/kg for hypotension;  
may repeat up to 40 ml/kg

**P** NOREPINEPHRINE 0.05-0.5 mcg/kg/  
min IV drip titrate to  
SBP >70 mm Hg + 2 x Age

**P** PUSH DOSE EPINEPHRINE 1:100,000  
10 mcg IV/IO, may repeat q 3-5 min  
titrate to SBP >70 mm Hg + 2 x Age  
(1 ml of a 1:100,000 solution)  
AND/OR EPINEPHRINE 0.05-0.3 mcg/  
kg/min IV drip

 Refer to Push Dose  
Epinephrine Guideline pg 156  
for mixing instructions



**History**

- Blood loss
- Fluid loss - vomiting, diarrhea, fever
- Infection
- Medications
- Allergic reaction
- History of poor oral intake

**Signs and Symptoms**

- Restlessness, confusion
- Weakness, dizziness
- Weak rapid pulse
- Pale, cool, clammy skin
- Poor skin turgor
- Delayed capillary refill
- Dry mucous membranes
- Hypotension
- Coffee-ground emesis
- Tarry stools

**Differential**

- Hypovolemic shock
- Cardiogenic shock
- Septic shock
- Neurogenic shock
- Anaphylactic shock
- Dysrhythmias
- Pulmonary embolism
- Tension pneumothorax
- Medication effect or overdose
- Vasovagal

**For patients with known adrenal insufficiency, administer patient's own HYDROCORTISONE (SOLU-CORTEF) as prescribed OR; METHYLPREDNISOLONE 2 mg/kg IV/IO/IM, max 125 mg**

**Causes of Adrenal Insufficiency:**

**Addison's Disease**

**Congenital Adrenal Hyperplasia**

**Long term administration of steroids**

**Others**

**• Pearls**

- Recommended exam: Mental Status, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro.
- Hypotension can be defined as a systolic < Estimated Minimum Systolic. This is not always reliable and should be interpreted in context and patient's typical BP, if known. Shock may present with a normal BP initially.
- Hypotension for age (lowest acceptable systolic blood pressure in mmHg):
  - Less than 1 years of age: 60
  - 1–10 years old: (age in years) (2) + 70
  - Greater than 10 years old: 90
- Shock often is present with normal vital signs and may develop insidiously. Tachycardia can be a late sign of shock in children and a tachycardic child may be close to cardiovascular collapse.
- Consider all possible causes of shock and treat per appropriate protocol.
- Physiologic targets of resuscitation: Systolic blood pressure at least fifth percentile for age, strong distal pulses, warm skin perfusion, capillary refill less than 2 seconds and improving mental status.

New onset DKA in pediatric patients commonly presents with nausea, vomiting, abdominal pain, and/or urinary frequency. Overly aggressive administration of fluid in hyperglycemic patients may cause cerebral edema or dangerous hyponatremia. Cerebral edema is a leading cause of death in children with DKA.

**Hypovolemic shock**

- Hemorrhage, trauma, GI bleeding, ruptured aortic aneurysm, or pregnancy related bleeding

**Cardiogenic shock**

- Heart failure, MI, cardiomyopathy, myocardial contusion, toxins, dysrhythmia, valvular disease


**Distributive shock**

- Sepsis, anaphylaxis, neurogenic, toxins, overdose, endocrine

**Obstructive shock**

- Pericardial tamponade, pulmonary embolus, tension pneumothorax

# Pediatric Smoke Inhalation



### History

- Exposed to smoke in a structure fire
- Exposed to smoke in a vehicle fire
- Exposed to smoke from other sources, industrial, confined space, wilderness fire etc.

### Signs and Symptoms

- Facial burns
- Singed nasal hairs or facial hair
- Shortness of breath
- Facial edema
- Stridor
- Grunting respirations

### Differential

- COPD
- CHF
- Toxic inhalation injury
- Caustic inhalation injury
- Pneumonitis

### Pearls

- Protect yourself and your crew.
- Have a high index of suspicion when treating patients at the scene of a fire.
- If the medication is not available on scene do not delay transport waiting for it.
- Carefully monitor respiratory effort and correct life threats immediately.
- Decide early on if you want to intubate as burned airways swell making intubation difficult.
- Profound altered mental status can be defined as a deficit that includes disorientation, bewilderment and difficulty following commands.

## Pediatric Cyanokit Instructions and Dosing

1. Reconstitute Cyanokit vial per the Instructions. **The new vial concentration will be 25mg/ml.**
2. See chart below to find the appropriate reconstituted dose.


APPROXIMATE AGE	NB-1 m	2 m	4-6 m	8-10 m	1-1.5 y	2-2.5 y	3-3.5 y	4-5 y	5.5-7 y	7.5-8 y	8.5-10 y
WEIGHT IN LBS	7-9 lb	11 lb	13-15 lb	18-20 lb	22-24 lb	26-31 lb	33-40 lb	42-48 lb	53-62 lb	66-70 lb	71-80 lb
WEIGHT IN KGS	3-4 kg	5 kg	6-7 kg	8-9 kg	10-11 kg	12-14 kg	15-18 kg	19-22 kg	24-28 kg	30-32kg	34-36 kg
RECONSTITUTED SYRINGE AMOUNT	10 ml	15 ml	20 ml	25 ml	30 ml	40 ml	50 ml	60 ml	75 ml	85 ml	100 ml
TOTAL DOSAGE OF MEDICATION	250mg	375mg	500mg	625mg	750mg	1000mg	1250mg	1500mg	1875mg	2125mg	2500mg
gtts / sec	1	1	1	5	5	5	5	5	5	6	6

GIVE DOSE IN 50ml NS BAG

GIVE DOSE IN 250ml NS BAG

3. Withdraw and waste the equivalent volume of Normal Saline from the bag size indicated.
4. Draw the appropriate dose from the vial referencing the chart, using the appropriate syringe size.
5. Inject the reconstituted medication into the appropriate sized bag of Normal Saline, below the Chart.
6. Spike the bag with 15 gtts/ml IV tubing.
7. Piggyback line into an IV/IO line and **Infuse over ~15 minutes** using the gtts/second noted above.

# Pediatric Tachycardia / Stable (Normal Mental Status, Palpable Radial Pulse)



**History**

- Medications
- Diet (caffeine)
- Drugs (cocaine, methamphetamines)
- Past medical history
- Syncope/near syncope
- History of palpitations/racing heart


**Signs and Symptoms**

- Heart rate  $\geq 180$  in children
- Heart rate  $\geq 220$  in infants
- Dizziness, CP, SOB
- Diaphoresis


**Differential**

- Heart disease (WPW, valvular)
- Sick sinus syndrome
- Electrolyte imbalance
- Exertion, fever, pain, emotional stress
- Hypoxia
- Hypovolemia
- Drug effect, overdose
- Hyperthyroidism

**Pearls**

- Pediatric synchronized cardioversion is by Medical Control order only. 
- Recommended exam: Mental Status, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro.
- Carefully monitor patients as you treat them; stable tachycardias may convert to unstable rhythms/conditions quickly.
- Sedate patients prior to cardioversion, if time allows.
- Administer ADENOSINE at a proximal IV site, rapidly followed by a saline flush.
- The most common tachyarrhythmia in children is sinus.

# Pediatric Tachycardia / Unstable (Mental Status Changes, No Palpable Radial Pulse)



**History**

- Medications
- Diet (caffeine)
- Drugs (cocaine, methamphetamines)
- Past medical history
- Syncope/near syncope
- History of palpitations/racing heart


**Signs and Symptoms**

- Cardiac Arrest
- Heart rate  $\geq 180$  in children
- Heart rate  $\geq 220$  in infants
- Dizziness, CP, SOB
- Diaphoresis

**Differential**



- Heart disease (WPW, valvular)
- Sick sinus syndrome
- Electrolyte imbalance
- Exertion, fever, pain, emotional stress
- Hypoxia
- Hypovolemia
- Drug effect, overdose
- Hyperthyroidism

**Pearls**

- Pediatric synchronized cardioversion is by Medical Control order only. 
- Recommended exam: Mental Status, Skin, Heart, Lungs, Abdomen, Back, Extremities, Neuro.
- If patient is in arrest, efforts should focus on quality chest compressions and rhythm correction.
- Administer ADENOSINE at a proximal IV site, rapidly followed by a saline flush.

# Pediatric Ventilation Management


Use supplemental oxygen to maintain an oxygen saturation of >94%; >90% for patients on home oxygen for chronic conditions



Intervention effective?

No

Yes



Consider Post-Intubation Sedation: Administer  
MIDAZOLAM 0.1 mg/kg IM/IV/IO, max dose 5 mg. May repeat after 5 min OR;  
KETAMINE 2 mg/kg IV, max 100 mg. May repeat after 10 min.

Consider Post-Intubation Analgesia: Administer  
FENTANYL 1 mcg/kg; max dose 50 mcg. May repeat after 10 min.

Supraglottic/ETT placement successful?

YES

NO

Able to ventilate without supraglottic/ETT in place?

YES

NO

P Needle Cricothyrotomy

Continue General Pediatric Assessment



Always weigh the risks and benefits of endotracheal intubation in the field against transport. All prehospital endotracheal intubations are considered high risk. If ventilation/oxygenation is adequate, transport may be the best option. The most important airway device and the most difficult to use correctly and effectively is the Bag Valve Mask (not the laryngoscope). Few prehospital airway emergencies cannot be temporized or managed with proper BVM techniques.

#### Pearls

- **The Montana Board of Medical Examiners DOES NOT allow drug assisted intubation (DAI) or rapid sequence intubation (RSI) for standard ground ALS paramedics.**
- Consider preoxygenation/lung denitrogenation with a non-rebreather or a nasal cannula at 15 LPM prior to intubation (as patient condition allows).
- Severe hypotension ( $SBP < 70 + (2 \times \text{age})$ ) should be addressed with IV fluids and/or pressors (as appropriate) prior to intubation in order to reduce likelihood of post-intubation cardiovascular decline.
- Capnometry (color) or capnography (waveform) is mandatory with all methods of intubation. Document results.
- Continuous capnography (ETCO<sub>2</sub>) is mandatory for the monitoring of all patients with an advanced airway.
- If an effective airway is being maintained by BVM and/or basic airway adjuncts (e.g. nasopharyngeal airway) with continuous pulse oximetry values of  $\geq 90\%$ , or values expected based on pathophysiologic condition with otherwise reassuring vital signs (e.g. pulse oximetry of 85% with otherwise normal vitals in a post drowning patient), it is acceptable to continue with basic airway measures instead of using a supraglottic airway device or intubation.
- For the purposes of this protocol, a secure airway is achieved when the patient is receiving appropriate oxygenation and ventilation.
- An Intubation Attempt is defined as passing the laryngoscope blade or endotracheal tube past the teeth or inserted into the nasal passage.
- An appropriate ventilatory rate is one that maintains an ETCO<sub>2</sub> of 35 - 45. Avoid hyperventilation.
- Paramedics should use a supraglottic airway device if oral-tracheal intubation is unsuccessful.
- Maintain C-spine stabilization for patients with suspected spinal injury.
- Gastric tube placement should be considered in all intubated patients if time allows.
- It is important to secure the endotracheal tube well.

#### QI Metrics

- Mandatory notification of Medical Director within 24 hours of any failed intubation attempt. For simplicity, may use this online form: <https://forms.gle/G3RPJQ8fEugRafMTA>
- Mandatory notification of Medical Director within 24 hours of any attempted, successful, and/or failed cricothyroidotomy in the field. For simplicity, may use this online form: <https://forms.gle/G3RPJQ8fEugRafMTA>

# **OPERATIONS GUIDELINES**

# Air Medical Activation

## INDICATIONS:

- The decision for mode of transport from both field and inter-facility transfer patients is based on the premise that the time to definitive care and quality of care are critical to achieving optimal outcomes.
- Factors of distance, injury severity, road conditions, weather, and traffic patterns must be considered when choosing between air or ground transport. The skill level of the transport team must also be considered.
- The potential benefit to the patient should outweigh the risks associated with air transport.

In general, use of air ambulances should be considered in the following situations:


**Skill-based reasons:** The skill set or medical equipment of the critical care air medical team is required for the patient's condition. Example: Expected need for rapid sequence intubation in a patient with airway difficulties.

**Aircraft-based reasons:** The specific qualities of aircraft transport, such as speed of transport over large distances, are expected to benefit the patient. The National Association of State EMS Officials' National Model of EMS Guidelines (2019 Version 2.2, pg. 11) recommends considering "air medical transport, if available, for patients with time-critical conditions where ground transport time exceeds 45 minutes." See map below for Gallatin Valley highlighting 45 minute drive time to BHDRMC.

**Provider discretion:** Local providers are often aware of resource limitations and/or challenges specific to certain incidents or locations. Use of air ambulance resources may be considered per incident command discretion based on these circumstances.

## CANCELING AIR MEDICAL TRANSPORT

- When air medical transport (AMT) has been requested, consideration for canceling it should only be made by trained EMS providers who are on scene and able to evaluate the situation and patient care needs.
- Resources are often dispatched before the full details of a situation are known. If AMT arrives on scene, it is appropriate to make a decision to transport the patient by ground EMS if it is felt by all providers that the patient does not require the higher level services of the AMT team.



# BHDRMC Trauma Activation

## Trauma Activation Criteria

### LEVEL 1

High likelihood of needing immediate procedures/MTP/OR  
(Need services in <30mins)

- Traumatic Cardiac Arrest with or without ROSC
- Intubated/compromised airway
- Assisted Ventilations
- Respiratory
  - RR <10 or >30
  - 0-12mo. RR <20
  - Concern for new hypoxia
    - SpO<sub>2</sub> <90% on room air in the setting of trauma
- Chest Injury
  - Flail chest/ chest deformity
  - Signs of tension pneumothorax
- Shock/Hypotension/Hemodynamic instability
  - >64 y.o. w/ SBP <110,
  - 10-64 y.o. w/ SBP <90
  - 0-9 y.o. w/ SBP < 70+(2 x age in years)
    - capillary refill time > 3 sec., mottling, weak pulses
- Shock index >1 (HR/SBP)
- GCS less than or equal to 8 with traumatic mechanism
- Active bleeding requiring tourniquet proximal to wrist/ankle
- Amputation proximal to wrist/ankle
- Pulseless/mangled extremity
- Suspected spinal cord injury with paralysis/motor function loss
- GSW/ penetrating trauma to head, neck, chest, abdomen, extremities proximal to elbow/knee
- Receipt of transfer patients from another hospital who require ongoing blood transfusion and/or ongoing respiratory compromise
- Physician Discretion

### LEVEL 2

High likelihood of serious injury but less likely to need time sensitive intervention  
(Needs services in < 60 mins)

- Suspected fractures of two or more long bones
- Burns >25% TBSA or signs of inhalation injury
- Pelvic instability or significant pain with pelvic compression
- Blunt abdominal trauma with concerning exam
- Chest injury with any of the following:
  - Asymmetric or absent lung sounds
- GCS 9-12 with concerning mechanism
- Open or depressed skull fracture(s)

### LEVEL 3

Concerning mechanism but patient does not meet criteria for level 1 or level 2

- **MVC with High Risk of Severe Injury**
  - Rollover
  - Partial or complete ejection
  - Vehicle intrusion: >12 in at occupant site, >18 in at any other site
  - Death in the same vehicle
  - Unrestrained child in MVC
  - Vehicle telemetry data consistent with significant injury (ex. OnStar alert)
  - Need for extrication
  - High speed with significant impact
- **Other High-Risk Mechanisms / Criteria**
  - Pedestrian or bicyclist vs. auto
  - Fall >10 feet
  - Rider separation from motorcycle, ATV, horse, bike, etc.
  - Blast or explosion
  - Hanging or drowning
  - Head trauma with GCS 13-14
  - Suspected hip fracture
  - EMS provider judgment

#### **Geriatric Trauma Activation Considerations (>64 years)**

Anatomic: Proximal long bone fracture from MVC, multiple body regions injured

Mechanism: Fall from any height with head/brain injury, pedestrian struck by vehicle

Risk Factors: Anticoagulation use, significant comorbidities (cardiac disease, COPD, dialysis, immunosuppression)

Note: Comorbidities/anticoagulation alone are not automatic triggers but should prompt escalation when combined with mechanism or physiologic/anatomic findings

# Big Sky Area Trauma Field Triage

## Trauma Field Triage Job Aid

Priority 1	Priority 2	Priority 3
<b>STEP 1: Traumatic mechanism with unstable vital signs / altered mental status/ noted injuries</b>	<b>STEP 2: Traumatic mechanism with isolated injury and stable vital signs</b>	<b>STEP 3: Traumatic mechanism with stable vital signs and no noted injury / single orthopedic injury</b>
Activate helicopter and transport to the closest Level 3 or higher facility	Bypass Big Sky Medical Center —can be ground	Transport to Big Sky Medical Center
<p><u>Airway</u>: unable to maintain airway or need for vent</p> <p><u>Breathing</u>: RR&lt;10 or &gt;29, SpO2&lt;88% despite supp. O2</p> <p><u>Circulation</u>: confirmed reading at any time</p> <ul style="list-style-type: none"> <li>• Age 0-9 &lt;70mmHg + (2 x age in years)</li> <li>• Age 10-64 ≤90mmHg</li> <li>• Age 65+ &lt;110mmHg</li> </ul> <p><u>Disability</u>: GCS &lt;9</p> <p>Post-arrest with ROSC</p> <p><b>Injury Patterns</b></p> <ul style="list-style-type: none"> <li>-Skull deformity, suspected skull fracture</li> <li>-Suspected spinal injury with motor or sensory loss</li> <li>-Significant penetrating injuries to head, neck, chest, abdomen, back, groin, buttocks or extremities proximal to elbow or knee</li> <li>-Chest wall instability, deformity or suspected flail chest</li> <li>-Unstable pelvic fracture (unable to ambulate)</li> <li>-Suspected fracture of two or more proximal long bones</li> <li>-Crushed, degloved, mangled or pulseless extremity</li> <li>-Amputation proximal to wrist or ankle</li> <li>-Active bleeding requiring a tourniquet or wound packing with continuous pressure</li> <li>-Burns &gt;20% TBSA COMBINED with other injury and trauma</li> <li>-Chest wall injury with suspected pneumothorax</li> <li>-Blunt abdominal trauma with significant firmness, distention or tenderness</li> <li>-High voltage electrical injury with or without significant injury</li> </ul> <p><b>Transport to nearest Level 2 facility</b></p> <ul style="list-style-type: none"> <li>-Isolated neurological injury</li> <li>-Isolated burns &gt;20% TBSA with no associated trauma</li> </ul>	<p>-Pregnancy &gt;20 weeks with stable vital signs and without priority 1 injuries</p> <p>-Geriatric (&gt;65) with SBP &gt;110 with multiple body regions injured</p> <p><b>Injury Patterns</b></p> <ul style="list-style-type: none"> <li>-Suspected hip fracture</li> <li>-Orthopedic injury needing OR: Obvious femur fracture, obvious tib/fib fracture, open fractures, etc.</li> </ul> <p><b>Provider Judgement</b></p>	<p><b>Mechanism of Injury</b></p> <ul style="list-style-type: none"> <li>-High-risk auto crash <ul style="list-style-type: none"> <li>• Partial or complete ejection</li> <li>• Death in passenger compartment</li> <li>• Significant intrusion (including roof) <ul style="list-style-type: none"> <li>• &gt;12 inches occupant site OR</li> <li>• &gt;18 inches any site</li> </ul> </li> <li>• Child (age 0-9) unrestrained or in unsecured child safety seat</li> </ul> </li> <li>-Pedestrian/bicycle rider thrown, run over or with significant impact</li> <li>-Fall from height &gt;10ft (all ages)</li> <li>-Rider separated from transport vehicle with significant impact (motorcycle, ATV, horse, snowmobile etc.)</li> <li>-High energy dissipation/rapid deceleration incidences striking fixed object with momentum (Skier, mountain biker, etc.)</li> </ul> <p><b>Injury Patterns</b></p> <ul style="list-style-type: none"> <li>-Orthopedic injury with neurovascular compromise for relocation</li> <li>-Stable, isolated orthopedic injury; upper extremity fractures, ankle fractures, etc.</li> <li>-GCS 10-13 (suspected concussion)</li> <li>-Penetrating injury to hand or foot</li> </ul>



# BSMC Trauma Activation

## BSMC Trauma Activation Criteria

High likelihood of needing immediate procedures/RTP/Transport  
(Need services in < 30mins)

### LEVEL 1

#### Anatomic/Physiologic

##### **Neuro/Head:**

- GCS  $\leq$  13
- Suspected spinal cord injury with paralysis/motor function loss
- Open/depressed skull fractures

##### **Cardiac/Chest:**

- Traumatic Cardiac Arrest with or without ROSC including hanging/drowning
- Flail chest/signs of tension pneumothorax

##### **Respiratory:**

- Intubated/compromised airway
- Assisted ventilations
- Respiratory Rate <10 or >30

##### **Shock/Hypotension/Hemodynamic instability**

- >64 y.o. w/ SBP <110
- 10-64 y.o. w/ SBP <90
- 0-9 y.o. w/ SBP < 70+(2 x age in years)
- capillary refill time > 3 sec., mottling, weak pulses
- Shock index >1 (HR/SBP)

##### **Abdomen:**

- Blunt abdominal trauma with tenderness/distension
- GSW/ penetrating trauma to head, neck, chest, abdomen, extremities proximal to elbow/knee

##### **Pelvis:**

- Suspected pelvis fracture

##### **Extremities:**

- Active bleeding requiring tourniquet proximal to wrist/ankle
- Amputation proximal to wrist/ankle
- Pulseless/mangled extremity
- Suspected fracture of two or more proximal long bone fractures

##### **Burns:**

- Burns  $\geq$  25% TBSA or high-energy electrical injury

Mass casualty 2 or more major trauma

##### **Physician/RN Discretion**

High likelihood of serious injury but less likely to need time sensitive intervention  
(Needs services in < 60 mins)

### LEVEL 2

#### Mechanism

+ Any complaint or s/s of injury to head, neck, spine, chest, abdomen, pelvis

##### **MVC:**

- Partial/complete ejection
- Death in same vehicle
- Child 0-9 years old unrestrained or in unsecured child safety seat
- Need for extrication for entrapped patient
- Auto vs pedestrian or bicycle

##### **Head/Neuro:**

Head trauma with GCS 14 and on anticoagulants

##### **Ejection:**

- Motorcycle, ATV, Snowmobile, Horse/Bull

##### **Falls:**

- >10 feet
- >64 y.o. fall from any height (including standing) with evidence of traumatic head injury

##### **Hip:**

Suspected hip fracture

Blast/ explosion

##### **Physician/RN Discretion**

# Defining a Patient

## **FIRST & SECOND PARTY CALLS:**

All first party calls (the patient calls to summon assistance for themselves) and second party calls (someone who knows the patient or who is involved in the situation summons EMS) should generate a refusal of care (including lift assists). If the patient refuses vitals/assessment/etc., that should be documented.

## **EXAMPLES OF SECOND PARTY CALLS:**

A family member calls for a choking relative who is better by arrival.

A motorist calls for someone they have hit in an MVC.

Law enforcement calls for someone involved in an MVC.

## **THIRD PARTY CALLS:**

Third party calls are calls where the reporting party does NOT know the person(s) involved or the situation.

## **EXAMPLES OF THIRD PARTY CALLS:**

Someone passes an accident on the road and calls 911 without knowledge of the individuals involved.

Someone driving down the road calls for someone unknown to them who was lying in the grass.

If it is a third party call and the parties refuse EMS and state there is no medical problem (as in the person sleeping in the park, or multiple parties involved in an MVC), and they do not appear obviously injured, ill, or impaired, then the incident may be documented as “no patient found”. Providers are encouraged to document that they made a visual assessment of the scene and the person(s) involved in the paperwork to demonstrate due diligence.

# Destination Hospital Choice

## **GUIDELINE:**

Patients should be transported to the closest appropriate receiving hospital unless:

- In consultation with medical control it is determined that a more distant hospital is more appropriate to meet the needs of the patient;

**-OR-**

- The patient meets criteria or published EMS guidelines for transport to a specialty care center (i.e. Trauma, STEMI);

**-OR-**

- The patient requests a specific hospital,
- **AND** The patient's condition is considered stable to tolerate additional transport time without the need for more urgent stabilization before more lengthy transport;
- **AND** The EMS transport service has determined that such a transport would not unreasonably remove the unit from its primary area of response causing a decrease of 911 coverage to the local area;
- **AND** The patient has been informed that the transport to a more distant location will be more expensive and may not be covered by insurance if the added transport is not felt to be medically necessary by the insurance company.



# Interfacility Transfers

ECPs may transfer patients between medical facilities if they possess the knowledge and skills necessary to manage the needs of the patient. Consultation with the transferring physician is required to assure the potential needs of the patient are met while conducting the transfer. The ECPs scope of practice may not be expanded to meet the needs of the patient. Appropriate personnel must be obtained to assure continuity of patient care. Questions regarding whether the patient qualifies for IFT via ground ambulance at a certain level of care should be directed to a supervisor/battalion chief/assistant chief depending on the structure of your agency or the EMS Medical Director.

**EMT (Emergency Medical Technician): GENERAL GUIDANCE:** If the patient has active medications running in an IV drip, has recently been administered sedating medications, has active airway or circulation issues, or has undergone advanced procedures (beyond simple IV starts, for example), this suggests that the patient falls outside the scope of the EMT for transfer and a higher-level provider or other means of transport should be considered. Status and care required for each patient must be evaluated individually to assess if the EMT is the correct provider for interfacility transport. An EMT (without the IV/IO maintenance endorsement) may transfer a patient BLS with a saline lock left in place. No monitoring of the saline lock is expected during transport. An EMT with IV/IO maintenance endorsement may transfer a patient BLS with clear non-medicated IV fluids running (eg D5W, LR, or NS).

**AEMT (Advanced Emergency Medical Technician): GENERAL GUIDANCE:** If the patient is being ventilated, requires continuous ECG monitoring, has cardiac rhythm disturbances or active ischemia, has active medications running in an IV drip, has recently been administered sedating medications, has active airway or circulation issues, or has undergone advanced procedures, this suggests that the patient falls outside the scope of the AEMT for transfer and a higher level provider or other means of transport should be considered. Status and care required for each patient must be evaluated individually to assess if the AEMT is the correct provider for interfacility transport.

**PARAMEDIC: GENERAL GUIDANCE:** If the patient is being transferred from an ICU or inpatient unit, has multiple IV medications or infusions, or has labile condition or vital signs, this suggests that the patient falls outside the scope of the Paramedic for transfer and Paramedic(s) with the Critical Care Endorsement or other means of transport should be considered. Status and care required for each patient must be evaluated individually to assess if the Paramedic is the correct provider for interfacility transport. The Paramedic may continue administration of medications initiated in the emergency department such as antibiotics, steroids, ACLS drugs, vitamins, magnesium (including OB mag), nitroglycerin, fractionated heparin, etc. via subcutaneous, intramuscular, intraosseous, and/or intravenous routes. The Paramedic may utilize central venous lines if previously trained and authorized by the EMS Medical Director. For the transport medication the Paramedic should be familiar with pharmacology & indications as well as signs, symptoms and treatment of any major adverse drug reactions. Infusion may be discontinued if significant adverse reaction is noted and should be reported as soon as possible.

Paramedics may initiate or continue the infusion of blood products.

Paramedics may maintain and monitor patients with a chest tube in place.

**PARAMEDIC, CRITICAL CARE ENDORSEMENT: GENERAL GUIDANCE:** Transport of intubated patients maintained on a mechanical ventilator and requiring the administration of sedation/paralytic medications are limited to critical care endorsed paramedics only.

# Medication List

The following is a list of approved medications for use in Gallatin County. While other medications are listed in the NASEMSO National Model EMS Clinical Guidelines version 3.0, only the following are approved for administration in Gallatin County.

Acetaminophen  
Activated charcoal  
Adenosine  
Albuterol  
Amiodarone  
Aspirin  
Atropine  
Brilinta (BSFD, HBRFD, and YMCRFD only)  
Calcium chloride  
Calcium gluconate  
Dextrose  
Diazepam  
Diphenhydramine  
Droperidol  
Epinephrine  
Fentanyl  
Glucagon  
Haloperidol  
Heparin (BSFD, HBRFD, and YMCRFD only)  
Hydroxocobalamin  
Ibuprofen  
Ipratropium  
Ketamine  
Ketorolac  
Lidocaine  
Lorazepam  
Magnesium sulfate  
Methylprednisolone  
Metoclopramide  
Metoprolol  
Midazolam  
Morphine  
Naloxone  
Nitroglycerin  
Norepinephrine  
Ondansetron  
Oral glucose  
Oxygen  
Oxymetazoline  
Oxytocin  
Sodium bicarbonate  
Tranexamic acid

# Prehospital Death Determination



For all emergency scenes where patient needs exceed available EMS resources, initial assessment and treatment shall be in accordance with the START triage methodology.

1. Patients who appear to have expired will not be resuscitated or transported by EMS personnel if any of the following obvious signs of death are present:

- A. Body decomposition
- B. Decapitation
- C. Transection of thorax (hemicorpectomy)
- D. Incineration
- E. Other obvious fatal injuries



Contact medical control if in doubt traumatic injuries are suspected to be incompatible with life.

**OR if ALL five (5) presumptive signs of death AND AT LEAST one (1) conclusive sign of death are identified.**

The (5) presumptive signs of death that **MUST** be present are:

- 1) Unresponsiveness
- 2) Apnea
- 3) Pulselessness
- 4) Fixed, dilated pupils
- 5) For Non-Traumatic Arrests, Asystole in at least 2 (two) leads or a "No Shock Advised" prompt from an AED



Conclusive signs of death include:

- 1) Dependent lividity
- 2) Rigor mortis

If any of the findings are different than those described above, clinical death is not confirmed, and resuscitative measures should be immediately initiated or continued.

2. Once it has been determined that the patient has expired and resuscitation will not be attempted:

- A) Immediately notify the appropriate authority;
- B) **DO NOT** leave a body unattended. You may be excused once a responsible person (i.e. coroner, law enforcement, fire, etc.) is present;
- C) **DO NOT** remove any property from the body or the scene for any purpose;
- D) **NEVER** transport/move a body without permission from the coroner except for assessment or its protection.



If the body is in the public view and cannot be isolated, screened, or blocked from view, and is creating an unsafe situation with citizens/family, the body can be covered with a clean sheet or STERILE BURN SHEET obtained from the EMS vehicle.

# Prehospital Report

## **PURPOSE:**

The purpose of the prehospital report is to give the receiving hospital notification of an inbound ambulance. The report should be brief and concise. It allows the receiving hospital to properly assign a room and assemble appropriate staff and resources depending on the acuity level. A full detailed report can be given at bedside to the receiving staff.

## **GUIDELINE:**

In the case of an MCI, early notification is key so the hospital has time to assemble extra staff and resources. As soon as triage is complete, the triage officer or incident command should contact the hospital with a patient count and corresponding triage categories (e.g. 2 red, 2 yellow, and 3 green).

## **METHOD:**

The preferred method of contacting BHDRMC is via Pulsara. It is also acceptable to call on the designated EMS landline. As a backup or as primary during an MCI, communication to BHDRMC will be via the GC MED COM radio channel.

The preferred method of contacting BSMC is via Pulsara. It is also acceptable to call the general ED phone number. As a backup or as primary during an MCI, communication to BSMC will be via the GC MED COM radio channel.

## **PREHOSPITAL REPORT FORMAT:**

The expected prehospital report format is MIVT for trauma and SSVT for medical. A full set of vitals signs should be reported including blood pressure, heart rate, respiratory rate, oxygen saturation, and GCS if a trauma patient. Terms such as “within normal limits” or “stable” are not acceptable.

- Name and DOB
- Activation? (Trauma, STEMI, Stroke, Sepsis)
- ETA

### **Trauma (MIVT)**

- Mechanism
- Injuries
- Vital Signs
- Treatment

### **Medical (SSVT)**

- Signs
- Symptoms
- Vital Signs
- Treatment

# POLST / Palliative Care

POLST forms have replaced the previous program of Comfort One in Montana. However, Comfort One forms are still valid and still present in the community. They should be honored if one is presented to EMS in the course of patient care.

## POLST:

### Out-of Hospital Guideline when presented with POLST Documentation

- POLST documentation, if presented to the prehospital provider, **MUST** be followed.
- POLST documentation **MUST** accompany the patient and be presented to other health care providers who subsequently attend the patient.
- The prehospital patient care report (PCR) must include the POLST documentation and care provided based on the POLST documentation.
- Never delay patient care to determine if the patient has POLST documentation.
- Comfort One bracelet identifies a patient who has a POLST document and a DNR (section A).
- A POLST document can be disregarded if the **PATIENT** requests or if the terminal condition no longer exists.
- A verbal DNR order from a physician **MUST** be followed.
- If there is a question regarding POLST, contact online medical control.

## End-of-Life Care/Palliative Care:

**PATIENT CARE GOALS:** When providing care for a patient near end-of-life:

- Provide relief from pain and other distressing symptoms.
- Affirm dying as a normal process.
- Integrate psychological and spiritual aspects of patient care.
- Offer a support system to help the family cope during the patient's illness and in their own bereavement.

### Inclusion Criteria

- Patient enrolled in hospice or palliative care, or who have advanced care directives or POLST, experiencing complaints related to the illness for which the patient is receiving those services.

### Exclusion Criteria

- Complaints unrelated to the illness for which they are receiving those services.

# POLST / Palliative Care

## End-of-Life Care/Palliative Care (conti):

### **PATIENT MANAGEMENT:**

#### **Patients with decision making capacity:**

If the patient is able to communicate and has the capacity to make decisions regarding treatment and transport, consult directly with the patient before treatment or transport.

#### **Patients without decision making capacity:**

If the patient lacks the capacity to make decisions regarding treatment or transport, identify any advanced care planning in place for information and consent for treatment, including:

- POLST or similar form
- Advanced care directives
- Guardian, healthcare power of attorney, or other accepted healthcare proxy
- In collaboration with hospice or palliative care provider, coordinate with guardian, healthcare power of attorney, or other accepted healthcare proxy if refusal of transport is considered.

### **TREATMENT CONSIDERATIONS:**

If the patient has **EXCESSIVE SECRETIONS**: provide suction

If the patient requires **PAIN RELIEF**: see Pain Management guideline

If the patient has **NAUSEA**: see Abdominal/Flank pain, Nausea & Vomiting guideline

### **PATIENT SAFETY CONSIDERATIONS:**

Careful and thorough assessments should be performed to identify complaints not related to the illness for which the patient is receiving hospice or palliative care. Care should be delivered with the utmost patience and compassion.

### **KEY CONSIDERATIONS:**

- Social interactions with family may affect end-of-life care.
- Scene safety should be considered when deciding on management.

### **PERTINENT ASSESSMENT FINDINGS:**

- Vital signs
- Pain score
- Neurological exam
- Lung sounds

### **KEY DOCUMENTATION ELEMENTS:**

- Interaction with hospice or palliative care provider.
- Confirmation of advanced directive or POLST
- Pain score, if applicable

# Refusal

## PURPOSE:

To define the requirements a patient must meet in order to refuse treatment and/or transport to the hospital.

## GUIDELINE:

Any patient refusing treatment must be informed of the risk of potential worsening of their condition, and the possibility it could lead to death or permanent disability.

A patient may refuse care:

- **IF** the patient has **CAPACITY** (see below);
- **AND** has no signs of being under the influence of an intoxicating substance;
- **AND** is alert and oriented to person, place, and time;
- **AND** is not a minor;
- **AND** is not showing signs of suicidal ideation or homicidal intent;
- **AND** still refuses;

**THEN** the patient must sign a refusal form indicating they understand and are accepting the risk of refusal and cannot hold anyone responsible for any negative outcome as a result of their refusal. If there are any questions or concerns about a patient's state of mind (capacity, intoxication, or altered mental status) who is refusing care or transport, EMS providers should involve online medical direction, enlist the help of family members or friends, and/or notify law enforcement for potential protective custody as appropriate.

**For the purpose of EMS, a patient with CAPACITY is defined as:**

- At least 18 years old (unless emancipated minor);
- **AND** is alert, responsive, oriented to person, place, time, and situation;
- **AND** has no signs of injury or illness which may impair the ability to make an informed decision;
- **AND** displays no signs of the patient's judgment being impaired by an intoxicating/mind altering substance (including carbon monoxide);
- **AND** is not suicidal or homicidal and does not want to hurt themselves;
- **AND** the patient demonstrates an understanding of:

**1. Diagnosis, possible diagnosis, or current medical problem:** Does the patient understand the condition/medical problem for which the specific treatment/transport is being offered?

**2. Nature and purpose of treatment:** Is the patient able to explain the nature of the treatment and understand relevant information?

**3. Risk and benefits of proposed treatment/transport:**

- Is the patient aware of the possible outcomes of treatment, alternatives, or lack of treatment, and is able to verbalize the potential danger/risk to their health and well-being by refusing transport/care?
- Is the patient able to make a decision and communicate a choice, and/or the expectations realistic? Are they able to manipulate the information rationally?

# Refusal

## **DOCUMENTATION:**

- Documentation of the refusal requires a patient care report (PCR) with as much information regarding the patient's evaluation as possible, including, but not limited to:
  - Any history obtained and any physical exam or objective observations. This may include visual descriptions if the patient declined any physical exam.
  - Documentation describing the discussions about the risks of refusal and options presented to the patient.

## **A BLS Provider may obtain a refusal:**

- If a BLS level call (ie patient overall well appearing with no significant complaints or high acuity complaints).
- If there is no ALS Provider on scene, including before ALS has arrived to the scene, if en route.
- If the ALS Provider is occupied with care of a more seriously ill or injured patient(s) on scene.
- If there are multiple patient refusals within the same call.

**NOTE:** Multiple services do not need to obtain refusals from the same patient, and the responsibility to obtain a refusal should fall to the agency having jurisdiction (AHJ) for the call, the agency holding incident command, or the transporting EMS agency.



# Termination of Resuscitation

1. Licensed EMS personnel are not obligated to continue resuscitation efforts that have been started by other persons at the scene if the patient meets the criteria listed in the Prehospital Death Determination guideline. This includes telephone CPR initiated by Emergency Medical Dispatchers.
2. Resuscitation should be terminated/not initiated if a valid DNR/POLST or physician written order is provided.
3. Resuscitation started in the field may be discontinued when the following conditions have been met:
  - A. For **medical arrest**:

The patient remains in persistent asystole or agonal rhythm after twenty (20) minutes of appropriate resuscitation, to include:

    - 1) CPR
    - 2) Effective ventilation with 100% oxygenation
    - 3) Administration of appropriate ACLS medications, if available
    - 4) Confirm asystole, no organized rhythm, PEA < 40, or a “No Shock Advised” on AED
  - B. For **traumatic arrest**:
    - 1) Hemorrhage control
    - 2) Airway management with least-invasive approach needed to maintain patency and effective ventilations with 100% oxygenation for two (2) minutes
    - 3) Perform bilateral needle decompression only if signs of chest trauma or if tension pneumothorax suspected
    - 4) Provide CPR *ONLY* after above life saving interventions have been performed
    - 5) Confirm asystole, no organized rhythm, PEA < 40, or a “No Shock Advised” on AED
  - C. The patient develops, or is found to have one of the following conclusive signs of death at any point during the resuscitative effort:
    - 1) Lividity
    - 2) Rigor mortis



Contact Medical Control for cases not meeting above criteria for Termination of Resuscitation.

4. When resuscitation has been terminated in the field, all medical interventions shall be left in place.
5. If possible, do not leave a body unattended. Once a responsible person (i.e. coroner, law enforcement, fire, etc.) is present at the scene, you may be excused.
6. **NEVER** transport/move a body without permission from the coroner, except for assessment or its protection.



If the body is in the public view and cannot be isolated, screened, or blocked from view, and is creating an unsafe situation with citizens/family, the body can be covered with a clean sheet or *STERILE BURN SHEET* obtained from the EMS vehicle.

# **PROCEDURES GUIDELINES**

# Continuous Positive Airway Pressure (CPAP)

LEVEL: EMT/AEMT/Paramedic

## Indications:

This process may be performed on any patient 15 years old or older in severe respiratory distress, asthma, COPD, CHF, pulmonary edema, HAPE, or pneumonia, who has *TWO* of the following:

- A. Retractions, accessory muscle use, and/or increased work of breathing
- B. Respiratory rate >20 per minute
- C. SpO<sub>2</sub> ≤ 94%

## Contraindications:

- A. Inadequate spontaneous respiratory effort or apnea
- B. Altered mental status or unable to follow commands
- C. Unable to protect airway
- D. Systolic blood pressure < 90 mmHg
- E. Major trauma/pneumothorax
- F. Vomiting or active GI bleed

Use device per  
manufacturer  
instructions

## Key procedural considerations:

- A. Assess patient and document VS, SpO<sub>2</sub> and ETCO<sub>2</sub> if available prior to applying oxygen. Paramedics should document ETCO<sub>2</sub>.
- B. Select the appropriate size face mask for the patient.
- C. Inform the patient about procedure process.
- D. If using CPAP, gradually increase the flow rate, slowly reaching the desired CPAP pressure. Secure face mask onto patient's face using the head harness.
- E. Check the mask and tubing for leaks.
- F. Reassess patient and document every five minutes.
- G. If the patient develops any of the contraindications or requires definitive airway control, discontinue CPAP and provide necessary airway control.

# Electrical Therapy/Defibrillation

LEVEL: Paramedic

## Indications:

This procedure may be performed on any patient experiencing:

- A. Ventricular fibrillation
- B. Pulseless ventricular tachycardia
- C. Torsades de Pointes

**Contraindications:** None

Use device per  
manufacturer  
instructions

## Key procedural considerations:

- A. The initial and subsequent attempts shall be at the energy level(s) suggested by the device manufacturer and/or the agency's medical director.
- B. Defibrillation should be immediately provided in an arrest **WITNESSED** by EMS personnel. In an arrest that is **UNWITNESSED** by EMS personnel, two minutes of CPR should be provided prior to defibrillation.
- C. Patients with automatic implantable cardioverter-defibrillators (AICD) will need external defibrillation if the AICD is ineffective.
- D. If defibrillation is needed on a patient with a permanent implanted pacemaker, the defibrillator paddles or self adhesive electrodes should be placed at least one inch from the pulse generator of the pacemaker.



Initial attempt at pediatric defibrillation shall be at 2 J/kg. If unsuccessful, defibrillation should be attempted at 4 J/kg. Repeated defibrillations should be at >4 J/kg up to 10 J/kg until conversion occurs. Adult paddles/pads may be used in children weighing more than 10 kg.

# Electrical Therapy/Synchronized Cardioversion

LEVEL: Paramedic



The patient **MUST** be on a cardiac monitor and **SHOULD** have Vascular Access

## Indications:

This procedure may be performed on any patient experiencing:

- A. Ventricular tachycardia with inadequate perfusion
- B. Supraventricular tachycardia with inadequate perfusion
- C. Ventricular tachycardia with adequate perfusion, but refractory to drug therapy

**Contraindications:** None

## Adjunctive therapy:

In the conscious patient with a systolic blood pressure of >90mmHg consider:

**Sedation:** Midazolam 0.1 mg/kg IN/IM/IV/IO, max dose 5 mg OR;

Lorazepam 0.1 mg/kg IM/IV/IO, max dose 2 mg OR;

Diazepam 5 mg IV/IO.

**Analgesia:** Morphine Sulfate up to 0.1 mg/kg slow IV/IO to a maximum single dose of 10 mg OR;

Fentanyl up to 1 mcg/kg IN/IM/IV/IO to a maximum single dose of 100 mcg OR;

Ketamine 0.2 mg/kg IN/IM/IV/IO to a maximum single dose of 25 mg

## Key procedural considerations:

- A. **Biphasic device:** The initial and subsequent attempts shall be at the energy level(s) suggested by the device manufacturer and/or the agency's medical director.
- B. **Monophasic device:**
  - 1. Ventricular dysrhythmias: 100 J escalating to 200, 300, and 360
  - 2. Supraventricular dysrhythmias: 50 J with subsequent attempts at 100 J



Pediatric synchronized cardioversion is by Medical Control order only.



Initial attempt at pediatric cardioversion shall be at 0.5 J/kg.

If unsuccessful, cardioversion should be attempted at 2 J/kg.

Adult paddle/pads may be used in children weighing more than 10 kg.

# Electrical Therapy/Transcutaneous Pacing

LEVEL: Paramedic

## Indications:

This procedure may be performed on any patient experiencing:

- A. Hemodynamically unstable bradycardia
- B. Unstable clinical condition that is likely because of bradycardia
- C. For pacing readiness (i.e. standby mode) in the setting of MI with bradycardia, second degree type II AV block, third degree AV block, new left or right alternating BBB or bifascicular block
- D. Overdrive pacing of tachycardias refractory to drug therapy or electrical cardioversion

**Contraindications:** None

## Adjunctive therapy:

In the conscious patient with a systolic blood pressure of >90mmHg consider:

**Sedation:** Midazolam 0.1 mg/kg IN/IM/IV/IO, max dose 5 mg OR;

Lorazepam 0.2 mg IM/IV/IO, max dose 2 mg OR;

Diazepam 5 mg IV/IO.

**Analgesia:** Morphine Sulfate up to 0.1 mg/kg slow IV/IO to a maximum single dose of 10 mg OR;

Fentanyl up to 1 mcg/kg IN/IM/IV/IO to a maximum single dose of 100 mcg OR;

Ketamine 0.2 mg/kg IN/IM/IV/IO to a maximum single dose of 25 mg

## Key procedural considerations:

- A. Apply pacing pads, begin pacing at a rate of 60 beats per minute at the lowest available current.
- B. Increase current by 20 milliamp increments until electrical capture.
- C. In the event of electrical capture and no pulses, continue pacing and CPR.



Pediatric pacing is by Medical Control order only.

# Endotracheal Intubation

LEVEL: Paramedic

1. All intubations **MUST** have initial, en route, and at transfer of care End-Tidal CO2 detection/capnography performed and recorded on the PCR.
2. All intubation attempts **MUST** be documented on the PCR.

## Indications:

This procedure may be performed on any patient in whom attempts at basic airway and ventilatory support are unsuccessful **AND** who has at least one of the following:

- A. Hypoxia
- B. Respiratory arrest/failure
- C. Inability to maintain airway patency

## Contraindications:

Absolute contraindications: None

Relative contraindications:

- A. Presence of gag reflex
- B. Suspected narcotic overdose/hypoglycemia prior to administration of NALOXONE/GLUCOSE

Check and prepare the  
endotracheal airway device prior  
to insertion

## Key procedural considerations:

- A. Position head properly.
- B. Insert blade while displacing tongue and elevate mandible with laryngoscope.
- C. Introduce ET tube and advance to proper depth.
- D. Inflate cuff to proper pressure and disconnect syringe.
- E. Ventilate patient and confirm proper placement.
- F. Verify proper tube placement by secondary confirmation such as capnography or colorimetric device.
- G. Secure device or confirm that the device remains properly secured.

**Mandatory notification of Medical Director within 24 hours of any failed intubation attempt.**  
For simplicity, may use this online form: <https://forms.gle/G3RPJQ8fEugRafMTA>

# Hemorrhage Control

LEVEL: EMR/EMT/AEMT/Paramedic \*Note: Use of TRANEXAMIC ACID (TXA) is Paramedic only\*

## Hemorrhage:

This procedure may be performed on any patient that has bleeding from an extremity, junctional hemorrhage, or torso hemorrhage.

## Extremity Hemorrhage – Tourniquet Application:

- A. Apply tourniquet proximal to the bleeding site.
- B. Absolute contraindication: Bleeding has stopped or controlled with less invasive measure.
- C. If bleeding is not controlled, consider additional tightening or applying a second tourniquet proximal side by side to the first.
- D. Wound packing does not preclude you from placing a tourniquet.


## Junctional Hemorrhage – Wound Packing:

- A. Junctional Hemorrhage Defined: hemorrhage occurring at the junction of an extremity with the torso, and/or the base of the neck.
- B. Use direct pressure and an appropriate pressure dressing with deep wound packing (plain gauze or, if available, hemostatic gauze).
- C. Absolute Contraindication: Hemostatic gauze use on hemorrhaging abdominal wounds.


## Torso Hemorrhage – Consider TRANEXAMIC ACID (TXA) for blunt or penetrating chest or abdominal trauma with suspected blood loss. Paramedic Administration Only.

- 1. Dose: Adults (age > 16 yo OR weight > 45 kg): 2 g IV/IO over 10 min for patients with SBP<90 and/or HR>110.
- 2. TRANEXAMIC ACID is ideally given within the first 90 minutes of injury.
- 3. TRANEXAMIC ACID administration is contraindicated if the traumatic injury occurred more than 3 hours prior to proposed administration.

Tourniquet Placement



Junctional Hemorrhage





# High Flow Nasal Cannula (HFNC)

LEVEL: Paramedic

## Indications:

Acute hypoxic respiratory failure (examples include asthma, pneumonia, COVID-19 infection, pulmonary embolism, pulmonary hypertension, interstitial lung disease, or other respiratory process)

## Contraindications:

- A. Abnormalities or surgery of the face, nose, or airway that preclude appropriate-fitting nasal cannula
- B. Basilar skull fracture or severe facial trauma
- C. Upper airway surgery to avoid theoretical risk that high pressure may precipitate a venous thromboembolism
- D. Circumstances in which CPAP, endotracheal intubation, or surgical airway is indicated
- E. Inability to provide continuous, heated humidification using an approved delivery device
- F. Inability to provide therapy through an appropriate sized nasal prongs
- G. Insufficient supply of oxygen to complete the transport

Use device per  
manufacturer  
instructions

## Key procedural considerations:

- A. Ensure that an adequate supply of oxygen is available.
  - 1. Calculate the amount of oxygen needed for interfacility transport based on current therapy:
$$\text{Minutes of available oxygen} = \frac{(\text{Tank PSI} \times \text{Tank Factor})}{(\text{Flow rate} \times \text{FiO}_2)}$$

Tank PSI: pounds  
Tank Factor: D size = 0.16; E size = 0.28; G size = 2.41; H and K size = 3.14; M size = 1.56  
Flow rate: liters per minute  
FiO<sub>2</sub>: expressed as a fraction (eg 40% = 0.40)
  - 2. Estimate total duration of interfacility transport and ensure there is at least twice the amount of necessary oxygen available.
- B. Perform appropriate patient assessment, including obtaining vital signs, pulse oximeter, and cardiac rhythm.
- C. Set FiO<sub>2</sub> to maintain SpO<sub>2</sub> at or above 90% (range 21-100%).

# High Flow Nasal Cannula (HFNC)

## Key procedural considerations (conti):

- D. Set flow rate in liters per minute (L/min) to decrease work of breathing:
  - 1. Utilize facility settings as starting point, if available.
  - 2. Pediatric flow calculation: 2 L/kg/min up to adult flow rate.
  - 3. Adult flow calculation: initial 50 L/min (range 5-60 L/min).
- E. In general, maximize the flow rate first and attempt to keep the  $\text{FiO}_2 \leq 60\%$ .
- F. Reassess vital signs, work of breathing, mental status, and breath sounds.
- G. Consider the need for escalation of respiratory support if patient remains in respiratory failure on more than 2 L/kg/min of flow or maximum settings for the delivery device.

## Complications:

- A. Abdominal distension
- B. Aspiration
- C. Rarely barotrauma (eg pneumothorax)



**FOR CIRCUMSTANCES IN WHICH THE PATIENT DOES NOT IMPROVE OR CONTINUES TO DETERIORATE ON HFNC, TERMINATE HFNC ADMINISTRATION AND ESCALATE RESPIRATORY SUPPORT TO PROVIDE POSITIVE PRESSURE VENTILATION VIA CPAP, BVM, SUPRAGLOTTIC AIRWAY, OR ENDOTRACHEAL INTUBATION, IF NECESSARY.**

# Legal Blood Draw

LEVEL: EMT w/ IV/IO Initiation Endorsement/AEMT/Paramedic

**Guideline:**

Steps for Legal Blood Draws:

1. Obtain patient ID
2. Ask their full name and DOB
3. Use the kit law enforcement provides
4. Prepare the site with a non-alcohol prep
5. Draw gray topped tube from the kit, label it, and hand the tube to law enforcement
6. Fill out the paperwork from the kit and hand the paperwork to law enforcement
7. Document a patient care record (PCR)

Example: "On May 7th at 1800 I drew blood from Tom Smith, DOB 1/2/99, into a gray topped vacutainer, which I subsequently labeled with the name Tom Smith and my initials (BB). Tube and associated paperwork handed to officer John Doe of the Bozeman Police Department, Badge #123."

# LUCAS Device

LEVEL: EMT/AEMT/Paramedic

## Rationale:

The LUCAS is a non-invasive mechanical CPR device. It has a role in providing uninterrupted chest compressions at an appropriate rate and depth, but it has **NOT** been shown to be superior to well-performed manual chest compressions. Manual compressions are still considered the standard of care by the AHA. Consider the device a tool to use in non-traumatic cardiac arrest, but it should not distract from beginning manual chest compressions as soon as possible, or cause unneeded interruptions in compressions at any point. Use of a LUCAS device can make chest compressions during transport more effective and safer for the transporting crew. However, availability of a LUCAS should not prompt transport when it would not have otherwise been considered. Cardiac arrests are still best managed in place unless there are extenuating circumstances mandating transport such as v-fib resistant to multiple shocks.

## Indications:

Non-traumatic cardiac arrest requiring CPR

## Contraindications:

- Traumatic arrest
- Patient too small for the pressure pad in the suction cup to make contact with the chest OR patient too large for the legs to lock
- Patient is a child under 12 years old
- Pregnancy

## Key procedural considerations:

- The LUCAS shall be used in accordance with the manufacturer's recommendations.
- LUCAS should not be used until after two full cycles of manual compressions. A defibrillator should be applied before the LUCAS. Starting manual chest compressions and initial defibrillation should take precedence over placement of the LUCAS. Use a two-step application and minimize pauses in chest compressions.
- The machine is a tool, but not a priority. Placement of the device should be done to take opportunity of inevitable pauses and to minimize no-flow states.
- The LUCAS should **NOT** be paused for intubations.
- Do **NOT** attempt to lift the patient or the device by the arm straps.
- A member of the agency that placed the device and has been trained on the LUCAS must remain with the patient at all times until the LUCAS is removed. That person shall be responsible for the safe operation of the device.
- One agency's hood may be exchanged for another, but keep the initial back plate in place and minimize any interruptions.



## Device malfunction:

If there is a device malfunction, immediately remove the device and resume high performance CPR. The device may be reapplied **ONLY** after the problem has been addressed.

# Medication Administration

LEVEL: EMR/EMT/EMT w/Medication Endorsement/AEMT/Paramedic (based on medication)

## Indications:

This procedure may be performed on any patient that requires the administration of a medication.

## Key procedural considerations (GENERAL):

- A. Inquire about allergies and previous medication reactions
- B. Check and recheck medication
- C. Solution clarity and expiration date
- D. Right drug
  - Right patient
  - Right dose
  - Right time
  - Right route
  - Right documentation
- E. Dispose of syringe and other material in proper container

## Intravenous and Intraosseous Bolus Medications

### Key procedural considerations:

- A. Identify and cleanse injection site closest to the patient
- B. Administer correct dose at proper push rate
- C. Turn IV on and adjust drip rate to TKO/KVO

## Intramuscular and Subcutaneous Drug Administration

### Key procedural considerations:

- A. Needle should be 20 gauge or smaller
  - B. Locate administration site
    - Deltoid muscle
    - Vastus lateralis (lateral thigh) muscle
    - Ventrogluteal or dorsogluteal muscles (buttocks)
- | <u>IM</u>                                | <u>SQ</u>                                |
|------------------------------------------|------------------------------------------|
| Pull skin tight                          | Pinch to lift skin slightly              |
| Insert needle at a 90° angle to the skin | Insert needle at a 45° angle to the skin |
| Advance into muscle layer                | Advance into subcutaneous layer          |

## Mucosal Atomizer Device (MAD) Administration

**Medications: FENTANYL, GLUCAGON, KETAMINE, MIDAZOLAM, NALOXONE**

### Key procedural considerations:

- A. Using the free hand, hold the crown of the head stable.
- B. Place the tip of the MAD snugly against the nostril, aiming slightly up and outward (toward the top of the ear).
- C. Briskly compress the syringe to deliver half the medication into the nostril.
- D. Move the device over to the opposite nostril and administer the remaining medication.

# Needle Cricothyroidotomy

LEVEL: Paramedic

## Indications:

This procedure may be performed on any patient with:

- A. Total airway obstruction by any BLS or ALS procedures, *OR*
- B. Inability to adequately ventilate AND oxygenate with any provider level emergency care procedures prior to the attempt.

## Contraindications:

- A. Inability to identify landmarks (cricothyroid membrane)
- B. Underlying anatomical abnormality (tumor)
- C. Tracheal transection
- D. Acute laryngeal disease due to infection or trauma



You **MUST** use a 14 gauge over-the-needle catheter attached to a 10 cc syringe or commercial cricothyroidotomy device.

## Key procedural considerations:

\*Please follow Manufacturer's Instructions\*

- A. Position patient supine (if possible), hyperextending the head.
- B. Locate cricothyroid membrane and clean site thoroughly.
- C. Stabilize cricoid and thyroid cartilages with one hand.
- D. Puncture needle/catheter at a 90° angle and then change insertion angle to 45° up to the stopper; gently aspirate with attached syringe.
- E. When syringe is able to aspirate air, stop advancing needle.
- F. Remove the stopper from the cannula and advance the cannula only until the phlange is flush with the patient's neck. Remove the metal needle from the cannula. Remove the syringe.
- G. Secure the cannula with the neck strap.
- H. Apply connecting tube and attach to BVM and ventilate patient.

**Mandatory notification of Medical Director within 24 hours of any attempted, successful, and/or failed cricothyroidotomy in the field. For simplicity, may use this online form:**

<https://forms.gle/G3RPJQ8fEugRafMTA>

# Needle Decompression

LEVEL: Paramedic

## Indications:

This procedure may be performed on any patient who has evidence of a tension pneumothorax, demonstrated by the following criteria:

1. Severe/progressive respiratory distress and/or increased resistance to bagging, AND unilateral diminished/absent breath sounds, AND:
  - A. Hypotension with signs of shock, or
  - B. Persistent hypoxia despite supplemental oxygen, or
  - C. Jugular venous distention, or
  - D. Tracheal deviation (late sign)
2. Any traumatic cardiac arrest with chest or abdominal trauma and undergoing resuscitation should have bilateral needle thoracostomy performed as soon as possible.

**Contraindications:** None



Needle Decompression is permitted in pediatric patients.

## Key procedural considerations:

- A. Select and identify insertion site:
  1. Primary site is the 4<sup>th</sup> or 5<sup>th</sup> intercostal space in the mid-axillary line of the affected side.
    - a. Needle should be placed within the “triangle of safety”. Insertion site must be above the nipple line as the nipple lies flat against the chest wall with the arm abducted.
    - b. In females, the breast can displace the nipple inferiorly. If displaced, the clinician should identify where the nipple would lie if flat against the chest wall. This will be superior to the inframammary fold/crease. When in doubt, a more superior site is preferred.
  2. Alternate site is the 2<sup>nd</sup> intercostal space in the mid-clavicular line of the affected side.
- B. Use appropriate size needle and length.
- C. Prep site with appropriate disinfectant (e.g. iodine, chlorhexidine, alcohol).
- D. Place tip of needle on top of appropriate rib and insert over top of rib into intercostal space.
- E. Advance needle into pleural space and remove needle. Leave catheter in place. Do not hub the needle.
- F. Consider attaching a one-way valve, if available.

# Push Dose Epinephrine

LEVEL: Paramedic

## History:

When norepinephrine was added and dopamine removed from the stock of medications carried in the prehospital setting, concern was raised about the ACLS algorithm for unstable bradycardia as norepinephrine does not primarily increase heart rate and is not indicated for bradycardia.

## Indications:


- Unstable or symptomatic bradycardia with a pulse
- Shock unresponsive to fluid resuscitation
- Post-ROSC with persistent hypotension
- Anaphylaxis/allergic reaction not responding to IM epinephrine 1:1000
- Sepsis while initiating norepinephrine drip

## Preparation:

- Mix 1 ml of epinephrine 1:10,000 (cardiac epi) with 9 ml of normal saline (NS) in a 10 ml syringe
- Results in a 10 mcg/ml concentration

## Dosage:

10 mcg (1 ml) every 2-5 minutes titrated to SBP > 90 or MAP > 65





# Spinal Motion Restriction

LEVEL: EMR(c-collar only)/EMT/AEMT/Paramedic



Spinal motion restriction is indicated in any patient who meets the indications (A-E) below:

## Indications:

This procedure may be performed on any patient with potential for spinal injury based upon the following (NEXUS) criteria:

- A. Midline cervical spinal tenderness
- B. Focal neurologic deficit
- C. Altered mental status
- D. Evidence of drug and/or alcohol intoxication
- E. Any painful, distracting injury

## Contraindications:

Spinal motion restriction is **NOT** performed in the following conditions:

- A. Penetrating trauma to the head and/or neck and no evidence of spinal injury
- B. Injuries where placement of the cervical collar might compromise patient assessment, airway management, ventilation and/or hemorrhage control
- C. Patients in cardiac arrest

## Key procedural considerations:

- A. If (A-E) above are **ALL NEGATIVE**, spinal motion restriction is not required.
- B. If required, spinal motion restriction is the placement of an approved, properly-sized cervical collar before the patient is moved.
- C. Tape, head straps, wedges, and head and/or neck support devices are generally not recommended.
- D. Patients found in motor vehicles should be asked if they are able to exit the motor vehicle on their own. If so, they should be assisted to a soft stretcher and secured for transport. Patients unable to exit the vehicle on their own accord should be removed by the appropriate extrication method.
- E. Once on the stretcher, the patient may be moved to a semi-Fowler's or high-Fowler's position for comfort.
- F. If a backboard is used for extrication or movement, the patient should be immediately moved to a soft mattress, if possible.
- G. In special situations, alternate stabilization devices (e.g. vacuum mattress, KED, etc.) may be used as indicated.
- H. Pediatric patients may be stabilized in an approved car seat or with a commercial pediatric stabilization device.

# Supraglottic Airway Device

LEVEL: EMT w/ Airway Endorsement/AEMT/Paramedic

## Indications:

This procedure may be performed on any patient in which attempts at basic airway and ventilatory support are unsuccessful *AND* who has at least one of the following:

- A. Hypoxia
- B. Respiratory arrest/failure
- C. Obtundation
- D. Failed endotracheal intubation

## Contraindications:

- A. Gag reflex
- B. History of esophageal trauma, or known esophageal disease
- C. Recent ingestion of a caustic substance
- D. Tracheostomy or laryngectomy
- E. Suspected foreign body obstruction

Check and prepare the  
supraglottic airway  
device prior to insertion

## Key procedural considerations:

- A. Pre-oxygenate the patient.
- B. Position the patient's head in a neutral or slightly flexed position if no suspected spinal injury (if a spine injury is suspected, maintain a neutral, in-line head position).
- C. Perform a tongue-jaw lift.
- D. Insert device to proper depth. *NEVER* force. If device does not advance, readjust the insertion.
- E. Secure device in the patient.
- F. Ventilate patient and confirm proper ventilation (correct lumen and proper insertion depth) by auscultation bilaterally over lungs and over epigastrium.
- G. Adjust ventilation as necessary (slightly withdraw tube until ventilation is optimized).
- H. Verify proper tube placement by secondary confirmation such as capnography or colorimetric device.
- I. Secure device or confirm that the device remains properly secured.

# Surgical Cricothyroidotomy

LEVEL: Paramedic

## Indications:

This procedure may be performed on any patient with:

- A. Total airway obstruction by any BLS or ALS procedures, *OR*
- B. Inability to adequately ventilate AND oxygenate with any provider level emergency care procedures prior to the attempt.

## Contraindications:


- A. Inability to identify landmarks (cricothyroid membrane)
- B. Underlying anatomical abnormality (tumor)
- C. Tracheal transection
- D. Acute laryngeal disease due to infection or trauma

## Key procedural considerations:

### **\*\*Must use North American Rescue Bougie-Aided Cricothyroidotomy Pack (BAC-Pack)\*\***

- A. Position patient supine (if possible), hyperextending the head.
- B. Locate cricothyroid membrane and clean site thoroughly.
- C. Stabilize cricoid and thyroid cartilages with one hand.
- D. Insert scalpel through skin, subcutaneous tissue, and cricothyroid membrane.
- E. Extend incision laterally in both directions.
- F. Prior to removing scalpel, insert tracheal hook.
- G. Insert the bougie down the trachea. Advance tube and inflate cuff. Remove bougie.
- H. Attach to BVM and ventilate patient.

Mandatory notification of Medical Director within 24 hours of any attempted, successful, and/or failed cricothyroidotomy in the field. For simplicity, may use this online form: <https://forms.gle/G3RPJQ8fEugRafMTA>



# Tracheostomy Tube Replacement

LEVEL: Paramedic

**Indications:**

This procedure may be performed on any patient that has A *TRACHEOSTOMY TUBE* and *WHO HAS*:

- A. Hypoxia
- B. Respiratory arrest/failure
- C. Obtundation
- D. Secretions unable to be cleared by suctioning

**Contraindications:** None

**Key procedural considerations:**

- A. If the patient or family has a replacement tube available, it may be used. If a replacement tube is not available, an endotracheal tube of a similar outer diameter may be used.
- B. Premoisten the tube with water soluble lubricant.
- C. Extend the neck and, if necessary, place a roll between the patient's shoulder blades to aid in visualizing the stoma.
- D. If the tube cannot be placed easily, withdraw the tube; administer oxygen and positive pressure ventilation. *NEVER* force the tube.
- E. Secure the device to the patient.
- F. If the tube cannot be easily placed, a suction catheter may be used as a guide.

# Traction Splint

LEVEL: EMT/AEMT/Paramedic

**Indications:**

This procedure may be performed on any patient with an isolated midshaft femur fracture.

**Contraindications:**

- A. Pelvic fracture or instability
- B. Knee, lower leg, or ankle instability

**Key procedural considerations:**

- A. Assess motor, sensory, and circulatory function in the involved extremity.
- B. Apply traction splint per the manufacturer's guidelines.
- C. Initiate mechanical traction to match manual traction.
- D. Reassess motor, sensory, and circulatory function in the involved extremity.
- E. Exercise care when applying traction not to reintroduce bone ends into the body.

**Pearls**

- Femur fractures can be difficult to identify and difficult to exclude comorbid injuries.
- No benefit of traction in preventing shock or reducing need for blood transfusion.
- Traction only minimally better for pain (based on a single study with weak evidence).
- Static (non-traction) splinting is acceptable alternative.

# Ultrasound

LEVEL: Paramedic

## Procedure:

Ultrasound will be available for the following modalities outlined below:

- A. EFAST
- B. Vascular Access
- C. Cardiac Arrest

## EFAST

### Indications:

May perform on patients suspected of having intra-abdominal free fluid, pericardial effusion, thoracic free fluid, or pneumothorax, which may guide appropriate therapy and resuscitation.

### Contraindications:

Higher priority critical actions needed, i.e., stabilization of ABCs.

### Technique:

Obtain the following views:

- a) Abdominal RUQ (right upper quadrant)
- b) Abdominal LUQ (left upper quadrant)
- c) Cardiac Subcostal OR Parasternal Long
- d) Pelvis
- e) Right lateral chest
- f) Left lateral chest
- g) Right anterior chest
- h) Left anterior chest

### Complications:

False positive exams:

- a) Intra-abdominal free fluid may not always be acute hemoperitoneum (e.g. patients with liver cirrhosis may have ascites).
- b) Pericardial effusion may not always be acute hemopericardium (e.g. patients with chronic pericardial effusions due to CHF, uremia, malignancy, inflammatory processes like lupus).
- c) Thoracic fluid may not always be acute hemothorax (e.g. patients with volume overload from heart/liver/kidney failure, parapneumonic effusions).
- d) Lack of lung sliding may not always be due to a pneumothorax (e.g. patient with subcutaneous air, prior pleurodesis, hyper-inflation from severe COPD/asthma, main-stem intubation, apnea).

# Ultrasound

## **Vascular**

### **Indications:**

- a) Multiple failed attempts at peripheral access.
- b) Need for large bore access.
- c) Need for central access for vasopressors.

### **Contraindications:**

- a) Should not be used to replace IO (intraosseous) lines in trauma patients that need fast vascular access.
- b) Same contraindications as non-ultrasound-guided vascular access (ie overlying cellulitis, suspected ipsilateral extremity fracture, ipsilateral lymph node dissection).

### **Technique:**

- a) Evaluate possible vascular access locations and identify compressible vein.
- b) Prep the skin with Chlorhexidine swab.
- c) Cleanse the ultrasound probe with Chlorhexidine swab.
- d) Place sterile ultrasound gel on the clean skin.
- e) Under ultrasound guidance, place angiocath into the vein and thread the catheter.
- f) Attach angiocath cap.
- g) Flush the catheter.
- h) Apply tegaderm and tape to secure the IV.

### **Complications:**

- a) Local hematoma
- b) Arterial stick
- c) Pain
- d) Infection

# Ultrasound

## **Cardiac Arrest**

### **Indications:**

The performance of ultrasound during cardiac arrest is controversial due to evidence that it is associated with prolonged pulse checks. However ultrasound during arrest has also been shown to identify reversible causes and also may be prognostic and aid in the decision to continue resuscitative efforts. For the performance of ultrasound during cardiac arrest a regimented approach should be followed to ensure that rhythm checks are not prolonged by it's usage, and that specific questions are answered which may directly impact the resuscitation.

### **Relative Contraindications:**

- a) Higher priority critical actions needed, i.e., administration of epinephrine or defibrillation.
- b) Lack of available provider to perform exam. Code leader should avoid becoming task fixated on performing scan.

### **Technique:**

Operational Approach to Image Acquisition

- (1) Do it early so there is time to act on data
- (2) Find subcostal or apical window during compressions
  - (a) If no subcostal or apical window then be ready to look parasternal at pause

Interpretation and Action Based on Findings

- (1) Pericardial Effusion
- (2) Absence of organized cardiac activity
  - (a) Absence of cardiac activity is a poor prognostic indicator, however, it should not be used in isolation to cease resuscitative efforts as there are documented survivors in this group and additionally there is variability in provider interpretation of cardiac standstill.

### **Complications:**

- a) Prolonged interruption in chest compressions.
- b) False positives:
  - (1) Falsely identifying a pericardial effusion.
  - (2) Falsely calling cardiac standstill in patient with severe bradycardia or profoundly reduced contractility.



# Vagal Maneuvers

LEVEL: Paramedic



The patient **MUST** be attached to a cardiac monitor and **MUST** have vascular access prior to performing the procedure

## Indications:

This procedure may be performed on any patient who is experiencing Supraventricular Tachycardia with adequate perfusion.

**Contraindications:** None

## Key procedural considerations:

A. Approved methods include:

1. Valsalva maneuver
2. Head-down tilt with deep inspiration
3. Activation of the “diving reflex” by facial immersion in ice water (unless ischemic heart disease is present)
4. Carotid massage (only on patients under 40 years of age)



B. In infants and young children, the most effective vagal maneuver is the application of ice to the face. IV access is not mandatory prior to vagal maneuvers in children.

# Vascular Access

LEVEL: EMT w/ IV/IO Initiation Endorsement/AEMT/Paramedic



Vascular access attempts should not unnecessarily delay transport, attempts should be completed enroute.  
All attempts are to be documented on the PCR.

## Indications for Peripheral Vascular Access:

This procedure may be performed on any patient whenever there is a potential need for:

- A. Intravenous drug administration.
- B. Need to administer IV fluids for volume expansion.

**Contraindications:** None

## Key procedural considerations:

- A. Saline locks may be used when appropriate and flushed with a 3 cc bolus of NS as needed.
- B. In general extension tubing should be used on IV lines. May consider hooking IV line directly to IV catheter when need the fastest flow possible (ie no needle-free access hub or connector).

## Indications for Intraosseous Access:

Critically ill or injured patient who requires IV drugs/fluids and in whom a peripheral line cannot be immediately established.

## Contraindications:

- A. Placement in, or distal to, a fractured bone.
- B. Previous significant orthopedic procedure at the site; prosthetic limb or joint.
- C. IO catheter use in past 48 hours of the target bone.
- D. Infection at the area of insertion.
- E. Absence of adequate anatomical landmarks.

**Paramedic may administer LIDOCAINE 1% preservative-free for anesthetic if IO is to be used for a conscious patient.**

- 1) Administer 0.5 mg/kg of LIDOCAINE 0.1 mg/mL with slow push through IO needle to a maximum of 40 mg to mitigate pain from IO medication administration.
- 2) Allow LIDOCAINE to dwell in IO space for 60 seconds.
- 3) Flush IO with 5-10 ml normal saline.
- 4) Consider systemic pain medication for patients not responding to IO LIDOCAINE.

**Key procedural considerations:** Only 1 (one) attempt is permitted per extremity

## Indications for use of Previously Established Central Line Access:

This procedure may be performed on any critically ill or injured patient who requires IV drugs or IV fluids AND in whom a peripheral line cannot be established.

**Contraindications:** Inability to freely aspirate blood out of the catheter

## Key procedural considerations: Central line access (Implantable Ports, Port-A-Caths, Mediports)

A. May only be used if the device has already been accessed and IV fluid set-up has been established and running.



B. These devices require special needles (non-coring type) for access. The device may be damaged if standard jumper (conventional) needles are used to access the ports.

# FORMULARY

### **ACETAMINOPHEN (TYLENOL)**

**PROVIDER LEVEL:** EMT (PO only), AEMT, Paramedic.

**CLASS:** Analgesic, antipyretic.

**PHARMACOLOGY/ACTIONS:** Acetaminophen is a para-aminophenol derivative. Analgesic effects are believed to be due to activation of descending serotonergic inhibitory pathways in the CNS. Interactions with other nociceptive systems may also be involved. Antipyretic effects are produced from inhibition of the hypothalamic heat-regulating center.

**ONSET/DURATION:** Onset: 5-10 minutes / Duration: 4-6 hours.

**INDICATIONS:** Pain control of mild to moderate pain.

**CONTRAINDICATIONS:** Hypersensitivity/allergy, patient who has received acetaminophen in any form within the previous 6 hours, acetaminophen overdose, severe hepatic impairment, and severe active liver disease.

**SIDE EFFECTS:** Acute hepatotoxicity, nausea, and vomiting.

**DRUG INTERACTIONS:** Alcohol and Sorafenib.

**ROUTE:** PO, IV, IO.

**DOSAGE:**

**ADULTS:** 325-1000 mg PO; 1000 mg IV/IO infusion over 15 minutes

**PEDS:** 15 mg/kg (max 1000 mg) PO; 15 mg/kg (max 1000 mg) IV/IO infusion over 15 minutes

**PREGNANCY SAFETY:** Category A – considered safe during pregnancy.

**COMMENTS:** Use with caution in patients with severe hypovolemia (ie dehydrated or blood loss). Use with caution in patients with known G6PD deficiency. Generally avoid in alcoholics. Limit acetaminophen dose from all sources and all routes of administration to  $\leq 4$  grams/day for adults.

### **ACTIVATED CHARCOAL**

**PROVIDER LEVEL:** EMT, AEMT, Paramedic.

**CLASS:** Absorbent.

**PHARMACOLOGY/ACTIONS:** Activated charcoal is a fine black powder that binds and absorbs ingested toxins still present in the GI tract. It has a tremendous surface area. Once it binds and absorbs the ingested toxin, the combined complex is excreted from the body.

**ONSET/DURATION:** Onset: immediate / Duration: unknown.

**INDICATIONS:** Acute poisoning in alert patients within 1 hour of ingestion.

**CONTRAINDICATIONS:** Poisonings by corrosive agents, cyanide, iron, mineral acids, or organic solvents. Patients with altered mental status, active vomiting, absence of bowel sounds, and GI perforation.

**SIDE EFFECTS:** Vomiting, abdominal cramping, bloating, constipation, and risk of aspiration.

**DRUG INTERACTIONS:** None significant.

**ROUTE:** PO.

**DOSAGE:**

**ADULTS:** 1 gram/kg PO

**PEDS:** 1 gram/kg PO

**PREGNANCY SAFETY:** Undetermined. Generally considered safe during pregnancy as no systemic absorption.

**COMMENTS:** Consider pre-medication with antiemetic. May be combined with sorbitol, which acts as a laxative decreasing GI transit time. Consider contacting Poison Control at 1-800-222-1222 to determine if recommend administration of activated charcoal.

### **ADENOSINE (ADENOCARD)**

**PROVIDER LEVEL:** Paramedic.

**CLASS:** Antidysrhythmic, endogenous nucleoside.

**PHARMACOLOGY/ACTIONS:** Adenosine primarily is formed from the breakdown of adenosine triphosphate (ATP) which is found in every cell of the body and has a wide range of metabolic roles. Adenosine slows SVT by decreasing electrical conduction through the AV node without causing negative inotropic effects. It also acts directly on sinus pacemaker cells and vagal nerve terminals to decrease chronotropic activity.

**ONSET/DURATION:** Onset: 20-30 seconds / Duration: 30 seconds.

**INDICATIONS:** - Regular narrow-complex PSVT.

- Regular wide-complex undifferentiated tachycardia.

- if ventricular fibrillation, adenosine will likely have no effect.

- Dysrhythmias associated with bypass tracts such as WPW syndrome.

### **ADENOSINE (ADENOCARD) (Cont.)**

**CONTRAINDICATIONS:** Second- or third-degree heart block, sick sinus syndrome, known hypersensitivity.

**SIDE EFFECTS:** Facial flushing, headache, chest pain, SOB, dizziness, and nausea.

**DRUG INTERACTIONS:** Methylxantines (eg Aminophylline, Theophylline) may decrease the effectiveness, thus requiring larger doses. Dipyridamole (Persantine, Aggrenox) can potentiate the effects, thus dosage may need to be reduced.

**ROUTE:** Rapid IV push.

**DOSAGE:**

**ADULTS:** Initial dose: 6 mg rapid IV/IO, followed by 20 ml flush; elevate extremity

Additional dose: 12 mg rapid IV/IO if PSVT continues, may repeat x1

Unstable initial dose if IV established: 12 mg rapid IV/IO

**PEDS:** Initial dose: 0.1 mg/kg (max 6 mg) rapid IV/IO, followed by 5-10 ml flush; elevate extremity

Additional dose: 0.2 mg/kg (max 12 mg) rapid IV/IO

**PREGNANCY SAFETY:** Category C – give only if potential benefits justifies risk to fetus.

**COMMENTS:** Due to extremely short half-life, adenosine must be administered rapid IV push, preferable via large bore IV closest to central circulation as possible (ie AC, EJ).

### **ALBUTEROL**

**PROVIDER LEVEL:** EMT, AEMT, Paramedic.

**CLASS:** Sympathomimetic, bronchodilator,  $\beta_2$ -agonist.

**PHARMACOLOGY/ACTIONS:** Albuterol is a sympathomimetic that is selective for  $\beta_2$ -adrenergic receptors. It relaxes smooth muscles of the bronchial tree and peripheral vasculature by stimulating adrenergic receptors of the sympathetic nervous system.

**ONSET/DURATION:** Onset: 5-15 minutes / Duration: 3-4 hours.

**INDICATIONS:** Relief of bronchospasm in patients with reversible obstructive airway disease (ie acute asthma exacerbation, acute COPD exacerbation, anaphylactic reaction), drowning, and treatment of hyperkalemia.

**CONTRAINDICATIONS:** Hypersensitivity, cardiac dysrhythmias associated with tachycardia.

**SIDE EFFECTS:** Restlessness, apprehension, dizziness, palpitations, tachycardia, dysrhythmias.

**DRUG INTERACTIONS:** Other sympathomimetics may exacerbate adverse cardiovascular effects. Antidepressants may potentiate effects on the vasculature (vasodilation).  $\beta$ -blockers may antagonize albuterol. May potentiate diuretic-induced hypokalemia.

**ROUTE:** Inhalation.

**DOSAGE:**

**ADULTS:** MDI: 1-2 puffs (90-180 mcg) every 4 hr, use of spacer is preferred; Nebulizer: 2.5 mg in 3 ml NS administer over 5-15 min, may repeat as needed.

Hyperkalemia: 5-10 mg in 3 ml NS continuous SVN.

**PEDS:** <10 kg 1.25 mg in 3 ml NS, >10 kg 2.5 mg in 3 ml NS; may repeat as needed.

**PREGNANCY SAFETY:** Category C – give only if potential benefits justifies risk to fetus.

**COMMENTS:** Consider use in combination with Ipratropium (Atrovent). May precipitate angina pectoris and dysrhythmias. Use in caution in patients with diabetes mellitus, hyperthyroidism, prostatic hypertrophy, seizure disorder, or cardiovascular disorders. Spacer use is desirable with meter dose inhalers as it will increase drug delivery.

### **AMIODARONE (CORDARONE)**

**PROVIDER LEVEL:** Paramedic.

**CLASS:** Class III antidysrhythmic agent (with multiple other class properties).

**PHARMACOLOGY/ACTIONS:** Prolongs the action potential duration and effective refractory period, and when given short-term IV, probably induces noncompetitive  $\beta$ -adrenoreceptor and calcium channel blocker activity. The IV formulation relaxes vascular smooth muscle, decreases peripheral vascular resistance, and increases coronary blood flow.

**ONSET/DURATION:** Onset: within minutes / Duration: varies.

**INDICATIONS:**

- Ventricular fibrillation (V-fib)/pulseless ventricular tachycardia (V-tach).
- Hemodynamically unstable V-tach with a pulse.
- Other atrial and ventricular dysrhythmias.

**CONTRAINDICATIONS:** No contraindications if in cardiac arrest or unstable dysrhythmia. Relative: pulmonary congestion, cardiogenic shock, hypotension, bradycardia, advanced AV block, known hypersensitivity, and sick sinus syndrome. Caution in patients with severe liver disease.

### **AMIODARONE (CORDARONE) (Cont.)**

**SIDE EFFECTS:** Hypotension, headache, dizziness, bradycardia, AV conduction abnormalities, flushing, and abnormal salivation.

**DRUG INTERACTIONS:** Multiple complex drug interactions.

**ROUTE:** IV, IO.

**DOSAGE:**

**ADULTS:** Cardiac Arrest: Initial: 300 mg rapid IV bolus. Additional: 150 mg in 3-5 min if needed.

Wide Complex Tachycardia: 150 mg in 50 ml NS over 10 minutes.

**PEDS:** Cardiac Arrest: Initial: 5 mg/kg (max 300 mg) rapid IV bolus, may repeat x2.

Wide Complex Tachycardia: 5 mg/kg (max 150 mg) IV/IO over 20 minutes.

**PREGNANCY SAFETY:** Category D – positive risk to fetus, maternal benefit may outweigh risk to fetus.

**COMMENTS:** Continuous ECG monitoring is required. Slow infusion or discontinue if bradycardia or AV blocks occur. Maintain drug at room temperature and protect from excessive heat. Administer 150 mg in 100 ml over 10 minutes if ventricular cardiac arrest with return of spontaneous circulation (ROSC) and did not previously push amiodarone as part of ACLS v-fib/pulseless v-tach algorithm.

### **ASPIRIN (ASA)**

**PROVIDER LEVEL:** EMT, AEMT, Paramedic.

**CLASS:** Platelet aggregator inhibitor and anti-inflammatory agent.

**PHARMACOLOGY/ACTIONS:** Aspirin blocks the formation of the substance thromboxane A2, which causes platelets to aggregate and arteries to constrict.

**ONSET/DURATION:** Onset: 15-30 minutes / Duration: 4-6 hours.

**INDICATIONS:** Chest pain suggestive of acute coronary syndrome (ACS).

**CONTRAINDICATIONS:** Absolute: known hypersensitivity, known hemorrhagic stroke. Relative: GI bleeding, bleeding disorders, active ulcer disease, aortic aneurysm, asthma.

**SIDE EFFECTS:** Heartburn, GI bleeding, nausea, vomiting, wheezing, and prolonged bleeding.

**DRUG INTERACTIONS:** Other anti-inflammatory agents; ↓ effects with antacids and steroids; ↑ effects with anticoagulants, insulin, oral hypoglycemics, fibrinolytic agents.

**ROUTE:** PO – chewable.

**DOSAGE:**

**ADULTS:** Chew 324 mg (81 mg x 4). If patient has taken at least 324 mg in past 12 hours, do not give additional dose. If patient has taken less than 324 mg – give additional dose to supplement total dose to 324 mg.

**PREGNANCY SAFETY:** Category D – positive risk to fetus, maternal benefit may outweigh risk to fetus.

**COMMENTS:** Do not substitute acetaminophen or ibuprofen.

### **ATROPINE**

**PROVIDER LEVEL:** Paramedic.

**CLASS:** Anticholinergic (parasympatholytic).

**PHARMACOLOGY/ACTIONS:** Atropine acts by blocking acetylcholine receptors, thus inhibiting parasympathetic stimulation. Antagonizes excess muscarinic receptor stimulation caused by organophosphate insecticides or chemical nerve agents (eg sarin).

**ONSET/DURATION:** Onset: immediate / Duration: 2-6 hours

**INDICATIONS:** Hemodynamically significant bradycardia, organophosphate or nerve gas poisoning (large doses usually required).

**CONTRAINDICATIONS:** Tachycardia, hypersensitivity.

**SIDE EFFECTS:** Blurred vision, dilated pupils, dry mouth, tachycardia, drowsiness, and confusion.

**DRUG INTERACTIONS:** There are few interactions in the prehospital setting.

**ROUTE:** IV, IO.

**DOSAGE:**

**ADULTS:** Symptomatic Bradycardia: 1 mg IV/IO. Repeat every 3-5 min for a maximum dose of 0.04 mg/kg or 3 mg.

Anticholinesterase Poisoning: 1-2 mg IV/IO every 5-15 min until atropine effects are observed.

**PEDS:** Symptomatic Bradycardia: 0.02 mg/kg IV/IO (minimum 0.1 mg, maximum 0.5 mg). May repeat once after 5 min.

Anticholinesterase Poisoning: 0.02 mg/kg IV/IO (minimum 0.1 mg, maximum 0.5 mg). Repeat every 15 min until atropine effects are observed.

**PREGNANCY SAFETY:** Category C – give only if potential benefits justifies risk to fetus.

### **ATROPINE (Cont.)**

**COMMENTS:** May worsen bradycardia associated with second-degree Mobitz II and third-degree AV blocks. A maximum dose of 0.04 mg/kg or 3 mg should not be exceeded except in the case of organophosphate poisoning.

### **BRILINTA (generic name TICAGRELOR)**

**PROVIDER LEVEL:** Paramedic.

**CLASS:** Antiplatelet; P2Y12 inhibitor.

**PHARMACOLOGY/ACTIONS:** Reversibly inhibits the P2Y12 receptor on the surface of platelets which prevents activation of the GPIIb/IIIa receptor complex reducing platelet aggregation.

**ONSET/DURATION:** Onset: about 30 minutes / Duration: 2-8 hours after loading dose.

**INDICATIONS:** STEMI.

**CONTRAINDICATIONS:** Hypersensitivity, active pathological bleeding, and history of intracranial hemorrhage (ICH). Relative: Bleeding disorders, heparin induced thrombocytopenia (HIT), hepatic impairment, and renal impairment.

**SIDE EFFECTS:** Dyspnea, nausea, hemorrhage, dizziness, and ventricular pause.

**DRUG INTERACTIONS:** Antiplatelet therapies, anticoagulants, antifibrinolytics, and strong CYP3A4 inhibitors.

**ROUTE:** PO.

**DOSAGE:**

**ADULTS:** 180 mg PO.

**PREGNANCY SAFETY:** Not recommended.

**COMMENTS:** May be administered by BSFD, YMCERFD, and HBRFD **only** when bypassing BSMC and transporting (ground or flight) directly to hospital with cardiac catheterization capability (i.e. BHDRMC, EIRMC) **and** once physician confirms STEMI on 12-Lead ECG.

### **CALCIUM**

**PROVIDER LEVEL:** Paramedic.

**CLASS:** Electrolyte.

**PHARMACOLOGY/ACTIONS:** Calcium chloride or calcium gluconate provides elemental calcium in the form of the cation  $\text{Ca}^{2+}$  which is necessary for many physiologic activities. Calcium causes a significant increase in myocardial contractility. Calcium also increases cardiac muscle tone and force of systolic contractions (positive inotropic effect) making it especially useful for patients with sympathetic blockade.

**ONSET/DURATION:** Onset: immediate / Duration: unknown.

**INDICATIONS:**

- Hyperkalemia.
- Calcium channel blocker overdose with hypotension and bradycardia.
- Toxicity from magnesium sulfate overdose.
- Tetany associated with black widow spider bite.
- Hydrofluoric acid burns.

**CONTRAINDICATIONS:** Digitalis toxicity, ventricular fibrillation, and hypercalcemia.

**SIDE EFFECTS:** Hypotension, bradycardia, arrhythmias, syncope, cardiac arrest, and tissue irritation.

**DRUG INTERACTIONS:** Will interact with sodium bicarbonate forming a precipitate.

**ROUTE:** IV, IO, topical.

**DOSAGE:**

**ADULTS:** Hyperkalemia: Calcium chloride 1 g IV/IO over 5 min or calcium gluconate 2 g IV/IO over 5 min.

Magnesium toxicity: Calcium Chloride 1 g IV or calcium gluconate 3 g IV.

CCB OD: calcium chloride 0.5-1 g slow IV/IO (50 mg/min) or calcium gluconate 2-6 g slow IV/IO over 10 min.

**PEDS:** CCB OD: calcium chloride (10%) 0.2 ml/kg slow IV/IO.

**PREGNANCY SAFETY:** Category C – give only if potential benefits justifies risk to fetus.

**COMMENTS:** Calcium is no longer recommended for routine administration during cardiac arrest. Calcium should be administered slowly through a patent IV in a large vein to avoid possibility of extravasation and resultant tissue necrosis.

### **DEXTROSE**

**PROVIDER LEVEL:** AEMT, Paramedic.

**CLASS:** Carbohydrate.

**DEXTROSE (Cont.)**

**PHARMACOLOGY/ACTIONS:** The term dextrose is used to describe the six-carbon sugar d-glucose, the principal form of carbohydrate used by the body. 50% dextrose solution is used in emergency care to treat hypoglycemia and in the management of coma of unknown origin.

**ONSET/DURATION:** Onset: 1 minute / Duration: dependent.

**INDICATIONS:** - Altered mental status of unknown origin.

- Hypoglycemia.
- Seizures.
- Head trauma with decreased mental status if unable to check blood glucose.
- Hypothermia if unable to take oral nutrition safely (prolonged field setting).

**CONTRAINDICATIONS:** None if documented or suspected hypoglycemia. Intracranial hemorrhage, ICP, and known or suspected CVA in the absence of hypoglycemia.

**SIDE EFFECTS:** Warmth, pain, burning from medication infusion, hyperglycemia, and thrombophlebitis.

**DRUG INTERACTIONS:** None significant.

**ROUTE:** IV, IO.

**DOSAGE:**

**ADULTS:** D10: 25 g (250 ml of 10% solution) IV/IO, may repeat x1 in 5 min.

D50: 12.5-25 g IV/IO, may repeat x1 in 5 min.

**PEDS:** D10: 5 ml/kg (max 25 g) IV/IO

**PREGNANCY SAFETY:** Category C – give only if potential benefits justifies risk to fetus.

**COMMENTS:** An infusion of dextrose 10% (D10) is preferred over dextrose 50% (D50). D10 is more physiologic (D50 is hypertonic). D10 results in lower post-treatment blood glucose levels. Extravasation may cause tissue necrosis, use large vein and aspirate occasionally to ensure route patency. Draw blood sample and check BGL prior to administration if possible.

**DIAZEPAM (VALIUM)**

**PROVIDER LEVEL:** Paramedic.

**CLASS:** Benzodiazepine.

**PHARMACOLOGY/ACTIONS:** Acts on the limbic, thalamic, and hypothalamic regions of the CNS to potentiate the effects of inhibitory (GABA) neurotransmitters, raising the seizure threshold in the motor cortex.

**ONSET/DURATION:** Onset: 1-5 minutes IV / Duration: 15 minutes – 1 hour.

**INDICATIONS:** - Seizure activity.  
- Acute anxiety states.  
- Premedication before cardioversion or TCP.  
- Acute alcohol withdrawal.  
- Skeletal muscle relaxation.

**CONTRAINDICATIONS:** Hypersensitivity, shock, coma, CNS depression due to head injury, and respiratory depression. Use with caution in substance abuse patients.

**SIDE EFFECTS:** Hypotension, respiratory depression, ataxia, psychomotor impairments, confusion, and nausea.

**DRUG INTERACTIONS:** May precipitate CNS depression and psychomotor impairment in patients who are taking other CNS depressant medications. Diazepam should not be administered with other drugs because of possible precipitation.

**ROUTE:** IV, IO, Rectal.

**DOSAGE:**

**ADULTS:** Seizures: 0.2 mg/kg IV (max 10 mg), may repeat x 1 after 5 min.

Sedation/Pain: 5 mg IV.

**PEDS:** Seizures: 0.2 mg/kg IV (max 10 mg), may repeat x 1 after 5 min.

**PREGNANCY SAFETY:** Category D – positive risk to fetus, maternal benefit may outweigh risk to fetus.

**COMMENTS:** DEA schedule IV drug with potential for abuse. May cause local venous irritation. Has short duration of anticonvulsant effect. Reduce dose by 50% in elderly patients or with known liver disease. Resuscitation equipment should be readily available.

**DIPHENHYDRAMINE (BENADRYL)**

**PROVIDER LEVEL:** EMT w/medication endorsement and AEMT (PO only), Paramedic.

**CLASS:** Antihistamine.



### **DIPHENHYDRAMINE (BENADRYL) (Cont.)**

**PHARMACOLOGY/ACTIONS:** Antihistamines prevent the physiological actions of histamines by blocking H1 and H2 receptor sites. It may inhibit MAST cell damage, preventing more histamine release. It also has anticholinergic and antiemetic effects.

**ONSET/DURATION:** Onset: maximum effects 1-3 hours / Duration: 6-12 hours.

**INDICATIONS:**

- Allergic reactions.
- Anaphylaxis.
- Acute extrapyramidal (dystonic) reactions.
- Nausea/vomiting (related to motion sickness and other causes).

**CONTRAINDICATIONS:** Hypersensitivity, patients taking MAO inhibitors, narrow-angle glaucoma (relative), newborns and nursing mothers.

**SIDE EFFECTS:** Drowsiness, disturbed coordination, hypotension, palpitations, tachycardia, bradycardia, thickening of bronchial secretions, dry mouth and throat, paradoxical excitement in children.

**DRUG INTERACTIONS:** CNS depressants may increase depressant effects, MAO inhibitors may prolong and intensify anticholinergic effects.

**ROUTE:** PO, IV, IO, IM.

**DOSAGE:**

**ADULTS:** 25-50 mg PO/IM, slow IV/IO.

**PEDS:** 1 mg/kg (max 50 mg) PO/IM, slow IV/IO.

**PREGNANCY SAFETY:** Category C – give only if potential benefits justifies risk to fetus.

**COMMENTS:** Use with caution in patients with CNS depression or lower respiratory disease such as asthma. Intramuscular epinephrine is the first line agent for treatment of allergic reactions/anaphylaxis, whereas diphenhydramine is a second line agent.

### **DROPERIDOL (INAPSINE)**

**PROVIDER LEVEL:** AEMT (antiemetic only) and Paramedic.

**Class:** First generation antipsychotic.

**PHARMACOLOGY/ACTIONS:** Droperidol is a butyrophenone antipsychotic. Antiemetic effect is a result of blockade of dopamine stimulation of the chemoreceptor trigger zone. Other effects include alpha-adrenergic blockade, peripheral vascular dilation, and reduction of the pressor effect of epinephrine resulting in hypotension and decreased peripheral vascular resistance. May also reduce pulmonary artery pressure.

**ONSET/DURATION:** Onset: 3-10 minutes / Duration: 2-4 hours.

**INDICATIONS:**

- Nausea/vomiting.
- Migraine headache.
- Nonspecific abdominal pain.
- Chemical sedation.

**CONTRAINDICATIONS:** Hypersensitivity and QT prolongation (Black Box Warning). Relative: anticholinergic effects, CNS depression, esophageal dysmotility, extrapyramidal syndrome, and neuroleptic malignant syndrome.

**SIDE EFFECTS:** QT prolongation, hypotension, ventricular tachycardia, drowsiness, dysphoria, extrapyramidal reaction - akathisia, hallucination, hyperactive behavior, neuroleptic malignant syndrome, laryngospasm, bronchospasm, and respiratory depression.

**DRUG INTERACTIONS:** Anti-parkinson agents (dopamine agonists) and CNS depressants.

**ROUTE:** IM, IV.

**DOSAGE:**

**ADULTS:** Nausea/vomiting, headache, nonspecific abdominal pain: 0.625-1.25 mg IV/IM.

Chemical sedation: 2.5 mg IV or 5-10 mg IM.

**PREGNANCY SAFETY:** Category C – give only if potential benefits justifies risk to fetus.

**COMMENTS:** Continuous cardiac monitoring recommended given potential for QT prolongation. Use with caution in patients with bradycardia, cardiac disease, concurrent MAO inhibitor therapy, Class I and Class III dysrhythmics or other drugs that prolong the QT interval and cause electrolyte disturbances due to its adverse cardiovascular effects, e.g., QT prolongation, hypotension, tachycardia, and torsades de pointes. Dilute and push very slowly or place in bag of IV fluid and infuse to reduce risk of akathisia.

### **EPINEPHRINE 1:1,000**

**PROVIDER LEVEL:** EMR (only patient prescribed auto-injector), EMT (agency supplied auto-injector), EMT w/medication endorsement (IM), AEMT and Paramedic.

**CLASS:** Sympathetic agonist (sympathomimetic).

**PHARMACOLOGY/ACTIONS:** Epinephrine acts directly on  $\alpha$ - and  $\beta$ -adrenergic receptors. Its effect on  $\beta$ -receptors is much more profound than its effect on  $\alpha$ -receptors. Effects include:  $\uparrow$  HR,  $\uparrow$  cardiac contractile force,  $\uparrow$  electrical activity in the myocardium,  $\uparrow$  systemic vascular resistances,  $\uparrow$  BP, and  $\uparrow$  automaticity.

**ONSET/DURATION:** Onset: < 2 minutes / Duration: 5-10 minutes.

**INDICATIONS:**

- Anaphylaxis/Allergic reaction.
- Severe asthma.

**CONTRAINDICATIONS:** Hypovolemic shock, severe hypertension. Relative: coronary artery disease, pregnancy, severe tachydysrhythmias, PVC's, hyperthyroidism, and cerebrovascular insufficiency.

**SIDE EFFECTS:** Tachycardia including V-tach and V-fib, palpitations, anxiety, tremor, headache, nausea, weakness, restlessness, and hypertension.

**DRUG INTERACTIONS:** MAO inhibitors may potentiate effects.  $\beta$ -adrenergic antagonists may blunt inotropic response. Sympathomimetics and phosphodiesterase inhibitors may exacerbate dysrhythmia response.

**ROUTE:** IM.

**DOSAGE:**

**ADULTS:** Auto-injector: 0.3 mg IM. Asthma/allergic reaction: 0.3-0.5 mg IM, may repeat every 15 min.

**PEDS:** Auto-injector: 0.15 mg IM. Asthma/allergic reaction: 0.01 mg/kg IM (max 0.3 mg), may repeat every 15 min.

**PREGNANCY SAFETY:** Category C – give only if potential benefits justifies risk to fetus.

**COMMENTS:** Epinephrine 1:1000 should only be given intramuscular (IM), NEVER intravenous (IV). Be extremely cautious with dosage calculations and administration. Use with caution in patients with peripheral vascular insufficiency. Epinephrine is pH dependent and can be deactivated by alkaline solutions such as sodium bicarbonate. Effects can be intensified in patients who are taking antidepressants. Intramuscular epinephrine is the first line agent for treatment of allergic reactions/anaphylaxis. Use with caution in patients over 50 years old due to cardiovascular stress, however do not withhold for severe allergic reaction/anaphylaxis.

### **EPINEPHRINE 1:10,000**

**PROVIDER LEVEL:** Paramedic.

**CLASS:** Sympathetic agonist (sympathomimetic).

**PHARMACOLOGY/ACTIONS:** Epinephrine acts directly on  $\alpha$ - and  $\beta$ -adrenergic receptors. Its effect on  $\beta$ -receptors is much more profound than its effect on  $\alpha$ -receptors. Effects include:  $\uparrow$  HR,  $\uparrow$  cardiac contractile force,  $\uparrow$  electrical activity in the myocardium,  $\uparrow$  systemic vascular resistances,  $\uparrow$  BP, and  $\uparrow$  automaticity.

**ONSET/DURATION:** Onset: < 2 minutes / Duration: 5-10 minutes.

**INDICATIONS:**

- Cardiac arrest (asystole, v-fib, pulseless v-tach, PEA).
- Anaphylaxis.
- Unstable bradycardia.

**CONTRAINDICATIONS:** None in cardiac arrest.

**SIDE EFFECTS:** Tachycardia including V-tach and V-fib, palpitations, anxiety, tremor, headache, nausea, weakness, restlessness, and hypertension.

**DRUG INTERACTIONS:** MAO inhibitors may potentiate effects.  $\beta$ -adrenergic antagonists may blunt inotropic response. Sympathomimetics and phosphodiesterase inhibitors may exacerbate dysrhythmia response.

**ROUTE:** IV, IO.

**DOSAGE:**

**ADULTS:** Cardiac arrest: 1 mg IV/IO push every 3-5 min.

Anaphylaxis: 10 mcg IV every 2-5 min and/or 0.5 mcg/kg/min IV infusion.

Unstable bradycardia: 10 mcg IV every 2-5 min and/or 0.02-0.2 mcg/kg/min IV infusion.

Shock: 10 mcg IV every 3-5 min and/or 0.05-0.3 mcg/kg/min IV infusion.

**PEDS:** Cardiac arrest: 0.01 mg/kg (max 1 mg) IV/IO push every 3-5 min.

Anaphylaxis: 0.1 mcg/kg (max 5 mcg) IV and/or 0.5 mcg/kg/min IV infusion.

Unstable bradycardia: 10 mcg IV, may repeat every 3-5 min.

Shock: 10 mcg IV every 3-5 min and/or 0.05-0.3 mcg/kg/min IV infusion.

**PREGNANCY SAFETY:** Category C – give only if potential benefits justifies risk to fetus.

### **EPINEPHRINE 1:10,000 (Cont.)**

**COMMENTS:** Epinephrine is pH dependent and can be deactivated by alkaline solutions such as sodium bicarbonate. Effects can be intensified in patients who are taking antidepressants. Epinephrine use in cardiac arrest has been shown to increase ROSC, but has not been shown to increase neurologically intact survival. Some studies have shown a correlation between increased epinephrine use and worse neurological outcome. While there is not clear definitive data to support an optimal dose of epinephrine, there is data supporting possible harm from high end dosing. Epinephrine should be limited to no more than 3 mg in cardiac arrest.

### **FENTANYL (SUBLIMAZE)**

**PROVIDER LEVEL:** Paramedic.

**CLASS:** Synthetic opioid agonist.

**PHARMACOLOGY/ACTIONS:** Analgesic with short duration of action. Minimal histamine release, so less hemodynamic compromise.

**ONSET/DURATION:** Onset: 5-8 minutes / Duration: 1-2 hours.

**INDICATIONS:**

- Pain control.
- Cardiac suspected chest pain **after** administration of oxygen, aspirin, and nitroglycerin according to guideline.
- Sedation for invasive procedures (TCP/cardioversion).

**CONTRAINDICATIONS:** Respiratory depression or insufficiency, uncorrected hypotension, and hypersensitivity.

**SIDE EFFECTS:** Respiratory depression, bradycardia, hypotension, hypertension, nausea and vomiting.

**DRUG INTERACTIONS:** Effects may be increased when given with other CNS depressants or skeletal muscle relaxants.

**ROUTE:** IV, IO, IM, IN.

**DOSAGE:**

**ADULTS:** Pain: 1 mcg/kg (max 100 mcg) IN/IM/IV/IO, may repeat dose after 10 min.

Cardioversion or TCP: 1 mcg/kg IN/IM/IV/IO to a maximum single dose of 100 mcg.

**PEDS:** 1 mcg/kg (max 50 mcg) IN/IM/IV/IO, may repeat dose after 10 min.

**PREGNANCY SAFETY:** Category C – give only if potential benefits justifies risk to fetus.

**COMMENTS:** DEA schedule II drug with potential for abuse. Fentanyl should be used with caution in elderly patients and in those with severe respiratory disorders, seizure disorders, cardiac disorders, or pregnancy. Rapid administration or large doses may cause skeletal muscle (chest wall) rigidity so severe that ventilation is difficult or impossible.

### **GLUCAGON**

**PROVIDER LEVEL:** EMT w/medication endorsement and AEMT (hypoglycemic reaction only), Paramedic.

**CLASS:** Pancreatic hormone, insulin antagonist.

**PHARMACOLOGY/ACTIONS:** Glucagon is a protein secreted by the alpha cells of the pancreas. It increases blood glucose by converting glycogen in the liver into glucose. Glucagon also has positive inotropic action on the heart and decreases renal vascular resistance which makes it useful in beta-blocker and calcium channel blocker overdose.

**ONSET/DURATION:** Onset: 1 minute / Duration: 60-90 minutes.

**INDICATIONS:**

- Persistent hypoglycemia despite glucose supplementation.
- Hypoglycemic patient where unable to establish IV access.
- Beta-blocker or calcium channel blocker toxicity.

**CONTRAINDICATIONS:** Hypersensitivity (allergy to proteins). Relative is patient with no glycogen storage (malnutrition, alcoholism).

**SIDE EFFECTS:** Tachycardia, hypotension, urticaria, nausea, and vomiting.

**DRUG INTERACTIONS:** Effects of anticoagulants may be increased if given with glucagon. Do not mix with saline.

**HOW SUPPLIED:** Glucagon must be reconstituted with provided diluent before administration. Dilute 1 unit (1 mg) white powder in 1 ml of diluting solution (1 mg/ml).

**ROUTE:** IV, IO, IM, IN.

**DOSAGE:**

**ADULTS:** Hypoglycemia: 1 mg IM/IN.

β-blocker or CCB toxicity: 5 mg IV/IO, may repeat in 5-10 min, max 10 mg.

**PEDS:** Hypoglycemia: <20 kg = 0.5 mg IM/IN, >20 kg = 1 mg IM/IN.

β-blocker and CCB toxicity: 50-150 mcg/kg IV/IO.

**PREGNANCY SAFETY:** Category B – unproven or unknown risk to fetus, and no risk in later trimesters.

### **GLUCAGON (Cont.)**

**COMMENTS:** Glucagon should not be considered a first-line choice for hypoglycemia. IV glucose will need to be administered if the patient does not respond to a second dose of glucagon. Do not give more than 2 doses of Glucagon as maximal glycogen release from the liver has occurred.

### **HALOPERIDOL (HALDOL)**

**PROVIDER LEVEL:** Paramedic.

**CLASS:** Antipsychotic/neuroleptic.

**PHARMACOLOGY/ACTIONS:** The drug is thought to block dopamine (type 2) receptors in the brain, altering mood and behavior.

**ONSET/DURATION:** Onset: 10-60 minutes / Duration: 12-24 hours.

**INDICATIONS:**

- Acute psychotic episodes.
- Emergency sedation of severely agitated, aggressive, or delirious patients who present a danger to themselves or others.
- Cannabinoid hyperemesis syndrome.

**CONTRAINDICATIONS:** Hypersensitivity, CNS depression, combativeness from trauma, pregnancy, severe liver or cardiac disease.

**SIDE EFFECTS:** Dose-related extrapyramidal reactions: pseudoparkinsonism, akathisia, dystonias, orthostatic hypotension, allergic reactions, nausea and vomiting, blurred vision.

**DRUG INTERACTIONS:** Other CNS depressants may potentiate effects. May inhibit vasoconstrictor effects of epinephrine.

**ROUTE:** IM, IV.

**DOSAGE:**

**ADULTS:** Behavioral emergency: 5-10 mg IM or 5 mg IV.

Cannabinoid hyperemesis syndrome: 2.5-5 mg IV/IM.

**PREGNANCY SAFETY:** Category C – give only if potential benefits justifies risk to fetus.

**COMMENTS:** Do not use in patients with suspected head injury. Administer diphenhydramine for patients with dystonic reaction. Fluid challenge is indicated with significant drop in blood pressure. Cardiac monitoring for prolongation of QT interval required with IV administration.

### **HEPARIN**

**PROVIDER LEVEL:** Paramedic.

**CLASS:** Anticoagulant.

**PHARMACOLOGY/ACTIONS:** Potentiates the action of antithrombin III and thereby inactivates thrombin (II) and prevents conversion of fibrinogen to fibrin.

**ONSET/DURATION:** Onset: immediate / Duration: dose dependent 30-150 minutes.

**INDICATIONS:** STEMI.

**CONTRAINDICATIONS:** Hypersensitivity, thrombocytopenia, history of heparin induced thrombocytopenia (HIT), and uncontrolled active bleeding.

**SIDE EFFECTS:** Bleeding and thrombosis due to HIT.

**DRUG INTERACTIONS:** Antifibrinolytics, antiplatelets, and anticoagulants.

**ROUTE:** IV.

**DOSAGE:**

**ADULTS:** 60 units/kg (max 4000 units) IV bolus.

**PREGNANCY SAFETY:** may be used (enoxaparin preferred).

**COMMENTS:** May be administered by BSFD, YMCERFD, and HBRFD **only** when bypassing BSMC and transporting (ground or flight) directly to hospital with cardiac catheterization capability (i.e. BHDRMC, EIRMC) **and** once physician confirms STEMI on 12-Lead ECG.

### **HYDROXOCOBALAMIN (CYANOKIT)**

**PROVIDER LEVEL:** Paramedic.

**CLASS:** Cyanide antidote; vitamin.

**PHARMACOLOGY:** In the presence of cyanide, each hydroxocobalamin molecule can bind one cyanide ion by displacing it for the hydroxo ligand linked to the trivalent cobalt ion, forming cyanocobalamin, which is then excreted in the urine.

**ONSET/DURATION:** Onset: 30 minutes / Duration: 24 hours.

### **HYDROXOCOBALAMIN (CYANOKIT) (Cont.)**

**INDICATIONS:** Cyanide poisoning / smoke inhalation.

**CONTRAINDICATIONS:** Hypersensitivity.

**SIDE EFFECTS:** Rash, diarrhea, urine discoloration, infusion site reaction, and headache.

**DRUG INTERACTIONS:** Chloramphenicol.

**ROUTE:** IV, IO.

**DOSAGE:**

**ADULTS:** 5 g IV/IO over 15 min.

**PEDS:** refer to dosing chart under Pediatric Smoke Inhalation Guideline.

**PREGNANCY SAFETY:** Treatment not withheld due to pregnancy.

**COMMENTS:** Will cause discoloration of the skin and urine, can interfere with pulse oximetry. Due to its interference with certain diagnostic blood tests, the performance of prehospital phlebotomy is preferable prior to the administration of hydroxocobalamin.

### **IBUPROFEN (MOTRIN, ADVIL)**

**PROVIDER LEVEL:** EMT, AEMT, Paramedic.

**CLASS:** Non-steroidal anti-inflammatory drug (NSAID).

**PHARMACOLOGY/ACTIONS:** Inhibits synthesis of prostaglandins in body tissues by inhibiting at least 2 cyclo-oxygenase (COX) isoenzymes, COX-1 and COX-2. May inhibit chemotaxis, alter lymphocyte activity, decrease proinflammatory cytokine activity, and inhibit neutrophil aggregation; these effects may contribute to anti-inflammatory activity.

**ONSET/DURATION:** Onset: 30 minutes / Duration: 4-6 hours.

**INDICATIONS:**

- Mild/moderate pain.
- Antipyretic.
- Acute Mountain Sickness.

**CONTRAINDICATIONS:** Aspirin allergy, perioperative pain in setting of coronary artery bypass graft (CABG) surgery, preterm infants with untreated proven or suspected infection, bleeding with active intracranial hemorrhage or GI bleed, thrombocytopenia, coagulation defects, proven or suspected necrotizing enterocolitis, significant renal impairment, congenital heart disease where patency or the patent ductus arteriosus (PDA) is necessary for pulmonary or systemic blood flow.

**SIDE EFFECTS:** Acute kidney injury, stomach upset, nausea, vomiting, dizziness, blurred vision, dry mouth, and irritation at the injection site.

**DRUG INTERACTIONS:** Other NSAIDs, steroids, aspirin, and anticoagulants (ex Coumadin, Xarelto, Eliquis).

**ROUTE:** PO.

**DOSAGE:**

**ADULTS:** 200-800 mg PO.

**PEDS:** 10 mg/kg (max 800 mg) PO.

**PREGNANCY SAFETY:** Category C – give only if potential benefits justifies risk to fetus. Avoid in pregnancy after 20 weeks gestation.

**COMMENTS:** Use with caution in patients who are dehydrated or taking ACEIs or ARBs. Ensure patient has not taken oral NSAID in last 6 hours (ibuprofen, Motrin), 12 hours (naproxen), or 24 hours (meloxicam) given potential for cumulative risk of inducing serious NSAID-related side effects.

### **IPRATROPIUM BROMIDE (ATROVENT)**

**PROVIDER LEVEL:** EMT, AEMT, Paramedic.

**CLASS:**  $\beta_2$ -agonist.

**PHARMACOLOGY/ACTIONS:** Ipratropium is an anticholinergic bronchodilator which is chemically related to atropine. Ipratropium is a parasympatholytic used in the treatment of respiratory emergencies. It causes bronchodilation and dries respiratory tract secretions. Ipratropium acts by blocking acetylcholine receptors, thus inhibiting parasympathetic stimulation.

**ONSET/DURATION:** Onset: 5-15 minutes / Duration: 4-6 hours.

**INDICATIONS:** Treatment of bronchial asthma, reversible bronchospasm associated with chronic bronchitis or emphysema.

**CONTRAINDICATIONS:** Known hypersensitivity to atropine or its derivatives. Not indicated as a single agent for acute treatment of bronchospasm where rapid response is required.

**SIDE EFFECTS:** Tachycardia, paradoxical bronchospasm may occur.

**DRUG INTERACTIONS:** None reported.

### **IPRATROPIUM BROMIDE (ATROVENT) (Cont.)**

**ROUTE:** Inhalation.

**DOSAGE:**

**ADULTS:** 500 mcg SVN once or DuoNeb 3 ml SVN once.

**PEDS:** > 2 yo with h/o wheezing 2.5 ml 0.02% solution SVN once or DuoNeb 3 ml SVN once.

**PREGNANCY SAFETY:** Category B – unproven or unknown risk to fetus, and no risk in later trimesters.

**COMMENTS:** Generally used in conjunction with first Albuterol dose, subsequent nebulizers should be with Albuterol only. Vital signs must be monitored during therapy with ipratropium. Caution should be used when administering to elderly patients and those with cardiovascular disease and hypertension. Use caution in patients with significant tachycardia (120+), prostatic hypertrophy, narrow angle glaucoma or bladder neck obstruction.

### **KETAMINE (KETALAR)**

**PROVIDER LEVEL:** Paramedic.

**CLASS:** NMDA antagonist.

**PHARMACOLOGY/ACTIONS:** Ketamine is a rapid-acting, general anesthetic producing an anesthetic state characterized by profound analgesia, amnesia, normal pharyngeal-laryngeal reflexes, normal or slightly enhanced skeletal muscle tone, cardiovascular and respiratory stimulation, and occasionally a transient and minimal respiratory depression. Ketamine bronchodilation relaxes bronchiolar musculature and prevents bronchoconstriction induced by histamine.

**ONSET/DURATION:** Onset: 30-60 seconds / Duration: 15 minutes.

**INDICATIONS:**

- Pain of traumatic origin.
- Opioid-tolerant patient with an acute exacerbation of pain.
- Pain that is refractory to opioids.
- Extreme agitation and/or combativeness.

**CONTRAINDICATIONS:** Hypersensitivity, chest pain of suspected cardiac origin, hypertensive crisis, amphetamine abuse, acute pulmonary edema, pregnancy, and history of psychiatric disorder (relative contraindication).

**SIDE EFFECTS:** Hallucinations, disorientation, delirium, agitation, dizziness, diplopia, dysphoria, rotary nystagmus, nausea, vomiting, hypertension, tachycardia, laryngospasms, and hypersalivation.

**DRUG INTERACTIONS:** Azelastine, Bromperidol, Flunarizine, Kratom, Olopatadine, Orphenadrine, Oxememazine, and Thalidomide.

**ROUTE:** IV, IO, IM, IN.

**DOSAGE:**

**ADULTS:** Pain: 0.2 mg/kg (max 25 mg) IM/IN/slow IV/IO push over 60 seconds or infusion over 10 minutes, may repeat every 10 minutes as needed, maximum cumulative dose 100 mg.

Extreme Agitation: 4 mg/kg IM (max 400 mg).

**ADULTS/PEDS:** Post-intubation sedation: 2 mg/kg IV (max 100 mg), may repeat every 10 minutes.

**PREGNANCY SAFETY:** Not classified.

**COMMENTS:** Sub-dissociative, low-dose ketamine used to treat pain. May be used in hypotensive patients although state protocols require SBP > 100 mmHg. Accidental overdose not associated with long-term morbidity. EtCO2 monitoring required if patient also receives narcotic pain medication. May treat adverse effects of agitation, aggression, and negative psychological reactions with Midazolam. Treat laryngospasm with ventilation with BVM, airway adjuncts, and suctioning as necessary. May treat hypersalivation with Atropine 0.5 mg IV.

### **KETOROLAC (TORADOL)**

**PROVIDER LEVEL:** AEMT, Paramedic.

**CLASS:** Nonsteroidal anti-inflammatory (NSAID).

**PHARMACOLOGY/ACTIONS:** Works by reducing hormones that cause inflammation and pain in the body. Possesses no sedative or anxiolytic properties. Also inhibits platelets.

**ONSET/DURATION:** Onset: 30 minutes / Duration: 4-6 hours.

**INDICATIONS:** Pain control of mild to moderate pain (age < 65 years old).

**CONTRAINDICATIONS:** NSAID allergy, ASA-sensitive asthma, active hemorrhage, hypotension, history of renal disease or kidney transplant, blood clotting disorder, multisystem trauma, closed head injury or bleeding in brain, history of GI bleed or ulcers, patient potentially needing surgery, open fracture or fracture deformity, and pregnancy.

**SIDE EFFECTS:** Acute kidney injury, stomach upset, nausea, vomiting, dizziness, blurred vision, dry mouth, and irritation at the injection site.

### **KETOROLAC (TORADOL) (Cont.)**

**DRUG INTERACTIONS:** Other NSAIDs, steroids, aspirin, and anticoagulants (ex Coumadin, Xarelto, Eliquis).

**ROUTE:** IV, IO, IM.

**DOSAGE:**

**ADULTS:** 15 mg IV/IO/IM once.

**PEDS:** age > 2 yo 0.5 mg/kg (max 15 mg) IV/IO.

**PREGNANCY SAFETY:** Category C – give only if potential benefits justifies risk to fetus. Avoid in pregnancy after 20 weeks gestation.

**COMMENTS:** Use with caution in patients who are dehydrated or taking ACEIs or ARBs. Ensure patient has not taken oral NSAID in last 6 hours (ibuprofen, Motrin), 12 hours (naproxen), or 24 hours (meloxicam) given potential for cumulative risk of inducing serious NSAID-related side effects.

### **LIDOCAINE (XYLOCAINE)**

**PROVIDER LEVEL:** Paramedic.

**CLASS:** Antidysrhythmic (Class IB), local anesthetic.

**PHARMACOLOGY/ACTIONS:** Lidocaine exerts its antidysrhythmic action by increasing the fibrillation threshold of the myocardium and decreasing automaticity, thus suppressing ventricular ectopy. Additional properties include decreasing stimulation of the CNS during intubation and providing local anesthetic effects in small doses.

**ONSET/DURATION:** Onset: 30-90 seconds / Duration: 10-20 minutes.

**INDICATIONS:** - Ventricular dysrhythmias / irritability.

- Following successful defibrillation or cardioversion.

- Anesthesia for initial IO line flush.

**CONTRAINDICATIONS:** Hypersensitivity to the drug, AV block, ventricular escape beats, accelerated idioventricular rhythm, and bradycardia.

**SIDE EFFECTS:** Hypotension, bradycardia, shock, CNS depression, confusion, dizziness, seizures, and slurred speech.

**DRUG INTERACTIONS:** Phenytoin increases its cardiac depressant effects. Procainamide compounds neurologic and cardiac effects. Cimetidine, beta-blockers, and quinidine increase the pharmacologic effects. Barbiturates decrease its activity.

**ROUTE:** IV, IO.

**DOSAGE:**

**ADULTS:** Ventricular dysrhythmias: 1-1.5 mg/kg IV/IO push, repeat 0.5-0.75 mg /kg IV/IO push, maximum 3 mg/kg.

Infusion: 1-4 mg/min IV/IO gtts.

IO anesthesia: 0.5 mg/kg (0.1 mg/ml of 1% preservative-free) (max 40 mg) slow push through IO.

**PEDS:** Ventricular dysrhythmias: 1 mg/kg IV/IO push, maximum 3 mg/kg.

IO anesthesia: 0.5 mg/kg (0.1 mg/ml of 1% preservative-free) (max 40 mg) slow push through IO.

**PREGNANCY SAFETY:** Category B – unproven or unknown risk to fetus, and no risk in later trimesters.

**COMMENTS:** Lidocaine is an alternative agent in ACLS algorithm for treatment of V-fib/pulseless V-tach. Reduce to half dose in the presence of decreased cardiac output, shock, patients older than 70 years and those with hepatic dysfunction.

### **LORAZEPAM (ATIVAN)**

**PROVIDER LEVEL:** Paramedic.

**CLASS:** Short-acting benzodiazepine.

**PHARMACOLOGY/ACTIONS:** Lorazepam is a water-soluble short-acting benzodiazepine with anticonvulsant, anxiolytic, sedative, and hypnotic effects. It acts on the limbic, thalamic, and hypothalamic regions of the CNS to potentiate the effects of inhibitory (GABA) neurotransmitters. Like diazepam, it suppresses the spread of seizure activity through the motor cortex of the brain while not abolishing the abnormal discharge focus.

**ONSET/DURATION:** Onset: 1-5 minutes IV, 15-30 minutes IM / Duration: 6-8 hours.

**INDICATIONS:** - Seizure activity.

- Acute anxiety states.

- Premedication before cardioversion or TCP.

- Chemical restraint in combative patient.

- Skeletal muscle relaxation.

**CONTRAINDICATIONS:** Hypersensitivity and pregnancy. Relative in respiratory depression, glaucoma, and psychosis.

**SIDE EFFECTS:** Hypotension, respiratory depression, apnea, and anterograde amnesia.

### **LORAZEPAM (ATIVAN) (Cont.)**

**DRUG INTERACTIONS:** Alcohol, CNS depressants, and anticonvulsants may precipitate CNS depression. Cimetidine increases plasma levels and potential toxicity.

**ROUTE:** IV, IO, IM.

**DOSAGE:**

**ADULTS:** Seizure: 0.1 mg/kg IV (max 4 mg), may repeat x 1 after 5 min.

Severe agitation: 2 mg IV or 4 mg IM, may repeat x 1 after 5 min.

Adjunct pain: 2 mg IV/IM.

Cardioversion or TCP: 0.1 mg/kg IM/IV/IO, max dose 2 mg.

**PEDS:** Seizure: 0.1 mg/kg IV (max 4 mg), repeat x 1 after 5 min.

**PREGNANCY SAFETY:** Category D – positive risk to fetus, maternal benefit may outweigh risk to fetus.

**COMMENTS:** DEA schedule IV drug with potential for abuse. Because lorazepam is a relatively short-acting drug, seizure activity may recur requiring additional dosing. Resuscitation equipment should be readily available. Shelf life of unrefrigerated lorazepam is 60 days, expiration dates and medication rotation must be monitored.

### **MAGNESIUM SULFATE**

**PROVIDER LEVEL:** Paramedic.

**CLASS:** Electrolyte.

**PHARMACOLOGY/ACTIONS:** Magnesium acts as a physiologic calcium channel blocker and blocks neuromuscular transmission, thereby providing electrical stability in the myocardium. Affects impulse formation and conduction time in the myocardium, thus reduces incidence of dysrhythmias associated with hypomagnesemia or prolonged QT interval. Also has anticonvulsant properties thought to be produced by CNS depression.

**ONSET/DURATION:** Onset: immediate / Duration: 30 minutes.

**INDICATIONS:**

- Torsades de pointes (polymorphic V-tach).
- Pre-eclampsia/eclampsia.
- Reactive airway disease/asthma refractory to other treatments.

**CONTRAINDICATIONS:** Myocardial damage, heart block, and renal disease.

**SIDE EFFECTS:** Respiratory depression, CNS depression, hypotension, pulmonary edema, flushing and swelling.

**DRUG INTERACTIONS:** Neuromuscular blocking agents and CNS depressants add to respiratory depression and apnea.

**ROUTE:** IV, IO.

**DOSAGE:**

**ADULTS:** Torsades de pointes: 2 g IV/IO (Stable: in 50 ml NS over 10 min; Unstable: push over 1-2 min).

Pre-eclampsia: 4 g in 50 ml NS over 20-30 min IV/IO. Eclampsia: 6 g in 50 ml NS over 5-10 min IV/IO.

Respiratory distress: 2 g in 50 ml NS over 10 min.

**PEDS:** Torsades de pointes: 25 mg/kg IV/IO in 50 ml NS over 10 min.

**PREGNANCY SAFETY:** Category A – considered safe during pregnancy.

**COMMENTS:** Administer magnesium sulfate slowly at no more than 1 gram/minute, no matter what the clinical condition. Calcium chloride or gluconate can be used as an antidote if serious side effects occur. Early indicators of magnesium toxicity include cathartic effect, profound thirst, feeling of warmth, sedation, confusion, depressed reflexes, and muscle weakness.

### **METHYLPREDNISOLONE (SOLU-MEDROL)**

**PROVIDER LEVEL:** EMT w/medication endorsement and AEMT (IM only), Paramedic.

**CLASS:** Glucocorticoid.

**PHARMACOLOGY/ACTIONS:** Methylprednisolone is a synthetic steroid that suppresses acute and chronic inflammation.

**ONSET/DURATION:** Onset: 1-2 hours / Duration: 8-24 hours.

**INDICATIONS:**

- Anaphylaxis/severe allergic reaction.
- Severe asthma exacerbation.
- COPD exacerbation.
- Patient with adrenal insufficiency in medical distress at risk of acute adrenal crisis.

**CONTRAINDICATIONS:** Use with caution in patients with gastrointestinal bleeding, diabetes mellitus, or severe infection.

**SIDE EFFECTS:** Headache, hypertension, sodium and water retention, hypokalemia, and alkalosis.

**DRUG INTERACTIONS:** Hypoglycemic responses to insulin and oral hypoglycemic agents may be blunted. Potassium-depleting agents may potentiate hypokalemia induced by corticosteroids.

**ROUTE:** IV, IO, IM.



**METHYLPREDNISOLONE (SOLU-MEDROL)****DOSAGE:**

**ADULTS:** 125 mg slow IV/IO/IM.

**PEDS:** 1-2 mg/kg (max 125 mg) slow IV/IO/IM.

**PREGNANCY SAFETY:** Category C – give only if potential benefits justifies risk to fetus.

**COMMENTS:** Methylprednisolone should not be considered first line therapy as onset of action takes 1-2 hours.

**METOCLOPRAMIDE (REGLAN)**

**PROVIDER LEVEL:** AEMT and Paramedic.

**CLASS:** Antiemetic.

**PHARMACOLOGY/ACTIONS:** Metoclopramide is a potent central and peripheral dopamine receptor antagonist. The exact mechanism of action is not clear, but it appears to sensitize GI smooth muscle to effects of acetylcholine by direct action. It increases the resting tone of the esophageal sphincter and the tone and amplitude of upper GI contractions. As a result, gastric emptying and intestinal transit are accelerated with little effect if any on gastric, biliary (ie gallbladder), or pancreatic secretions (eg promotility effect). Antiemetic action results from drug-induced elevation of CTZ threshold and enhanced gastric emptying.

**ONSET/DURATION:** Onset 1-3 minutes IV versus 10-15 minutes IM / Duration: 1-3 hours.

**INDICATIONS:** Nausea and vomiting.

**CONTRAINDICATIONS:** Prior sensitivity or intolerance to the drug, allergy to sulfite agents, and suspected small bowel obstruction or perforation. Use with caution in patients with concurrent use of drugs that can cause extrapyramidal symptoms.

**SIDE EFFECTS:** Mild sedation, dizziness, restlessness, agitation, extrapyramidal symptoms (acute dystonic type), blurry or double vision, and dry mouth.

**DRUG INTERACTIONS:** Alcohol and other CNS depressants may add to sedation. Phenothiazines (antipsychotic medications) may potentiate extrapyramidal symptoms. Hypertension may occur when metoclopramide is administered to patients taking MAO inhibitors.

**ROUTE:** IM, IV.

**DOSAGE:**

**ADULTS:** 5-10 mg slow IV/IM.

**PEDS:** if 8 years of age or older 5 mg slow IV/IM.

**PREGNANCY SAFETY:** Category B – unproven or unknown risk to fetus, and no risk in later trimesters. Generally considered safe in pregnancy.

**COMMENTS:** The injection form contains sodium metabisulfite as an antioxidant. If patient has history of allergy to sulfite agents, this product should be avoided. Extrapyramidal symptoms are most likely to occur in young adults, elderly, and with high-dose treatment of vomiting associated with cancer chemotherapy. Development of extrapyramidal symptoms can be reduced or eliminated by diluting the medication and slow administration over several minutes. Diphenhydramine (Benadryl) should be available as an antidote.

**METOPROLOL (LOPRESSOR)**

**PROVIDER LEVEL:** Paramedic.

**CLASS:** Antihypertensive, Beta Blocker, Beta 1 Selective.

**PHARMACOLOGY/ACTIONS:** Selective inhibitor of beta1-adrenergic receptors; competitively blocks beta1-receptors. Decreases cardiac output and heart rate.

**ONSET/DURATION:** Onset: immediate (peak effect 20 minutes) / Duration: 6 hours.

**INDICATIONS:** Atrial fibrillation/flutter with rapid ventricular response.

**CONTRAINDICATIONS:** Hypersensitivity, cardiogenic shock, hypotension, bradycardia, 2nd or 3rd degree heart block (unless functioning pacemaker), asthma, and COPD.

**SIDE EFFECTS:** Hypotension, bradycardia, bronchospasm, fatigue, and hypoglycemia masking.

**DRUG INTERACTIONS:** Dexmedetomidine and antihypertensives (other beta-blockers, calcium channel blockers).

**ROUTE:** IV.

**DOSAGE:**

**ADULTS:** 5 mg slow IVP over 2 min, may repeat every 5 minutes as needed if SBP>100, max cumulative dose 15 mg.

**PREGNANCY SAFETY:** Not preferred, can be continued if risks of discontinuing outweigh benefits.

**COMMENTS:** Use with caution in patients with impaired hepatic or renal function. Use with caution in patients with cardiomegaly and CHF controlled by digitalis and diuretics, observe closely for impending heart failure. During IV administration: BP, HR, and ECG should be monitored carefully.

## **MIDAZOLAM (VERSED)**

**PROVIDER LEVEL:** Paramedic.

**CLASS:** Short-acting benzodiazepine.

**PHARMACOLOGY/ACTIONS:** Midazolam is a water-soluble short-acting benzodiazepine that is metabolized by the liver and excreted in the urine. It binds benzodiazepine receptors and enhances GABA central nervous system effects.

**ONSET/DURATION:** Onset: 1-3 minutes / Duration: 2-6 hours.

**INDICATIONS:**

- Seizure activity.
- Acute anxiety states.
- Premedication before cardioversion or TCP.
- Chemical restraint in combative patient.
- Skeletal muscle relaxation.

**CONTRAINDICATIONS:** Hypersensitivity, glaucoma (relative), shock, coma, alcohol intoxication (relative), depressed vital signs and concomitant use of barbiturates, alcohol, narcotics, or other CNS depressants used by the patient.

**SIDE EFFECTS:** Respiratory depression, hiccups, cough, over sedation, nausea and vomiting, headache, blurred vision, fluctuation in vital signs, hypotension, and respiratory arrest.

**DRUG INTERACTIONS:** Sedative effect of midazolam accentuated by concomitant use of barbiturates, alcohol, or narcotics.

**ROUTE:** IV, IO, IM, IN.

**DOSAGE:**

**ADULTS:** Seizure: 0.2 mg/kg IN/IM (max 10 mg) or 0.1 mg/kg IV/IO (max 5 mg), may repeat x 1 after 5 min.

Severe agitation: 5 mg IN/IM/IV, may repeat x 1 after 5 min.

Adjunct pain: 2-4 mg IV/IM.

Cardioversion or TCP: 0.1 mg/kg IN/IM/IV/IO, max dose 5 mg.

**PEDS:** Seizure: 0.2 mg/kg IN/IM (max 10 mg) or 0.1 mg/kg IV/IO (max 5 mg), may repeat x 1 after 5 min.

**PREGNANCY SAFETY:** Category D – positive risk to fetus, maternal benefit may outweigh risk to fetus.

**COMMENTS:** Note: in elderly or patients with known liver disease, lower dosage should be used due to impaired metabolism. DEA schedule IV drug with potential for abuse. Provide continuous monitoring of respiratory and cardiac function. Have resuscitation equipment available. Never administer as rapid push IV bolus, which may lead to profound hypotension and/or respiratory impairment.

## **MORPHINE**

**PROVIDER LEVEL:** Paramedic.

**CLASS:** Narcotic analgesic.

**PHARMACOLOGY/ACTIONS:** Extremely potent narcotic analgesic. It dilates peripheral vasculature (reducing pre-load and after-load and decreasing myocardial oxygen demand). Morphine also tends to reduce the respiratory rate and tidal volume and causes pupils to constrict. It reduces apprehension and anxiety. The vasodilatation should cause no problems if patients are supine and not upright, not volume depleted, or have a decreased cardiac output. The onset of action is immediate if given IV. Peak effects are seen within 20 minutes.

**ONSET/DURATION:** Onset: 1-2 minutes / Duration: 2-7 hours.

**INDICATIONS:**

- Cardiac suspected chest pain **after** administration of oxygen, aspirin, and nitroglycerin according to guideline.
- Pain control in absence of hypotension.
- Analgesia for cardioversion or TCP.

**CONTRAINDICATIONS:** Major blood loss (hypovolemia), hypotension, head or abdominal injuries, increased ICP, and respiratory difficulties. Relative contraindication in patients who have taken alcohol, tricyclic antidepressants, MAO inhibitors, or other depressants.

**SIDE EFFECTS:** Respiratory depression, hypotension, nausea and vomiting, decreased LOC, constricted pupils, urinary retention, and histamine release.

**DRUG INTERACTIONS:** CNS depressants may potentiate effects of morphine (respiratory depression, hypotension, sedation). Phenothiazines and benzodiazepines may potentiate analgesia. MAO inhibitors may cause paradoxical excitation.

**ROUTE:** IM, IV, IO.

**DOSAGE:**

**ADULT:** Pain: 0.1 mg/kg IM/IV/IO (max 10 mg), may repeat after 10 min.

Cardioversion or TCP: 0.1 mg/kg slow IV/IO (max 10 mg)

**PEDS:** Pain: 0.1 mg/kg IM/IV/IO (max 5 mg), may repeat after 10 min.

**PREGNANCY SAFETY:** Category B – unproven or unknown risk to fetus, and no risk in later trimesters.

### **MORPHINE (Cont.)**

**COMMENTS:** DEA schedule II drug with potential for abuse. Closely monitor the patient's blood pressure before and after administration of morphine. Use with caution in the elderly, those with asthma, and in those susceptible to CNS depression. Naloxone should be readily available.

### **NALOXONE (NARCAN)**

**PROVIDER LEVEL:** EMR and EMT (IN/auto-injector), EMT w/medication endorsement, AEMT, and Paramedic (IN/IM/IV/IO).

**CLASS:** Narcotic antagonist.

**PHARMACOLOGY/ACTIONS:** Naloxone is a narcotic antagonist which competitively binds to narcotic sites, but which exhibits almost no pharmacologic activity of its own.

**ONSET/DURATION:** Onset: 2 minutes / Duration: 30-60 minutes.

**INDICATIONS:** Coma, altered mental status, ingestions, poisoning, drug overdose, head trauma with decreased mental status.

**CONTRAINDICATIONS:** Known hypersensitivity. Use with caution in narcotic-dependent patients who may experience withdrawal syndrome (including neonates of narcotic-dependent mothers).

**SIDE EFFECTS:** Tachycardia, hypertension, dysrhythmias, nausea, vomiting, diaphoresis, blurred vision, and opiate withdrawal.

**DRUG INTERACTIONS:** Incompatible with bisulfite and alkaline solutions.

**ROUTE:** IM, IV, IO, IN.

**DOSAGE:**

**ADULTS:** 2-4 mg IN, 0.4-2 mg IM/IV/IO, titrated to increase in respiratory effort, repeat as necessary.

**PEDS:** 0.1 mg/kg IN/IM/IV/IO (max 2 mg), titrated to increase in respiratory effort, repeat as necessary.

**PREGNANCY SAFETY:** Category B – unproven or unknown risk to fetus, and no risk in later trimesters.

**COMMENTS:** Consider repeated doses if transient response is noted. Naloxone may not reverse hypotension. Exercise caution when administering naloxone to narcotic addicts (may precipitate withdrawal with hypertension, tachycardia, and violent behavior).

### **NITROGLYCERIN (NITROSTAT)**

**PROVIDER LEVEL:** EMT (patient prescribed only), EMT w/medication endorsement and AEMT (SL), Paramedic (SL/TD/IV).

**CLASS:** Anti-anginal / vasodilator.

**PHARMACOLOGY/ACTIONS:** Nitrates and nitrites relax smooth muscle dilating arterioles and veins in the periphery (and coronary arteries in high doses). The resultant reduction in preload, and to a lesser extent in afterload, decreases the workload of the heart and lowers myocardial oxygen demand. Antianginal, anti-ischemic, and antihypertensive effects.

**ONSET/DURATION:** Onset: 1-3 minutes (SL/TD), immediate (IV) / Duration: 30-60 minutes (SL/TD), 3-5 minutes (IV).

**INDICATIONS:** Chest pain suggestive of acute coronary syndrome (ACS), acute pulmonary edema, congestive heart failure, hypertensive emergency with normal mental status and no focal weakness.

**CONTRAINDICATIONS:** Increased intracranial pressure, cerebral edema, severe hypotension, dehydration, children under 12, aortic stenosis, and recent use of erectile dysfunction drugs such as Viagra, Cialis, Levitra, Revatio, Staxyn, etc.

**SIDE EFFECTS:** Hypotension, headache, dizziness, tachycardia, nausea, vomiting and diaphoresis.

**DRUG INTERACTIONS:** Other vasodilators may have additive hypotensive effects.

**ROUTE:** SL, TD, IV.

**DOSAGE:**

**ADULTS:** 0.4 mg (1 tablet or spray) SL, may be repeated every 5 minutes up to three doses.

Consider NTG paste 0.5-1" to chest wall.

Maintenance: infusion 10 mcg/min IV, titrate 5 mcg/min every 3-5 min, maximum 200 mcg/min.

**PREGNANCY SAFETY:** Category C – give only if potential benefits justifies risk to fetus.

**COMMENTS:** Establish IV access. Use only if SBP > 100 mmHg. Recheck BP and pulse after administration. Therapeutic effect is enhanced but adverse effects are increased when patient is upright. Transient headache is common. If become hypotensive, stop drip, Trendelenburg, and consider fluid bolus. Use with caution or withhold if concern for RV ischemia/infarction (ECG changes in leads II, III, aVF, and V4R) and be prepared to augment preload if BP precipitously drops.

### **NOREPINEPHRINE (LEVOPHED)**

**PROVIDER LEVEL:** Paramedic.

**CLASS:** Sympathomimetic.

**NOREPINEPHRINE (LEVOPHED) (Cont.)**

**PHARMACOLOGY/ACTIONS:** Norepinephrine is a direct-acting sympathomimetic amine identical to the body catecholamine norepinephrine. It acts primarily on  $\alpha$ -adrenergic receptors, with little action on  $\beta$ -adrenergic receptors except in the heart (beta1 receptors). Main therapeutic effects are vasoconstriction and cardiac stimulation. Powerful vasoconstrictor action on resistance and capacitance blood vessels. Peripheral vasoconstriction and moderate inotropic stimulation of heart result in increased systolic and diastolic blood pressure, myocardial oxygenation, coronary artery blood flow, and work of heart.

**ONSET/DURATION:** Onset: 1-2 minutes / Duration: 1-2 minutes after termination of infusion.

**INDICATIONS:** - Hemodynamically significant hypotension in the absence of hypovolemia.  
- Shock.

**CONTRAINDICATIONS:** Hypovolemia. Use with caution in patients taking MAOIs, antihistamines, and antidepressants (imipramine or imipramine types) due to risk of prolonged hypertension.

**SIDE EFFECTS:** Headache, hypertension, reflexive bradycardia, anxiety, dyspnea, ischemic injury due to vasoconstriction, dysrhythmias, and tissue necrosis at injection site (with extravasation).

**DRUG INTERACTIONS:** Alpha and beta blockers antagonize pressor effects. TCAs may potentiate pressor effects.

**ROUTE:** IV, IO.

**DOSAGE:**

**ADULTS:** Shock/Sepsis: 0.05-0.5 mcg/kg/min IV/IO drip titrated to maintain SBP >90 or MAP >65.

Pulmonary edema/CHF: 0.02-2 mcg/kg/min IV/IO drip titrated to maintain SBP >90 or MAP >65.

**PEDS:** 0.05-0.5 mcg/kg/min IV/IO drip titrated to maintain SBP >70 mmHg + 2 x age.

**PREGNANCY SAFETY:** Category C – give only if potential benefits justifies risk to fetus.

**COMMENTS:** Utilize a large vein to decrease risk of extravasation with can cause tissue necrosis. Continually check IV site for patency and signs/symptoms of infiltration. Mix in D5W to protect against loss of potency due to oxidation. Do not use solution if discoloration or precipitate is present. Protect from light. Early consultation with Medical Control to discuss medical decision making and treatment is encouraged when possible.

**ONDANSETRON (ZOFTRAN)**

**PROVIDER LEVEL:** AEMT, Paramedic.

**CLASS:** Antiemetic.

**PHARMACOLOGY/ACTIONS:** Ondansetron's mechanism of action has not been fully characterized. The released serotonin may stimulate the vagal afferents through the 5-HT<sub>3</sub> receptors and initiate the vomiting reflex. Ondansetron selectively antagonizes the 5-HT<sub>3</sub> receptors. It has limited effectiveness for motion sickness, consider diphenhydramine (Benadryl) for refractive nausea/vomiting in those settings.

**ONSET/DURATION:** Onset: 30 minutes for peak effect / Duration: 5-7 hours.

**INDICATIONS:** Prevention and treatment of nausea and vomiting in adults and pediatrics.

**CONTRAINDICATIONS:** Hypersensitivity to drug/class (Kytril and Aloxi), gastric/abdominal surgery in pediatric patients.

**SIDE EFFECTS:** Headache, dizziness, diarrhea, rash, agitation, and prolonged QT interval.

**DRUG INTERACTIONS:** Apomorphine and Dronedarone.

**ROUTE:** IV, IO, PO (oral disintegrating tablet/ODT).

**DOSAGE:**

**ADULTS:** 4 mg ODT/IV/IO.

**PEDS:** 0.15 mg/kg ODT/IV/IO (max dose 4 mg), round up to nearest ½ pill.

**PREGNANCY SAFETY:** Category B – unproven or unknown risk to fetus, and no risk in later trimesters. Generally considered safe in pregnancy.

**COMMENTS:** Consider early in patients with spinal motion restriction to decrease risk of vomiting and aspiration. Use caution in patients with severe liver disease, the dose should not exceed 8 mg in 24 hours. Not commonly used in patients < 1 year of age.

**ORAL GLUCOSE (GLUTOSE)**

**PROVIDER LEVEL:** EMT, AEMT, Paramedic.

**CLASS:** Oral hypoglycemic agent, carbohydrate (sugar).

**PHARMACOLOGY/ACTIONS:** Elevates blood glucose. Causes hyperosmolar diuresis and decreases cerebral edema.

**ONSET/DURATION:** Onset: 10 minutes / Duration: varies.

**INDICATIONS:** Hypoglycemia, seizures, and/or altered mental status and unable to determine blood sugar level.

**CONTRAINDICATIONS:** Unconscious, unable to protect airway.

**ORAL GLUCOSE (GLUTOSE) (Cont.)**

**SIDE EFFECTS:** May be aspirated if patient is unable to protect airway.

**DRUG INTERACTIONS:** None.

**HOW SUPPLIED:** Gel in 15 gm/tube (single use).

**ROUTE:** Oral.

**DOSAGE:**

**ADULTS:** 10-20 grams, may be repeated in 10 minutes if necessary.

**PEDS:** 5-20 grams, may be repeated in 10 minutes if necessary.

**PREGNANCY SAFETY:** Category C – give only if potential benefits justifies risk to fetus.

**COMMENTS:** Because changes in level of consciousness can occur rapidly in patients with hypoglycemia, it is important to ascertain the patient's ability to swallow an oral preparation without airway compromise.

**OXYGEN**

**PROVIDER LEVEL:** EMR, EMT, AEMT, Paramedic.

**CLASS:** Gas.

**PHARMACOLOGY/ACTIONS:** Helps to oxidize glucose to produce ATP.

**ONSET/DURATION:** Onset: immediate / Duration: < 2 minutes.

**INDICATIONS:** May be used in any trauma and/or medical patient – specifically hypoxia, ischemic chest pain, dyspnea, CO poisoning, and cardiac arrest.

**CONTRAINDICATIONS:** Should never be withheld in any critically ill patient.

**SIDE EFFECTS:** High-concentrations of oxygen may cause decreased level of consciousness and respiratory depression in patients with chronic carbon dioxide retention (ie COPD).

**DRUG INTERACTIONS:** None.

**ROUTE:** Inhaled.

**DOSAGE:** Low: 1 – 6 lpm via NC. High: 10 – 15 lpm via NRB or BVM.

**PREGNANCY SAFETY:** N/A.

**COMMENTS:** Oxygen vigorously supports combustion. The Gallatin County EMS Patient Care Guidelines list goal oxygen saturation >94% to be consistent with the NASEMSO National Model EMS Clinical Guidelines. However, given the altitude throughout Gallatin County, maintaining an oxygen saturation >90% is an acceptable goal. In the setting of COPD, the goal oxygen saturation should be 88-90%.

**OXYMETAZOLINE (AFRIN)**

**PROVIDER LEVEL:** Paramedic.

**CLASS:** Decongestant.

**PHARMACOLOGY/ACTIONS:** Alpha-adrenergic agonist. Stimulates alpha-adrenergic receptors and produces vasoconstriction in the arterioles of the nasal mucosa.

**ONSET/DURATION:** Onset: within 10 minutes / Duration: up to 12 hours.

**INDICATIONS:** Management of epistaxis.

**CONTRAINDICATIONS:** Hypersensitivity.

**SIDE EFFECTS:** Temporary local discomfort such as burning, stinging, or sneezing.

**DRUG INTERACTIONS:** Fentanyl, esketamine, and zavegepant.

**ROUTE:** Topical.

**DOSAGE:**

**ADULTS:** 2 sprays to each nostril.

**PREGNANCY SAFETY:** Adverse fetal/neonatal events have been noted in case reports following large doses or extended use of oxymetazoline nasal spray in the first trimester of pregnancy.

**COMMENTS:** For intranasal use only. Shake well before use. Before using for the first time, prime the pump by firmly depressing the rim several times. Keep head upright and insert nozzle into nostril, depress rim firmly, and inhale deeply. Use with caution in patients with hypertension, heart disease, or diabetes mellitus.

**OXYTOCIN (PITOCIN)**

**PROVIDER LEVEL:** Paramedic.

**CLASS:** Hormone.

### **OXYTOCIN (PITOCIN) (Cont.)**

**PHARMACOLOGY/ACTIONS:** The medication is pharmacologically identical to naturally occurring oxytocin that is secreted by the posterior pituitary. Uterine sensitivity to oxytocin increases during pregnancy and peaks sharply before delivery. It produces phasic contractions characteristic of normal delivery.

**ONSET/DURATION:** Onset: immediately IV versus 3-5 minutes IM / Duration: 1 hour IV versus 2-3 hours IM.

**INDICATIONS:** Heavy postpartum hemorrhage following delivery of the placenta.

**CONTRAINDICATIONS:** Prior to delivery of fetus(es) and hypersensitivity to medication. When administered prior to delivery, may cause fetal hypoxia, fetal asphyxia, fetal dysrhythmias, and possible fetal intracranial bleeding.

**SIDE EFFECTS:** Tachycardia, cardiac dysrhythmias, hypertensive episodes, seizures, nausea, and vomiting. Hypersensitivity leads to uterine hypertonicity, tetanic contractions, uterine rupture, and/or anaphylactic reaction.

**DRUG INTERACTIONS:** Vasoconstrictors can cause severe hypertension.

**ROUTE:** IV, IM.

**DOSAGE:**

**ADULTS:** 20 units IV in 1000 ml NS or LR infused wide open.

**PREGNANCY SAFETY:** Category C – give only if potential benefits justifies risk to fetus.

**COMMENTS:** Ensure fetus and placenta have delivered and there is not an additional fetus in the uterus. Administer properly diluted IV solution by continuous infusion only. Fundus should be checked frequently. Incidence of hypersensitivity or allergic reactions is higher when given IM or IV injection rather than by IV infusion of diluted solution.

### **SODIUM BICARBONATE**

**PROVIDER LEVEL:** Paramedic.

**CLASS:** Buffer, alkalizing agent, and electrolyte supplement.

**PHARMACOLOGY/ACTIONS:** Sodium bicarbonate reacts with hydrogen ions to form water and carbon dioxide and thereby can act to buffer metabolic acidosis. As the plasma hydrogen ion concentration decreases, blood pH rises.

**ONSET/DURATION:** Onset: 2-10 minutes / Duration: 30-60 minutes.

**INDICATIONS:**

- Tricyclic antidepressant (TCA)/sodium channel blocker overdose with wide QRS.
- Known or suspected hyperkalemia.
- Alkalization for treatment of specific toxidromes/rhabdomyolysis (with medical control consultation).

**CONTRAINDICATIONS:** None if patient in extremis. Metabolic and respiratory alkalosis. Routine use in cardiac arrest.

**SIDE EFFECTS:** Metabolic alkalosis, hypoxia, rise in intracellular PCO<sub>2</sub> and increase in tissue acidosis, electrolyte imbalance (hypernatremia).

**DRUG INTERACTIONS:** Alkalization of urine may shorten elimination half-lives of certain drugs. Vasopressors may be deactivated.

**ROUTE:** IV, IO.

**DOSAGE:**

**ADULTS:** Suspected hyperkalemia: 1 mEq/kg IV/IO of 8.4% adult preparation.

TCA OD: 1-2 mEq/kg IV/IO of 8.4% adult preparation, may repeat x1 in 3-5 min if QRS remains wide.

**PEDS:** 1 mEq/kg IV/IO (max dose 50 mEq) of 7.5% pediatric preparation, infuse slowly and only if ventilations are adequate.

**PREGNANCY SAFETY:** Category C – give only if potential benefits justifies risk to fetus.

**COMMENTS:** Bicarb administration produces carbon dioxide, which crosses cell membranes more rapidly than bicarbonate (potentially worsening intracellular acidosis). Bicarb may worsen CHF. Maintain adequate ventilation (gas exchange) to correct most underlying metabolic/respiratory acidosis states. May use 8.4% adult preparation in pediatrics if 7.5% pediatric concentration unavailable; need to dilute 8.4% concentration 1:1 with normal saline to create 4.2% solution for age < 1 year old.

### **TRANEXAMIC ACID (TXA)**

**PROVIDER LEVEL:** Paramedic.

**CLASS:** Antifibrinolytic.

**PHARMACOLOGY/ACTIONS:** Potent antifibrinolytic drug. The main action is blocking the lysine-binding sites of the plasminogen molecule. This prevents activation of plasminogen by plasminogen activator. There is no evidence of a thrombogenic effect.

**ONSET/DURATION:** Onset: several minutes / Duration: 3 hours.

**TRANEXAMIC ACID (TXA) (Cont.)**

**INDICATIONS:** - Trauma with suspected hemorrhagic shock (HR > 110 or SBP < 90).  
- Postpartum hemorrhage.

**CONTRAINDICATIONS:** Time since injury > 3 hours, isolated spinal shock (cord injury without evidence of hemorrhage), known thromboembolic disease, and mechanical valve.

**SIDE EFFECTS:** Anaphylaxis, anaphylactoid reaction, seizure, headache, dizziness, and ocular effects.

**DRUG INTERACTIONS:** Anti-inhibitor coagulant complex, estrogen derivatives, factor IX complex, prothrombin complex concentrate, thrombolytic agents, and Tretinoin.

**ROUTE:** IV/IO.

**DOSAGE:**

**ADULTS:** 2 g IV/IO infusion over 10 minutes, preferable before IV fluids.

**PREGNANCY SAFETY:** Category B – adequate studies in pregnant women have not demonstrated a risk to the fetus in the 1<sup>st</sup> trimester and there is no risk in the last trimesters.

**COMMENTS:** Randomized placebo-controlled trial showed early administration of TXA in patients with significant hemorrhage reduced the risk of death from bleeding. Post-hoc analysis revealed treatment beyond 3 hours was shown to be significantly less effective and possibly associated with harm. In patients with subarachnoid hemorrhage (SAH), may cause cerebral edema and infarction. The 2 gram bolus dose has a slightly increased risk of seizures, but did not increase adverse outcomes. There is an increased risk of venous thromboembolism (VTE) in doses above 30 mg/kg.